

Basic Riemannian Geometry and Sobolev Estimates used in Symplectic Topology

Aleksey Zinger*

September 26, 2021

Abstract

This note collects a number of standard statements in Riemannian geometry and in Sobolev-space theory that play a prominent role in analytic approaches to symplectic topology. These include relations between connections and complex structures, estimates on exponential-like maps, and dependence of constants in Sobolev and elliptic estimates.

Contents

1	Connections in real vector bundles	2
1.1	Connections and splittings	2
1.2	Metric-compatible connections	5
1.3	Torsion-free connections	5
2	Complex structures	6
2.1	Complex linear connections	6
2.2	Generalized $\bar{\partial}$ -operators	8
2.3	Connections and $\bar{\partial}$ -operators	10
2.4	Holomorphic vector bundles	11
2.5	Deformations of almost complex submanifolds	12
3	Riemannian geometry estimates	15
3.1	Parallel transport	15
3.2	Poincare lemmas	19
3.3	Exponential-like maps and differentiation	21
3.4	Expansion of the $\bar{\partial}$ -operator	23
4	Sobolev and elliptic inequalities	24
4.1	Euclidian case	25
4.2	Bundle sections along smooth maps	29
4.3	Elliptic estimates	31

*Partially supported by DMS grant 0846978

1 Connections in real vector bundles

1.1 Connections and splittings

Suppose M is a smooth manifold and $\pi_E: E \rightarrow M$ is a vector bundle. We identify M with the zero section of E . Denote by

$$\mathbf{a}: E \oplus E \rightarrow E \quad \text{and} \quad \pi_{E \oplus E}: E \oplus E \rightarrow M$$

the associated addition map and the induced projection map, respectively. For $f \in C^\infty(M; \mathbb{R})$, define

$$m_f: E \rightarrow E \quad \text{by} \quad m_f(v) = f(\pi_E(v)) \cdot v \quad \forall v \in E. \quad (1.1)$$

In particular,

$$\pi_{E \oplus E} = \pi_E \circ \mathbf{a}, \quad \pi_E = \pi_E \circ m_f \quad \forall f \in C^\infty(M; \mathbb{R}).$$

The total spaces of the vector bundles

$$\pi_{E \oplus E}: E \oplus E \rightarrow M \quad \text{and} \quad \pi_E^* E \rightarrow E$$

consist of the pairs (v, w) in $E \times E$ such that $\pi_E(v) = \pi_E(w)$.

Define a smooth bundle homomorphism

$$\iota_E: \pi_E^* E \rightarrow TE, \quad \iota_E(v, w) = \left. \frac{d}{dt}(v + tw) \right|_{t=0}. \quad (1.2)$$

Since the restriction of ι_E to the fiber over $v \in E$ is the composition of the isomorphism

$$E_{\pi_E(v)} \rightarrow T_v E_{\pi_E(v)}, \quad w \rightarrow \left. \frac{d}{dt}(v + tw) \right|_{t=0},$$

with the differential of the embedding of the fiber $E_{\pi_E(v)}$ into E , ι_E is an injective bundle homomorphism. Furthermore,

$$d\pi_E \circ \iota_E = 0, \quad m_f^* \iota_E \circ \pi_E^* m_f = dm_f \circ \iota_E, \quad \mathbf{a}^* \iota_E \circ \pi_{E \oplus E}^* \mathbf{a} = d\mathbf{a} \circ \iota_{E \oplus E}, \quad (1.3)$$

$$TE|_M \approx TM \oplus \text{Im } \iota_E. \quad (1.4)$$

By the first statement in (1.3), the injectivity of ι_E , and surjectivity of $d\pi_E$,

$$0 \longrightarrow \pi_E^* E \xrightarrow{\iota_E} TE \xrightarrow{d\pi_E} \pi_E^* TM \longrightarrow 0 \quad (1.5)$$

is an exact sequence of vector bundles over E . By the second statement in (1.3), the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_E^* E & \xrightarrow{\iota_E} & TE & \xrightarrow{d\pi_E} & \pi_E^* TM \longrightarrow 0 \\ & & \downarrow \pi_E^* m_f & & \downarrow dm_f & & \downarrow \pi_E^* \text{id} \\ 0 & \longrightarrow & \pi_E^* E & \xrightarrow{m_f^* \iota_E} & m_f^* TE & \xrightarrow{m_f^* d\pi_E} & \pi_E^* TM \longrightarrow 0 \end{array} \quad (1.6)$$

of vector bundle homomorphisms over E commutes. By the third statement in (1.3), the diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \pi_{E \oplus E}^*(E \oplus E) & \xrightarrow{\iota_{E \oplus E}} & T(E \oplus E) & \xrightarrow{d\pi_{E \oplus E}} & \pi_{E \oplus E}^*TM \longrightarrow 0 \\
& & \downarrow \pi_{E \oplus E}^* \alpha & & \downarrow d\alpha & & \downarrow \pi_{E \oplus E}^* \text{id} \\
0 & \longrightarrow & \pi_{E \oplus E}^*E & \xrightarrow{\alpha^* \iota_E} & \alpha^*TE & \xrightarrow{\alpha^* d\pi_E} & \pi_{E \oplus E}^*TM \longrightarrow 0
\end{array} \tag{1.7}$$

of vector bundle homomorphisms over $E \oplus E$ commutes.

A connection in E is an \mathbb{R} -linear map

$$\begin{aligned}
\nabla: \Gamma(M; E) &\longrightarrow \Gamma(M; T^*M \otimes_{\mathbb{R}} E) \quad \text{s.t.} \\
\nabla(f\xi) &= df \otimes \xi + f\nabla\xi \quad \forall f \in C^\infty(M), \xi \in \Gamma(M; E).
\end{aligned} \tag{1.8}$$

The Leibnitz property (1.8) implies that any two connections in E differ by a 1-form on M . In other words, if ∇ and $\tilde{\nabla}$ are connections in E there exists

$$\begin{aligned}
\theta &\in \Gamma(M; T^*M \otimes_{\mathbb{R}} \text{Hom}_{\mathbb{R}}(E, E)) \quad \text{s.t.} \\
\tilde{\nabla}_v \xi &= \nabla_v \xi + \{\theta(v)\}\xi \quad \forall \xi \in \Gamma(M; E), v \in T_x M, x \in M.
\end{aligned} \tag{1.9}$$

If U is a neighborhood of $x \in M$ and f is a smooth function on M supported in U such that $f(x)=1$, then

$$\nabla \xi|_x = \nabla(f\xi)|_x - d_x f \otimes \xi(x) \tag{1.10}$$

by (1.8). The right-hand side of (1.10) depends only on $\xi|_U$. Thus, a connection ∇ in E is a local operator, i.e. the value of $\nabla \xi$ at a point $x \in M$ depends only on the restriction of ξ to any neighborhood U of x .

Suppose U is an open subset of M and $\xi_1, \dots, \xi_n \in \Gamma(U; E)$ is a frame for E on U , i.e.

$$\xi_1(x), \dots, \xi_n(x) \in E_x$$

is a basis for E_x for all $x \in U$. By definition of ∇ , there exist

$$\theta_l^k \in \Gamma(U; T^*U) \quad \text{s.t.} \quad \nabla \xi_l = \sum_{k=1}^{k=n} \xi_k \theta_l^k \equiv \sum_{k=1}^{k=n} \theta_l^k \otimes \xi_k \quad \forall l=1, \dots, n.$$

We call

$$\theta \equiv (\theta_l^k)_{k,l=1, \dots, n} \in \Gamma(U; T^*U \otimes_{\mathbb{R}} \text{Mat}_n \mathbb{R})$$

the connection 1-form of ∇ with respect to the frame $(\xi_k)_k$.

For an arbitrary section

$$\xi = \sum_{l=1}^{l=n} f^l \xi_l \in \Gamma(U; E),$$

by (1.8) we have

$$\nabla \xi = \sum_{k=1}^{k=n} \xi_k \left(df^k + \sum_{l=1}^{l=n} \theta_l^k f^l \right), \quad \text{i.e.} \quad \nabla(\underline{\xi} \cdot \underline{f}^t) = \underline{\xi} \cdot \{d + \theta\} \underline{f}^t, \quad (1.11)$$

$$\text{where} \quad \underline{\xi} = (\xi_1, \dots, \xi_n), \quad \underline{f} = (f^1, \dots, f^n). \quad (1.12)$$

This implies that

$$\nabla \xi|_x = \pi_2|_x \circ d_x \xi: T_x M \longrightarrow E_x \quad \forall \xi \in \Gamma(U; E) \text{ s.t. } \xi(x) = 0, \quad (1.13)$$

where $\pi_2|_x: T_x E \longrightarrow E_x$ is the projection to the second component in (1.4).

By (1.11), ∇ is a first-order differential operator. By (1.8), its symbol is given by

$$\sigma_\nabla: T^* M \longrightarrow \text{Hom}(E, T^* M \otimes_{\mathbb{R}} E), \quad \{\sigma_\nabla(\eta)\}(f) = \eta \otimes f.$$

Lemma 1.1. *Suppose M is a smooth manifold and $\pi_E: E \longrightarrow M$ is a vector bundle. A connection ∇ in E induces a splitting*

$$TE \approx \pi_E^* TM \oplus \pi_E^* E \quad (1.14)$$

of the exact sequence (1.5) extending the splitting (1.4) such that

$$\nabla \xi|_x = \pi_2|_x \circ d_x \xi: T_x M \longrightarrow E_x \quad \forall \xi \in \Gamma(M; E), \quad x \in M, \quad (1.15)$$

where $\pi_2|_x: T_x E \longrightarrow E_x$ is the projection onto the second component in (1.14). Furthermore,

$$dm_t \approx \pi_E^* \text{id} \oplus \pi_E^* m_t \quad \forall t \in \mathbb{R} \quad \text{and} \quad \mathbf{a} \approx \pi_{E \oplus E}^* \text{id} \oplus \pi_{E \oplus E}^* \mathbf{a}, \quad (1.16)$$

with respect to the splitting (1.14), i.e. it is consistent with the commutative diagrams (1.6) and (1.7).

Proof. Given $x \in M$ and $v \in E_x$, choose $\xi \in \Gamma(M; E)$ such that $\xi(x) = v$ and let

$$T_v E^{\text{h}} = \text{Im} \{d\xi - \nabla \xi\}|_x \subset T_v E.$$

Since $\pi_E \circ \xi = \text{id}_M$,

$$d_v \pi_E \circ \{d\xi - \nabla \xi\}|_x = \text{id}_{T_x M} \quad \implies \quad T_v E \approx T_v E^{\text{h}} \oplus E_x \approx T_x M \oplus E_x.$$

This splitting of $T_v E$ satisfies (1.15) at v .

With the notation as in (1.11),

$$\{d\xi - \nabla \xi\}|_x = \left(d_x \text{id}_M, - \sum_{l=1}^{l=n} f^l(x) \theta_l^1|_x, \dots, - \sum_{l=1}^{l=n} f^l(x) \theta_l^n|_x \right): T_x M \longrightarrow T_x M \oplus \mathbb{R}^n$$

with respect to the identification $E|_U \approx U \times \mathbb{R}^k$ determined by the frame $(\xi_k)_k$. Thus, $T_v E^{\text{h}}$ is independent of the choice of ξ . Furthermore, the resulting splitting (1.14) of (1.5) extends (1.4) and satisfies (1.16). \square

1.2 Metric-compatible connections

Suppose $E \rightarrow M$ is a smooth vector bundle. Let g be a metric on E , i.e.

$$g \in \Gamma(M; E^* \otimes_{\mathbb{R}} E^*) \quad \text{s.t.} \quad g(v, w) = g(w, v), \quad g(v, v) > 0 \quad \forall v, w \in E_x, \quad v \neq 0, \quad x \in M.$$

A connection ∇ in E is g -compatible if

$$d(g(\xi, \zeta)) = g(\nabla \xi, \zeta) + g(\xi, \nabla \zeta) \in \Gamma(M; T^*M) \quad \forall \xi, \zeta \in \Gamma(M; E).$$

Suppose U is an open subset of M and $\xi_1, \dots, \xi_n \in \Gamma(U; E)$ is a frame for E on U . For $i, j = 1, \dots, n$, let

$$g_{ij} = g(\xi_i, \xi_j) \in C^\infty(U).$$

If ∇ is a connection in E and θ_{kl} is the connection 1-form for ∇ with respect to the frame $\{\xi_k\}_k$, then ∇ is g -compatible on U if and only if

$$\sum_{k=1}^{k=n} (g_{ik} \theta_j^k + g_{jk} \theta_i^k) = dg_{ij} \quad \forall i, j = 1, 2, \dots, n. \quad (1.17)$$

1.3 Torsion-free connections

If M is a smooth manifold, a connection ∇ in TM is torsion-free if

$$\nabla_X Y - \nabla_Y X = [X, Y].$$

If $(x_1, \dots, x_n): U \rightarrow \mathbb{R}^n$ is a coordinate chart on M , let

$$\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n} \in \Gamma(U; TM)$$

be the corresponding frame for TM on U . If ∇ is a connection in TM , the corresponding connection 1-form θ can be written as

$$\theta_j^k = \sum_{i=1}^{i=n} \Gamma_{ij}^k dx^i, \quad \text{where} \quad \nabla_{\partial/\partial x_i} \frac{\partial}{\partial x_j} = \sum_{k=1}^{k=n} \Gamma_{ij}^k \frac{\partial}{\partial x_k}.$$

The connection ∇ is torsion-free on $TM|_U$ if and only if

$$\Gamma_{ij}^k = \Gamma_{ji}^k \quad \forall i, j, k = 1, \dots, n. \quad (1.18)$$

Lemma 1.2. *If (M, g) is a Riemannian manifold, there exists a unique torsion-free g -compatible connection ∇ in TM .*

Proof. (1) Suppose ∇ and $\tilde{\nabla}$ are torsion-free g -compatible connections in TM . By (1.9), there exists

$$\begin{aligned} \theta &\in \Gamma(M; T^*M \otimes_{\mathbb{R}} \text{Hom}_{\mathbb{R}}(TM, TM)) \quad \text{s.t.} \\ \tilde{\nabla}_X Y - \nabla_X Y &= \{\theta(X)\}Y \quad \forall Y \in \Gamma(M; TM), X \in T_x M, x \in M. \end{aligned}$$

Since ∇ and $\tilde{\nabla}$ are torsion-free,

$$\{\theta(X)\}Y = \{\theta(Y)\}X \quad \forall X, Y \in T_x M, x \in M. \quad (1.19)$$

Since ∇ and $\tilde{\nabla}$ are g -compatible,

$$\begin{cases} g(\{\theta(X)\}Y, Z) + g(Y, \{\theta(X)\}Z) = 0 \\ g(\{\theta(Y)\}X, Z) + g(X, \{\theta(Y)\}Z) = 0 \\ g(\{\theta(Z)\}X, Y) + g(X, \{\theta(Z)\}Y) = 0 \end{cases} \quad \forall X, Y, Z \in T_x M, x \in M. \quad (1.20)$$

Adding the first two equations in (1.20), subtracting the third, and using (1.19) and the symmetry of g , we obtain

$$2g(\{\theta(X)\}Y, Z) = 0 \quad \forall X, Y, Z \in T_x M, x \in M \quad \implies \quad \theta \equiv 0.$$

Thus, $\tilde{\nabla} = \nabla$.

(2) Let $(x_1, \dots, x_n) : U \longrightarrow \mathbb{R}^n$ be a coordinate chart on M . With notation as in the paragraph preceding Lemma 1.2, ∇ is g -compatible on $TM|_U$ if and only if

$$\sum_{l=1}^{l=n} (g_{il}\Gamma_{kj}^l + g_{jl}\Gamma_{ki}^l) = \partial_{x_k} g_{ij}; \quad (1.21)$$

see (1.17). Define a connection ∇ in $TM|_U$ by

$$\Gamma_{ij}^k = \frac{1}{2} \sum_{l=1}^{l=n} g^{kl} (\partial_{x_i} g_{jl} + \partial_{x_j} g_{il} - \partial_{x_l} g_{ij}) \quad \forall i, j, k = 1, \dots, n,$$

where g^{ij} is the (i, j) -entry of the inverse of the matrix $(g_{ij})_{i,j=1,\dots,n}$. Since $g_{ij} = g_{ji}$, Γ_{ij}^k satisfies (1.18); a direct computation shows that Γ_{ij}^k also satisfies (1.21). Therefore, ∇ is a torsion-free g -compatible connection on $TM|_U$. In this way, we can define a torsion-free g -compatible connection on every coordinate chart. By the uniqueness property, these connections agree on the overlaps. \square

2 Complex structures

2.1 Complex linear connections

Suppose M is a smooth manifold and $\pi : (E, i) \longrightarrow M$ is a complex vector bundle. Similarly to Section 1.1, there is an exact sequence

$$0 \longrightarrow \pi_E^* E \xrightarrow{\iota_E} TE \xrightarrow{d\pi_E} \pi_E^* TM \longrightarrow 0 \quad (2.1)$$

of vector bundles over E . The homomorphism ι_E is now \mathbb{C} -linear. If $f \in C^\infty(M; \mathbb{C})$ and $m_f: E \rightarrow E$ is defined as in (1.1), there is a commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \pi_E^* E & \xrightarrow{\iota_E} & TE & \xrightarrow{d\pi_E} & \pi_E^* TM \longrightarrow 0 \\
& & \downarrow \pi_E^* m_f & & \downarrow dm_f & & \downarrow \pi_E^* \text{id} \\
0 & \longrightarrow & \pi_E^* E & \xrightarrow{m_f^* \iota_E} & m_f^* TE & \xrightarrow{m_f^* d\pi_E} & \pi_E^* TM \longrightarrow 0
\end{array} \tag{2.2}$$

of bundle maps over E .

Suppose

$$\nabla: \Gamma(M; E) \longrightarrow \Gamma(M; T^*M \otimes_{\mathbb{R}} E)$$

is a \mathbb{C} -linear connection, i.e.

$$\nabla_v(\mathbf{i}\xi) = \mathbf{i}(\nabla_v \xi) \quad \forall \xi \in \Gamma(M; E), v \in TM.$$

If U is an open subset of M and $\xi_1, \dots, \xi_n \in \Gamma(U; E)$ is a \mathbb{C} -frame for E on U , then there exist

$$\theta_l^k \in \Gamma(M; T^*M) \quad \text{s.t.} \quad \nabla \xi_l = \sum_{k=1}^{k=n} \xi_k \theta_l^k \equiv \sum_{k=1}^{k=n} \theta_l^k \otimes \xi_k \quad \forall l=1, \dots, n.$$

We will call

$$\theta \equiv (\theta_l^k)_{k,l=1,\dots,n} \in \Gamma(\Sigma; T^*M \otimes_{\mathbb{R}} \text{Mat}_n \mathbb{C})$$

the complex connection 1-form of ∇ with respect to the frame $(\xi_k)_k$. For an arbitrary section

$$\xi = \sum_{l=1}^{l=n} f^l \xi_l \in \Gamma(U; E),$$

by (1.8) and \mathbb{C} -linearity of ∇ we have

$$\nabla \xi = \sum_{k=1}^{k=n} \xi_k \left(df^k + \sum_{l=1}^{l=n} \theta_l^k f^l \right), \quad \text{i.e.} \quad \nabla(\underline{\xi} \cdot \underline{f}^t) = \underline{\xi} \cdot \{d + \theta\} \underline{f}^t, \tag{2.3}$$

where $\underline{\xi}$ and \underline{f} are as (1.12).

Let g be a hermitian metric on E , i.e.

$$g \in \Gamma(M; \text{Hom}_{\mathbb{C}}(\bar{E} \otimes_{\mathbb{C}} E, \mathbb{C})) \quad \text{s.t.} \quad g(v, w) = \overline{g(w, v)}, \quad g(v, v) > 0 \quad \forall v, w \in E_x, v \neq 0, x \in M.$$

A \mathbb{C} -linear connection ∇ in E is g -compatible if

$$d(g(\xi, \zeta)) = g(\nabla \xi, \zeta) + g(\xi, \nabla \zeta) \in \Gamma(M; T^*M \otimes_{\mathbb{R}} \mathbb{C}) \quad \forall \xi, \zeta \in \Gamma(M; E).$$

With notation as in the previous paragraph, let

$$g_{ij} = g(\xi_i, \xi_j) \in C^\infty(U; \mathbb{C}) \quad \forall i, j = 1, \dots, n.$$

Then ∇ is g -compatible on U if and only if

$$\sum_{k=1}^{k=n} (g_{ik} \theta_j^k + \bar{g}_{jk} \bar{\theta}_i^k) = dg_{ij} \quad \forall i, j = 1, 2, \dots, n. \tag{2.4}$$

2.2 Generalized $\bar{\partial}$ -operators

If (Σ, \mathfrak{j}) is an almost complex manifold, let

$$T^*\Sigma^{1,0} \equiv \{\eta \in T^*\Sigma \otimes_{\mathbb{R}} \mathbb{C} : \eta \circ \mathfrak{j} = \mathfrak{i}\eta\} \quad \text{and} \quad T^*\Sigma^{0,1} \equiv \{\eta \in T^*\Sigma \otimes_{\mathbb{R}} \mathbb{C} : \eta \circ \mathfrak{j} = -\mathfrak{i}\eta\}$$

be the bundles of \mathbb{C} -linear and \mathbb{C} -antilinear 1-forms on Σ . If (Σ, \mathfrak{j}) and (M, J) are smooth almost complex manifolds and $u: \Sigma \rightarrow M$ is a smooth function, define

$$\bar{\partial}_{J,\mathfrak{j}}u \in \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} u^*TM) \quad \text{by} \quad \bar{\partial}_{J,\mathfrak{j}}u = \frac{1}{2}(du + J \circ du \circ \mathfrak{j}). \quad (2.5)$$

A smooth map $u: (\Sigma, \mathfrak{j}) \rightarrow (M, J)$ will be called (J, \mathfrak{j}) -holomorphic if $\bar{\partial}_{J,\mathfrak{j}}u = 0$.

Definition 2.1. *Suppose (Σ, \mathfrak{j}) is an almost complex manifold and $\pi: (E, \mathfrak{i}) \rightarrow \Sigma$ is a complex vector bundle. A $\bar{\partial}$ -operator on (E, \mathfrak{i}) is a \mathbb{C} -linear map*

$$\bar{\partial}: \Gamma(\Sigma; E) \rightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} E)$$

such that

$$\bar{\partial}(f\xi) = (\bar{\partial}f) \otimes \xi + f(\bar{\partial}\xi) \quad \forall f \in C^\infty(\Sigma), \xi \in \Gamma(\Sigma; E), \quad (2.6)$$

where $\bar{\partial}f = \bar{\partial}_{\mathfrak{i},\mathfrak{j}}f$ is the usual $\bar{\partial}$ -operator on complex-valued functions.

Similarly to Section 1.1, a $\bar{\partial}$ -operator on (E, \mathfrak{i}) is a first-order differential operator. If U is an open subset of M and $\xi_1, \dots, \xi_n \in \Gamma(U; E)$ is a \mathbb{C} -frame for E on U , then there exist

$$\theta_l^k \in \Gamma(U; T^*U^{0,1}) \quad \text{s.t.} \quad \bar{\partial}\xi_l = \sum_{k=1}^{k=n} \xi_k \theta_l^k \equiv \sum_{k=1}^{k=n} \theta_l^k \otimes \xi_k \quad \forall l=1, \dots, n.$$

We call

$$\theta \equiv (\theta_l^k)_{k,l=1,\dots,n} \in \Gamma(U; T^*U^{0,1} \otimes_{\mathbb{C}} \text{Mat}_n \mathbb{C})$$

the connection 1-form of $\bar{\partial}$ with respect to the frame $(\xi_k)_k$. For an arbitrary section

$$\xi = \sum_{l=1}^{l=n} f^l \xi_l \in \Gamma(U; E),$$

by (2.6) we have

$$\bar{\partial}\xi = \sum_{k=1}^{k=n} \xi_k \left(\bar{\partial}f^k + \sum_{l=1}^{l=n} \theta_l^k f^l \right), \quad \text{i.e.} \quad \bar{\partial}(\underline{\xi} \cdot \underline{f}) = \underline{\xi} \cdot \{\bar{\partial} + \theta\} \underline{f}, \quad (2.7)$$

where $\underline{\xi}$ and \underline{f} are as in (1.12). It is immediate from (2.6) that the symbol of $\bar{\partial}$ is given by

$$\sigma_{\bar{\partial}}: T^*\Sigma \rightarrow \text{Hom}_{\mathbb{C}}(E, T^*\Sigma^{0,1} \otimes_{\mathbb{C}} E), \quad \{\sigma_{\bar{\partial}}(\eta)\}(f) = (\eta + \mathfrak{i}\eta \circ \mathfrak{j}) \otimes f.$$

In particular, $\bar{\partial}$ is an elliptic operator (i.e. $\sigma_{\bar{\partial}}(\eta)$ is an isomorphism for $\eta \neq 0$) if (Σ, \mathfrak{j}) is a Riemann surface.

Lemma 2.2. *Suppose (Σ, \mathfrak{j}) is an almost complex manifold and $\pi: (E, \mathfrak{i}) \rightarrow \Sigma$ is a complex vector bundle. If*

$$\bar{\partial}: \Gamma(\Sigma; E) \longrightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} E)$$

*is a $\bar{\partial}$ -operator on (E, \mathfrak{i}) , there exists a unique almost complex structure $J = J_{\bar{\partial}}$ on (the total space of) E such that π is a (\mathfrak{j}, J) -holomorphic map, the restriction of J to the vertical tangent bundle $TE^{\vee} \approx \pi^*E$ agrees with \mathfrak{i} , and*

$$\bar{\partial}_{J, \mathfrak{j}} \xi = 0 \in \Gamma(U; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} \xi^*TE) \quad \iff \quad \bar{\partial} \xi = 0 \in \Gamma(U; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} E) \quad (2.8)$$

for every open subset U of Σ and $\xi \in \Gamma(U; E)$.

Proof. (1) With notation as above, define

$$\varphi: U \times \mathbb{C}^n \longrightarrow E|_U \quad \text{by} \quad \varphi(x, c^1, \dots, c^n) = \underline{\xi}(x) \cdot \underline{c}^t \equiv \sum_{k=1}^{k=n} c^k \xi_k(x) \in E_x.$$

The map φ is a trivialization of E over U . If $J \equiv J_{\bar{\partial}}$ is an almost complex structure on E , let \tilde{J} be the almost complex structure on $U \times \mathbb{C}^n$ given by

$$\tilde{J}_{(x, \underline{c})} = \{d_{(x, \underline{c})} \varphi\}^{-1} \circ J_{\varphi(x, \underline{c})} \circ d_{(x, \underline{c})} \varphi \quad \forall (x, \underline{c}) \in U \times \mathbb{C}^n. \quad (2.9)$$

The almost complex structure J restricts to \mathfrak{i} on TE^{\vee} if and only if

$$\tilde{J}_{(x, \underline{c})} w = \mathfrak{i} w \in T_{\underline{c}} \mathbb{C}^n \subset T_{(x, \underline{c})}(U \times \mathbb{C}^n) \quad \forall w \in T_{\underline{c}} \mathbb{C}^n. \quad (2.10)$$

If J restricts to \mathfrak{i} on TE^{\vee} , the projection π is (\mathfrak{j}, J) -holomorphic on $E|_U$ if and only if there exists

$$\begin{aligned} \tilde{J}^{\text{vh}} &\in \Gamma(U \times \mathbb{C}^n; \text{Hom}_{\mathbb{R}}(\pi_U^* TU, \pi_{\mathbb{C}^n}^* T\mathbb{C}^n)) \quad \text{s.t.} \\ \tilde{J}_{(x, \underline{c})} w &= \mathfrak{j}_x w + \tilde{J}_{(x, \underline{c})}^{\text{vh}} w \quad \forall w \in T_x U \subset T_{(x, \underline{c})}(U \times \mathbb{C}^n). \end{aligned} \quad (2.11)$$

If $\xi \in \Gamma(U; E)$, let

$$\tilde{\xi} \equiv \varphi^{-1} \circ \xi \equiv (\text{id}_U, \underline{f}), \quad \text{where} \quad \underline{f} \in C^\infty(U; \mathbb{C}^n).$$

By (2.9)-(2.11),

$$\begin{aligned} 2 \bar{\partial}_{J, \mathfrak{j}} \xi|_x &= d_{\tilde{\xi}(x)} \varphi \circ 2 \bar{\partial}_{\tilde{J}, \mathfrak{j}} \tilde{\xi}|_x = d_{\tilde{\xi}(x)} \varphi \circ \{(\text{Id}_{T_x U}, d_x \underline{f}) + \tilde{J}_{\tilde{\xi}(x)} \circ (\text{Id}_{T_x U}, d_x \underline{f}) \circ \mathfrak{j}_x\} \\ &= d_{\tilde{\xi}(x)} \varphi \circ (0, 2 \bar{\partial} \underline{f}|_x + \tilde{J}_{\tilde{\xi}(x)}^{\text{vh}} \circ \mathfrak{j}_x). \end{aligned} \quad (2.12)$$

On the other hand, by (2.7),

$$\begin{aligned} \bar{\partial} \xi|_x &= \bar{\partial}(\underline{\xi} \cdot f^t)|_x = \underline{\xi}(x) \cdot \{\bar{\partial} + \theta\} f^t|_x \\ &= \varphi(\bar{\partial} \underline{f}|_x + \theta_x \cdot f(x)^t). \end{aligned} \quad (2.13)$$

By (2.12) and (2.13), the property (2.8) is satisfied for all $\xi \in \Gamma(U; E)$ if and only if

$$\tilde{J}_{(x, \underline{c})}^{\text{vh}} = 2(\theta_x \cdot \underline{c}^t) \circ (-\mathfrak{j}_x) = 2\mathfrak{i} \theta_x \cdot \underline{c}^t \quad \forall (x, \underline{c}) \in U \times \mathbb{C}^n.$$

In summary, the almost complex structure $J = J_{\bar{\partial}}$ on E has the three desired properties if and only if for every trivialization of E over an open subset U of Σ

$$\begin{aligned} \tilde{J}_{(x, \underline{c})}(w_1, w_2) &= (j_x w_1, i w_2 + 2i\theta_x(w_1) \cdot \underline{c}^t) \\ \forall (x, \underline{c}) \in U \times \mathbb{C}^n, (w_1, w_2) \in T_x U \oplus T_{\underline{c}} \mathbb{C}^n &= T_{(x, \underline{c})}(U \times \mathbb{C}^n), \end{aligned} \quad (2.14)$$

where \tilde{J} is the almost complex structure on $U \times \mathbb{C}^n$ induced by J via the trivialization and θ is the connection 1-form corresponding to $\bar{\partial}$ with respect to the frame inducing the trivialization.

(2) By (2.14), there exists at most one almost complex structure J satisfying the three properties. Conversely, (2.14) determines such an almost complex structure on E . Since

$$\begin{aligned} \tilde{J}_{(x, \underline{c})}^2(w_1, w_2) &= \tilde{J}_{(x, \underline{c})}(j w_1, i w_2 + 2i\theta_x(w_1) \cdot \underline{c}^t) = (j^2 w_1, i(i w_2 + 2i\theta_x(w_1) \cdot \underline{c}^t) + 2i\theta_x(j w_1) \cdot \underline{c}^t) \\ &= -(w_1, w_2), \end{aligned}$$

\tilde{J} is indeed an almost complex structure on E . The almost complex structure induced by \tilde{J} on $E|_U$ satisfies the three properties by part (a). By the uniqueness property, the almost complex structures on E induced by the different trivializations agree on the overlaps. Therefore, they define an almost complex structure $J = J_{\bar{\partial}}$ on the total space of E with the desired properties. \square

2.3 Connections and $\bar{\partial}$ -operators

Suppose (Σ, j) is an almost complex manifold, $\pi: (E, i) \rightarrow \Sigma$ is a complex vector bundle, and

$$\bar{\partial}: \Gamma(\Sigma; E) \rightarrow \Gamma(\Sigma; T^* \Sigma^{0,1} \otimes_{\mathbb{C}} E)$$

is a $\bar{\partial}$ -operator on (E, i) . A \mathbb{C} -linear connection ∇ in (E, i) is $\bar{\partial}$ -compatible if

$$\bar{\partial} \xi = \bar{\partial}_{\nabla} \xi \equiv \frac{1}{2}(\nabla \xi + i \nabla \xi \circ j) \quad \forall \xi \in \Gamma(M; \Sigma). \quad (2.15)$$

Lemma 2.3. *Suppose (Σ, j) is an almost complex manifold, $\pi: (E, i) \rightarrow \Sigma$ is a complex vector bundle,*

$$\bar{\partial}: \Gamma(\Sigma; E) \rightarrow \Gamma(\Sigma; T^* \Sigma^{0,1} \otimes_{\mathbb{C}} E)$$

is a $\bar{\partial}$ -operator on (E, i) , and $J_{\bar{\partial}}$ is the complex structure in the vector bundle $TE \rightarrow E$ provided by Lemma 2.2. A \mathbb{C} -linear connection ∇ in (E, i) is $\bar{\partial}$ -compatible if and only if the splitting (1.14) determined by ∇ respects the complex structures.

Proof. Since $J_{\bar{\partial}} = \pi^* i$ on $\pi^* E \subset TE$, the splitting (1.14) determined by ∇ respects the complex structures if and only if

$$J_{\bar{\partial}}|_v \circ \{d\xi - \nabla \xi\}|_x = \{d\xi - \nabla \xi\}|_x \circ j_x: T_x \Sigma \rightarrow T_v E$$

for all $x \in \Sigma$, $v \in E_x$, and $\xi \in \Gamma(\Sigma; E)$ such that $\xi(x) = 0$; see the proof of Lemma 1.1. This identity is equivalent to

$$\bar{\partial}_{J_{\bar{\partial}}, j} \xi = \bar{\partial}_{\nabla} \xi \quad \forall \xi \in \Gamma(\Sigma; E). \quad (2.16)$$

On the other hand, by the proof of Lemma 2.2,

$$\bar{\partial}_{J_{\bar{\partial}}, j} \xi = \bar{\partial} \xi \quad \forall \xi \in \Gamma(\Sigma; E); \quad (2.17)$$

see (2.12)-(2.14). The lemma follows immediately from (2.16) and (2.17). \square

2.4 Holomorphic vector bundles

Let (Σ, j) be a complex manifold. A holomorphic vector bundle (E, i) on (Σ, j) is a complex vector bundle with a collection of trivializations that overlap holomorphically.

A collection of holomorphically overlapping trivializations of (E, i) determines a holomorphic structure J on the total space of E and a $\bar{\partial}$ -operator

$$\bar{\partial}: \Gamma(\Sigma; E) \longrightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} E).$$

The latter is defined as follows. If ξ_1, \dots, ξ_n is a holomorphic complex frame for E over an open subset U of M , then

$$\bar{\partial} \sum_{k=1}^{k=n} f^k \xi_k = \sum_{k=1}^{k=n} \bar{\partial} f^k \otimes \xi_k \quad \forall f^1, \dots, f^k \in C^\infty(U; \mathbb{C}).$$

In particular, for all $\xi \in \Gamma(M; E)$

$$\bar{\partial}_{J,j} \xi = 0 \quad \iff \quad \bar{\partial} \xi = 0.$$

Thus, $J = J_{\bar{\partial}}$; see Lemma 2.2.

Lemma 2.4. *Suppose (Σ, j) is a Riemann surface and $\pi: (E, i) \rightarrow \Sigma$ is a complex vector bundle. If*

$$\bar{\partial}: \Gamma(\Sigma; E) \longrightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} E)$$

is a $\bar{\partial}$ -operator on (E, i) , the almost complex structure $J = J_{\bar{\partial}}$ on E is integrable. With this complex structure, $\pi: E \rightarrow \Sigma$ is a holomorphic vector bundle and $\bar{\partial}$ is the corresponding $\bar{\partial}$ -operator.

Proof. By (2.8), it is sufficient to show that there exists a (J, j) -holomorphic local section through every point $v \in E$, i.e. there exist a neighborhood U of $x \equiv \pi(v)$ in Σ and $\xi \in \Gamma(U; E)$ such that

$$\xi(x) = v \quad \text{and} \quad \bar{\partial}_{J,j} \xi = 0.$$

By Lemma 2.2 and (2.13), this is equivalent to showing that the equation

$$\left\{ \bar{\partial} + \theta \right\} f^t = 0, \quad f(x) = v, \quad f \in C^\infty(U; \mathbb{C}^n), \quad (2.18)$$

has a solution for every $v \in \mathbb{C}^n$. We can assume that U is a small disk contained in S^2 . Let

$$\eta: S^2 \longrightarrow [0, 1]$$

be a smooth function supported in U and such that $\eta \equiv 1$ on a neighborhood of x . Then,

$$\eta\theta \in \Gamma(S^2; (T^*S^2)^{0,1} \otimes_{\mathbb{C}} \text{Mat}_n \mathbb{C}).$$

Choose $p > 2$. The operator

$$\Theta: L_1^p(S^2; \mathbb{C}^n) \longrightarrow L^p(S^2; (T^*S^2)^{0,1} \otimes_{\mathbb{C}} \mathbb{C}^n) \oplus \mathbb{C}^n, \quad \Theta(f) = (\bar{\partial}_{i,j} f, f(x)),$$

is surjective. If η has sufficiently small support, so is the operator

$$\Theta_\eta: L_1^p(S^2; \mathbb{C}^n) \longrightarrow L^p(S^2; (T^*S^2)^{0,1} \otimes_{\mathbb{C}} \mathbb{C}^n) \oplus \mathbb{C}^n, \quad \Theta_\eta(f) = (\{\bar{\partial}_{i,j} + \eta\theta\} f, f(x)).$$

Then, the restriction of $\Theta_\eta^{-1}(0, v)$ to a neighborhood of x on which $\eta \equiv 1$ is a solution of (2.18). By elliptic regularity, $\Theta_\eta^{-1}(0, v) \in C^\infty(S^2; \mathbb{C}^n)$. \square

2.5 Deformations of almost complex submanifolds

If (M, J) is a complex manifold, holomorphic coordinate charts on (M, J) determine a holomorphic structure in the vector bundle $(TM, i) \rightarrow M$. If $(\Sigma, j) \subset (M, J)$ is a complex submanifold, holomorphic coordinate charts on Σ can be extended to holomorphic coordinate charts on M . Thus, the holomorphic structure in $T\Sigma \rightarrow \Sigma$ induced from (Σ, j) is the restriction of the holomorphic structure in $TM|_\Sigma$. It follows that

$$\bar{\partial}_M = \bar{\partial}_\Sigma: \Gamma(\Sigma; T\Sigma) \rightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} T\Sigma) \subset \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} TM|_\Sigma),$$

where $\bar{\partial}_M$ and $\bar{\partial}_\Sigma$ are the $\bar{\partial}$ -operators in $TM|_\Sigma$ and $T\Sigma$ induced from the holomorphic structures in Σ and M . Therefore, $\bar{\partial}_M$ descends to a $\bar{\partial}$ -operator on the quotient

$$\bar{\partial}: \Gamma(\Sigma; \mathcal{N}_M \Sigma) = \Gamma(\Sigma; TM|_\Sigma) / \Gamma(\Sigma; T\Sigma) \rightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} \mathcal{N}_M \Sigma),$$

where

$$\mathcal{N}_M \Sigma \equiv TM|_\Sigma / T\Sigma \rightarrow \Sigma$$

is the normal bundle of Σ in M . This vector bundle inherits a holomorphic structure from that of $TM|_\Sigma$ and Σ . The above $\bar{\partial}$ -operator on $\mathcal{N}_M \Sigma$ is the $\bar{\partial}$ -operator corresponding to this induced holomorphic structure on $\mathcal{N}_M \Sigma$.

Suppose (M, J) is an almost complex manifold and $(\Sigma, j) \subset (M, J)$ is an almost complex submanifold. Let ∇ be a torsion-free connection in TM . Define

$$\begin{aligned} D_{J;\Sigma}: \Gamma(\Sigma; TM|_\Sigma) &\rightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} TM|_\Sigma) && \text{by} \\ D_{J;\Sigma}\xi &= \frac{1}{2}(\nabla\xi + J \circ \nabla\xi \circ j) - \frac{1}{2}J \circ \nabla_\xi J: T\Sigma \rightarrow TM|_\Sigma. \end{aligned} \quad (2.19)$$

If ∇ is the Levi-Civita connection (the connection of Lemma 1.2) for a J -compatible metric on M (and Σ is a Riemann surface), then $D_{J;\Sigma}$ is the linearization of the $\bar{\partial}_J$ -operator at the inclusion map $\iota: \Sigma \rightarrow M$; see [4, Proposition 3.1.1].

In fact, $D_{J;\Sigma}$ is independent of the choice of a torsion-free connection in TM . Let

$$\tilde{\nabla} = \nabla + \theta, \quad \theta \in \Gamma(M; T^*M \otimes_{\mathbb{R}} \text{Hom}_{\mathbb{R}}(TM, TM)), \quad (2.20)$$

be another torsion-free connection; see (1.9). Since $\tilde{\nabla}$ and ∇ are torsion-free connections,

$$\{\theta(X)\}Y = \{\theta(Y)\}X \quad \forall X, Y \in T_x M, x \in M. \quad (2.21)$$

If $x \in M$ and $X, Y \in \Gamma(M; TM)$,

$$\begin{aligned} \{\nabla_Y J\}X &= \nabla_Y(JX) - J\nabla_Y X, & \{\tilde{\nabla}_Y J\}X &= \tilde{\nabla}_Y(JX) - J\tilde{\nabla}_Y X && \implies \\ \{\tilde{\nabla}_Y J\}X - \{\nabla_Y J\}X &= \{\theta(Y)\}(JX) - J\{\theta(Y)\}X = \{\theta(JX)\}Y - J\{\theta(X)\}Y \end{aligned} \quad (2.22)$$

by (2.20) and (2.21). On the other hand, by (2.20) for all $X \in T\Sigma$ and $\xi \in \Gamma(\Sigma; TM|_\Sigma)$,

$$\begin{aligned} \{\tilde{\nabla}\xi + J \circ \tilde{\nabla}\xi \circ j\}(X) - \{\nabla\xi + J \circ \nabla\xi \circ j\}(X) &= \{\theta(X)\}\xi + J\{\theta(jX)\}\xi \\ &= J(\{\theta(JX)\}\xi - J\{\theta(X)\}\xi), \end{aligned} \quad (2.23)$$

since $\mathfrak{j} = J|_{T\Sigma}$ and $J^2 = -\text{Id}$. By (2.22) and (2.23), $D_{J,\Sigma}$ is independent of the choice of torsion-free connection ∇ .

Since any torsion-free connection on Σ extends to a torsion-free connection on M , the above observation implies that

$$D_{J,\Sigma} : \Gamma(\Sigma; T\Sigma) \longrightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} T\Sigma) \subset \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} TM|_{\Sigma}). \quad (2.24)$$

Thus, an almost complex submanifold (Σ, \mathfrak{j}) of an almost complex manifold (M, J) induces a well-defined generalized Cauchy-Riemann operator¹ on the normal bundle of Σ in M ,

$$D_{J,\Sigma}^{\mathcal{N}} : \Gamma(\Sigma; \mathcal{N}_M\Sigma) \longrightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} \mathcal{N}_M\Sigma), \quad D_{J,\Sigma}^{\mathcal{N}}(\pi(\xi)) = \pi(D_{J,\Sigma}(\xi)) \quad \forall \xi \in \Gamma(\Sigma; TM|_{\Sigma}),$$

where $\pi : TM|_{\Sigma} \longrightarrow \mathcal{N}_M\Sigma$ is the quotient projection map. The \mathbb{C} -linear part of $D_{J,\Sigma}^{\mathcal{N}}$ determines a $\bar{\partial}$ -operator on the normal bundle of Σ in M :

$$\begin{aligned} \bar{\partial}_{J,\Sigma}^{\mathcal{N}} : \Gamma(\Sigma; \mathcal{N}_M\Sigma) &\longrightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} \mathcal{N}_M\Sigma), \\ \bar{\partial}_{J,\Sigma}^{\mathcal{N}}(\xi) &= \frac{1}{2}(D_{J,\Sigma}^{\mathcal{N}}(\xi) - JD_{J,\Sigma}^{\mathcal{N}}(J\xi)) \quad \forall \xi \in \Gamma(\Sigma; \mathcal{N}_M\Sigma). \end{aligned}$$

Both operators are determined by the almost complex submanifold (Σ, \mathfrak{j}) of the almost complex manifold (M, J) only and are independent of the choice of torsion-free connection ∇ in (2.19).

Any connection ∇ in TM induces a J -linear connection in TM by

$$\nabla_X^J \xi = \nabla_X \xi - \frac{1}{2}J(\nabla_X J)\xi \quad \forall X \in TM, \xi \in \Gamma(M; TM). \quad (2.25)$$

If ∇ is as in (2.19),

$$\{D_{J,\Sigma}\xi\}(X) = \{\bar{\partial}_{\nabla^J}\xi\}(X) + A_J(X, \xi) - \frac{1}{4}\{(\nabla_{J\xi}J) + J(\nabla_{\xi}J)\}(X) \quad (2.26)$$

for all $\xi \in \Gamma(\Sigma; TM|_{\Sigma})$ and $X \in T\Sigma$, where A_J is the Nijenhuis tensor of J :

$$A_J(\xi_1, \xi_2) = \frac{1}{4}\left([\xi_1, \xi_2] + J[\xi_1, J\xi_2] + J[J\xi_1, \xi_2] - [J\xi_1, J\xi_2]\right) \quad \forall \xi_1, \xi_2 \in \Gamma(M; TM). \quad (2.27)$$

Since the sum of the terms in the curly brackets in (2.26) is \mathbb{C} -linear in ξ , while the Nijenhuis tensor is \mathbb{C} -antilinear, the \mathbb{C} -linear operator

$$\Gamma(\Sigma; TM|_{\Sigma}) \longrightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} TM|_{\Sigma}), \quad \xi \longrightarrow \bar{\partial}_{\nabla^J}(\xi) - \frac{1}{4}\{(\nabla_{J\xi}J) + J(\nabla_{\xi}J)\}, \quad (2.28)$$

takes $\Gamma(\Sigma; T\Sigma)$ to $\Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} T\Sigma)$ by (2.24). Thus, it induces a $\bar{\partial}$ -operator on $\mathcal{N}_M\Sigma$ and this induced operator is $\bar{\partial}_{J,\Sigma}^{\mathcal{N}}$. If the image of the homomorphism

$$TM \longrightarrow T^*\Sigma^{0,1} \otimes_{\mathbb{C}} TM|_{\Sigma}, \quad \xi \longrightarrow \nabla_{\xi}J - J\nabla_{J\xi}J,$$

¹see Section 4.3

is contained in $T^*\Sigma^{0,1} \otimes_{\mathbb{C}} T\Sigma$, then $\bar{\partial}_{\nabla^J}$ preserves $T\Sigma$ and induces a $\bar{\partial}$ -operator $\bar{\partial}_{\nabla^J}^{\mathcal{N}}$ on $\mathcal{N}_M\Sigma$ with $\bar{\partial}_{\nabla^J}^{\mathcal{N}} = \bar{\partial}_{J;\Sigma}^{\mathcal{N}}$. In this case,

$$D_{J;\Sigma}^{\mathcal{N}}(\pi(\xi)) = \pi(\bar{\partial}_{\nabla^J}\xi + A_J(\cdot, \xi)): T\Sigma \longrightarrow \mathcal{N}_M\Sigma \quad \forall \xi \in \Gamma(\Sigma; TM|_{\Sigma}).$$

This is the case in particular if J is compatible with a symplectic form ω on M and ∇ is the Levi-Civita connection for the metric $g(\cdot, \cdot) = \omega(\cdot, J\cdot)$, as the sum in the curly brackets in (2.26) then vanishes by [4, (C.7.5)].

It is immediate that A_J takes $T\Sigma \otimes_{\mathbb{R}} T\Sigma$ to $T\Sigma$ and thus induces a bundle homomorphism

$$A_J^{\mathcal{N}}: T\Sigma \otimes_{\mathbb{R}} \mathcal{N}_M\Sigma \longrightarrow \mathcal{N}_M\Sigma.$$

If ζ is any vector field on M such that $\zeta(x) = X \in T_x\Sigma$ for some $x \in \Sigma$, then

$$\begin{aligned} \{D_{J;\Sigma}\xi\}(X) &= \frac{1}{2}([\zeta, \xi] + J[J\zeta, \xi])|_x, \\ \left\{\bar{\partial}_{\nabla^J}(\xi) - \frac{1}{4}((\nabla_{J\xi}J) + J(\nabla_{\xi}J))\right\}(X) &= \frac{1}{4}([\zeta, \xi] + J[J\zeta, \xi] - J[\zeta, J\xi] + [J\zeta, J\xi])|_x, \end{aligned} \quad (2.29)$$

since ∇ is torsion-free.² These two identities immediately imply that the operators (2.19) and (2.28) preserve $T\Sigma \subset TM|_{\Sigma}$ and thus induce operators

$$\Gamma(\Sigma; \mathcal{N}_M\Sigma) \longrightarrow \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} \mathcal{N}_M\Sigma)$$

as claimed above.

If g is a J -compatible metric on $TM|_{\Sigma}$ and $\pi^{\perp}: TM|_{\Sigma} \longrightarrow T\Sigma^{\perp}$ is the projection to the g -orthogonal complement of $T\Sigma$ in $TM|_{\Sigma}$, the composition ∇^{\perp}

$$\Gamma(\Sigma; T\Sigma^{\perp}) \hookrightarrow \Gamma(\Sigma; TM|_{\Sigma}) \xrightarrow{\nabla^J} \Gamma(\Sigma; T^*\Sigma \otimes_{\mathbb{R}} TM|_{\Sigma}) \xrightarrow{\pi^{\perp}} \Gamma(\Sigma; T^*\Sigma \otimes_{\mathbb{R}} T\Sigma^{\perp}),$$

with ∇^J as in (2.25), is a g -compatible J -linear connection in $T\Sigma^{\perp}$. Via the isomorphism $\pi: T\Sigma^{\perp} \longrightarrow \mathcal{N}_M\Sigma$, it induces a J -linear connection $\nabla^{\mathcal{N}}$ in $\mathcal{N}_M\Sigma$ which is compatible with the metric $g^{\mathcal{N}}$ induced via this isomorphism from $g|_{T\Sigma^{\perp}}$. If the image of the homomorphism

$$T\Sigma^{\perp} \longrightarrow T^*\Sigma^{0,1} \otimes_{\mathbb{C}} TM|_{\Sigma}, \quad \xi \longrightarrow \nabla_{\xi}J - J\nabla_{J\xi}J, \quad (2.30)$$

is contained in $T^*\Sigma^{0,1} \otimes_{\mathbb{C}} T\Sigma$, then $\bar{\partial}_{\nabla^{\mathcal{N}}} = \bar{\partial}_{J;\Sigma}^{\mathcal{N}}$ and so

$$D_{J;\Sigma}^{\mathcal{N}}(\pi(\xi)) = \pi(\bar{\partial}_{\nabla^{\perp}}\xi + A_J(\cdot, \xi)): T\Sigma \longrightarrow \mathcal{N}_M\Sigma \quad \forall \xi \in \Gamma(\Sigma; T\Sigma^{\perp}).$$

This is the case if Σ is a divisor in M , i.e. $\text{rk}_{\mathbb{C}}\mathcal{N} = 1$, since $(\nabla_{\zeta}J)\xi$ is g -orthogonal to ξ and $J\xi$ for all $\xi, \zeta \in T_xM$ and $x \in M$ by [4, (C.7.1)]. This is also the case if J is compatible with a symplectic form ω on M and $g(\cdot, \cdot) = \omega(\cdot, J\cdot)$, as the homomorphism (2.30) is then trivial by [4, (C.7.5)].

²Since LHS and RHS of these identities depend only ξ and $X = \zeta(x)$, and not on ζ , it is sufficient to verify them under the assumption that $\nabla_{\zeta}|_x = 0$.

3 Riemannian geometry estimates

This section is based on [1, Chapter 1] and [2, Section 3] and culminates in a Poincare lemma for closed curves in Proposition 3.6 and an expansion for the $\bar{\partial}$ -operator in Proposition 3.13. If $u: \Sigma \rightarrow M$ is a smooth map between smooth manifolds and $E \rightarrow M$ is a smooth vector bundle, let

$$\Gamma(u; E) = \Gamma(\Sigma; u^*E), \quad \Gamma^1(u; E) = \Gamma(\Sigma; T^*\Sigma \otimes_{\mathbb{R}} u^*E).$$

We denote the subspace of compactly supported sections in $\Gamma(u; E)$ by $\Gamma_c(u; E)$.

An exponential-like map on a smooth manifold M is a smooth map $\exp: TM \rightarrow M$ such that $\exp|_M = \text{id}_M$ and

$$d_x \exp = (\text{id}_{T_x M} \text{id}_{T_x M}): T_x(TM) = T_x M \oplus T_x M \rightarrow T_x M \quad \forall x \in M,$$

where the second equality is the canonical splitting of $T_x(TM)$ into the horizontal and vertical tangent space along the zero section. Any connection ∇ in TM gives rise to a smooth map $\exp^\nabla: W \rightarrow M$ from some neighborhood W of the zero section M in TM ; see [1, Section 1.3]. If $\eta: TM \rightarrow \mathbb{R}$ is a smooth function which equals 1 on a neighborhood of M in TM and 0 outside of W , then

$$\exp: TM \rightarrow M, \quad v \rightarrow \exp^\nabla(\eta(v)v),$$

is an exponential-like map. If M is compact, then W can be taken to be all of TM and $\exp = \exp^\nabla$.

If (M, g, \exp) is a Riemannian manifold with an exponential-like map and $x \in M$, let $r_{\exp}(x) \in \mathbb{R}^+$ be the supremum of the numbers $r \in \mathbb{R}$ such that the restriction

$$\exp: \{v \in T_x M: |v| < r\} \rightarrow M$$

is a diffeomorphism onto an open subset of M . Set

$$r_{\exp}^g(x) = \inf \{d_g(x, \exp(v)): v \in T_x M, |v| = r_{\exp}(x)\} \in \mathbb{R}^+,$$

where d_g is the metric on M induced by g . If $K \subset M$, let

$$r_{\exp}^g(K) = \inf_{x \in K} r_{\exp}^g(x);$$

this number is positive if $\bar{K} \subset M$ is compact.

3.1 Parallel transport

Let $(E, \langle, \rangle, \nabla) \rightarrow M$ be a vector bundle, real or complex, with an inner-product \langle, \rangle and a metric-compatible connection ∇ . If $\alpha: (a, b) \rightarrow M$ is a piecewise smooth curve, denote by

$$\Pi_\alpha: E_{\alpha(a)} \rightarrow E_{\alpha(b)}$$

the parallel-transport map along α with respect to the connection ∇ . If $\exp: TM \rightarrow M$ is an exponential-like map, $x \in M$, and $v \in T_x M$, let

$$\Pi_v: E_x \rightarrow E_{\exp(v)}$$

be the parallel transport along the curve

$$\gamma_v: [0, 1] \longrightarrow M, \quad \gamma_v(t) = \exp(tv).$$

If $u: [a, b] \times [c, d] \longrightarrow M$ is a smooth map, let

$$\Pi_{\partial u}: E_{u(a,c)} \longrightarrow E_{u(a,c)}$$

be the parallel transport along u restricted to the boundary of the rectangle traversed in the positive direction. If $u: \Sigma \longrightarrow M$ is any smooth map, ∇ induces a connection

$$\nabla^u: \Gamma(u; E) \longrightarrow \Gamma^1(u; E)$$

in the vector bundle $u^*E \longrightarrow \Sigma$. If α is a smooth curve as above and $\zeta \in \Gamma(\alpha; E)$, let

$$\frac{D}{dt}\zeta = \nabla_{\partial_t}^{\alpha} \zeta \in \Gamma(\alpha; E),$$

where ∂_t is the standard unit vector field on \mathbb{R} .

Lemma 3.1. *If (M, g) is a Riemannian manifold and $(E, \langle \cdot, \cdot \rangle, \nabla)$ is a normed vector bundle with connection over M , for every compact subset $K \subset M$ there exists $C_K \in \mathbb{R}^+$ such that for every smooth map $u: [a, b] \times [c, d] \longrightarrow M$ with $\text{Im } u \subset K$*

$$\|\Pi_{\partial u} - \mathbb{I}\| \leq C_K \int_c^d \int_a^b |u_s| |u_t| ds dt,$$

where the norm of $(\Pi_{\partial u} - \mathbb{I}) \in \text{End}(E_{u(a,c)})$ is computed with respect to the inner-product in $E_{u(a,c)}$.

Proof. (1) Choose an orthonormal frame $\{v_i\}$ for $E_{u(a,c)}$. Extend each v_i to

$$\xi_i \in \Gamma(u|_{[a, b] \times [c, d]}; E)$$

by parallel-transporting along the curve $t \longrightarrow u(a, t)$ and then to $\zeta_i \in \Gamma(u; E)$ by parallel-transporting $\xi_i(a, t)$ along the curve $s \longrightarrow u(s, t)$; see Figure 1. By construction,

$$\frac{D}{ds}\zeta_i = 0 \in \Gamma(u; E).$$

Let A be the matrix-valued function on $[a, b] \times [c, d]$ such that

$$\frac{D}{dt}\zeta_i \Big|_{(s,t)} = \sum_{l=1}^{l=k} A_{il}(s, t) \zeta_l(s, t), \quad (3.1)$$

where k is the rank of E . Note that $A_{ij}(a, t) = 0$ and

$$\langle \mathcal{R}_{\nabla}(u_s, u_t) \zeta_i, \zeta_j \rangle = \left\langle \frac{D}{ds} \frac{D}{dt} \zeta_i - \frac{D}{dt} \frac{D}{ds} \zeta_i, \zeta_j \right\rangle = \sum_{l=1}^{l=k} \left\langle \left(\frac{\partial}{\partial s} A_{il} \right) \zeta_l, \zeta_j \right\rangle = \frac{\partial}{\partial s} A_{ij}, \quad (3.2)$$

where \mathcal{R}_{∇} is the curvature tensor of the connection of ∇ . Since K is compact and the image of u is contained in K , it follows that

$$|A_{ij}(b, t)| \leq C_K \int_a^b |u_s|_{(s,t)} |u_t|_{(s,t)} ds. \quad (3.3)$$




Figure 1: Extending a basis $\{v_i\}$ for $E_{u(a,c)}$ to a frame $\{\zeta_i\}$ over $[a, b] \times [c, d]$

(2) The parallel transport of ζ_i along the curves

$$\tau \longrightarrow u(\tau, c), \quad \tau \longrightarrow u(\tau, d), \quad \tau \longrightarrow u(a, \tau)$$

is ζ_i itself. Thus, it remains to estimate the parallel transport of each ζ_i along the curve $\tau \longrightarrow u(b, \tau)$. Let h_{ij} be the SO_k -valued function (U_k -valued function if E is complex) on $[c, d]$ such that

$$h(c) = \mathbb{I}, \quad \sum_{j=1}^{j=k} \frac{D}{dt} (h_{ij} \zeta_j) \Big|_{(b,t)} = 0 \quad \forall i, t.$$

The second equation is equivalent to

$$\sum_{j=1}^{j=k} h'_{ij}(t) \zeta_j(b, t) + \sum_{j=1}^{j=k} \sum_{l=1}^{l=k} h_{ij}(t) A_{jl}(b, t) \zeta_l(b, t) = 0 \quad \iff \quad h' = -hA(b, \cdot). \quad (3.4)$$

Since (the real part of) the trace of (A_{ij}) is zero by (3.2), equation (3.4) has a unique solution in SO_k (or U_k) such that $h(c) = \mathbb{I}$. Furthermore, by (3.3)

$$|h(d) - \mathbb{I}| \leq \int_c^d |h'(t)| dt \leq \int_c^d |h| |A| dt \leq k^2 \int_c^d \int_a^b C_K |u_s| |u_t| ds dt. \quad (3.5)$$

Since $\Pi_{\partial\alpha} v_i = \sum_{j=1}^{j=k} h_{ij}(d) v_j$ by the above, the claim follows from equation (3.5). \square

Corollary 3.2. *If (M, g) is a Riemannian manifold and $(E, \langle \cdot, \cdot \rangle, \nabla)$ is a normed vector bundle with connection over M , for every compact subset $K \subset M$ there exists $C_K \in \mathbb{R}^+$ such that for every smooth closed curve $\alpha: [a, b] \longrightarrow M$ with $\text{Im } \alpha \subset K$*

$$|\Pi_\alpha - \mathbb{I}| \leq C_K \min(\|\text{d}\alpha\|_1, (b-a)\|\text{d}\alpha\|_2^2).$$

Proof. Let $\exp: TM \longrightarrow M$ be an exponential-like map. Since the group SO_k (or U_k if E is complex) is compact and

$$\|\text{d}\alpha\|_1^2 \leq (b-a)\|\text{d}\alpha\|_2^2$$

by Hölder's inequality, it is enough to assume that

$$\|\text{d}\alpha\|_1 \leq \min(r_{\text{exp}}^g(K)/2, 1).$$

Thus, there exists

$$\tilde{\alpha} \in C^\infty([a, b]; T_{\alpha(a)}M) \quad \text{s.t.} \quad \alpha(t) = \exp(\tilde{\alpha}(t)), \quad |\tilde{\alpha}(t)|_{\alpha(a)} < r_{\exp}(\alpha(a)).$$

Define

$$u: [0, 1] \times [a, b] \longrightarrow K \subset M \quad \text{by} \quad u(s, t) = \exp(s\tilde{\alpha}(t)).$$

Using

$$\begin{aligned} |\tilde{\alpha}(t)| &\leq C_K d_g(\alpha(a), \alpha(t)) \leq C_K \|\mathrm{d}\alpha\|_1, \\ |\tilde{\alpha}'(t)| &= |\{\mathrm{d}_{\tilde{\alpha}(t)} \exp\}^{-1}(\alpha'(t))| \leq C_K |\mathrm{d}_t \alpha|, \end{aligned}$$

we find that

$$u_s(s, t) = \{\mathrm{d}_{s\tilde{\alpha}(t)} \exp\}(\tilde{\alpha}(t)) \implies |u_s|_{(s,t)} \leq C'_K \|\mathrm{d}\alpha\|_1; \quad (3.6)$$

$$u_t(s, t) = s\{\mathrm{d}_{s\tilde{\alpha}(t)} \exp\}(\tilde{\alpha}'(t)) \implies |u_t|_{(s,t)} \leq C'_K |\mathrm{d}_t \alpha|. \quad (3.7)$$

Thus, by Lemma 3.1,

$$|\Pi_\alpha - \mathbb{I}| = |\Pi_{\partial u} - \mathbb{I}| \leq C_K \int_0^1 \int_a^b |u_s| |u_t| \mathrm{d}s \mathrm{d}t \leq C'_K \|\mathrm{d}\alpha\|_1^2 \leq C'_K (b-a) \|\mathrm{d}\alpha\|_2^2.$$

Since $\|\mathrm{d}\alpha\|_1 \leq r_{\exp}^g(K)$, it follows that $|\Pi_\alpha - \mathbb{I}| \leq C_K \|\mathrm{d}\alpha\|_1$. \square

Corollary 3.3. *If (M, g, \exp) is a Riemannian manifold with an exponential-like map and $(E, \langle \cdot, \cdot \rangle, \nabla)$ is a normed vector bundle with connection over M , for every compact subset $K \subset M$ there exists $C_K \in C^\infty(\mathbb{R}; \mathbb{R})$ such that for all $x \in K$ and smooth maps $\tilde{\alpha}: (-\epsilon, \epsilon) \longrightarrow T_x M$ and $\xi: (-\epsilon, \epsilon) \longrightarrow E_x$*

$$\left| \frac{D}{\mathrm{d}t} \left(\Pi_{\tilde{\alpha}(t)} \xi(t) \right) \Big|_{t=0} - \Pi_{\tilde{\alpha}(0)} \xi'(0) \right| \leq C_K (|\tilde{\alpha}(0)|) |\tilde{\alpha}(0)| |\tilde{\alpha}'(0)| |\xi(0)|. \quad (3.8)$$

Proof. Define

$$u: [0, 1] \times [0, \epsilon/2] \longrightarrow K \subset M \quad \text{by} \quad u(s, t) = \exp(s\tilde{\alpha}(t)).$$

Let $\{v_i\}$ be an orthonormal basis for E_x . Extend each v_i to

$$\zeta_i \in \Gamma(u|_{[0,1] \times t; E})$$

by parallel-transporting along the curves $s \longrightarrow u(s, t)$. If

$$\xi(t) = \sum_{i=1}^{i=k} f_i(t) v_i,$$

where k is the rank of E , then

$$\begin{aligned} \Pi_{\tilde{\alpha}(t)} \xi(t) &= \sum_{i=1}^{i=k} f_i(t) \zeta_i(1, t) \implies \\ \frac{D}{\mathrm{d}t} \left(\Pi_{\tilde{\alpha}(t)} \xi(t) \right) \Big|_{t=0} &= \sum_{i=1}^{i=k} f'_i(0) \zeta_i(1, 0) + \sum_{i=1}^{i=k} f_i(0) \frac{D}{\mathrm{d}t} \zeta_i(1, t) \Big|_{t=0} \\ &= \Pi_{\tilde{\alpha}(0)} \xi'(0) + \sum_{i=1}^{i=k} f_i(0) \frac{D}{\mathrm{d}t} \zeta_i(1, t) \Big|_{t=0}. \end{aligned} \quad (3.9)$$

On the other hand, by (3.1), (3.3), and the first identities in (3.6) and (3.7),

$$\begin{aligned} \left| \frac{D}{dt} \zeta_i(1, t) \right|_{t=0} &= \sum_{j=1}^{j=k} |A_{ij}(1, 0)| \leq k C'_K(|\tilde{\alpha}(0)|) \int_0^1 |u_s|_{(s,0)} |u_t|_{(s,0)} ds \\ &\leq C_K(|\tilde{\alpha}(0)|) |\tilde{\alpha}(0)| |\tilde{\alpha}'(0)|. \end{aligned} \quad (3.10)$$

The claim follows from (3.9) and (3.10). \square

Remark 3.4. Note that (3.3) is applied above with K replaced by the compact set

$$\exp(\{v \in T_x M : x \in K, |v| \leq |\tilde{\alpha}(0)|\}).$$

Thus, the constants $C'_K(|\tilde{\alpha}(0)|)$ and $C_K(|\tilde{\alpha}(0)|)$ may depend on $|\tilde{\alpha}(0)|$. If M is compact, then the first constant does not depend on $|\tilde{\alpha}(0)|$, since (3.3) can then be applied with $K = M$. The second constant is then also independent of K and $|\tilde{\alpha}(0)|$ if $\exp = \exp^\nabla$ for some connection ∇ in TM . So, in this case, the function C_K in (3.8) can be taken to be a constant independent of K .

3.2 Poincare lemmas

Lemma 3.5. *If $\zeta : S^1 \rightarrow \mathbb{R}^k$ is a smooth function such that $\int_0^{2\pi} \zeta(\theta) d\theta = 0$,*

$$\int_0^{2\pi} |\zeta(\theta)|^2 d\theta \leq \int_0^{2\pi} |\zeta'(\theta)|^2 d\theta.$$

Proof. Write

$$\zeta(\theta) = \sum_{n > -\infty}^{n < \infty} \zeta_n e^{in\theta};$$

see [6, Section 6.16]. Since ζ integrates to 0, $\zeta_0 = 0$. Thus,

$$\int_0^{2\pi} |\zeta(\theta)|^2 d\theta = 2\pi \sum_{n > -\infty}^{n < \infty} |\zeta_n|^2 \leq 2\pi \sum_{n > -\infty}^{n < \infty} |n\zeta_n|^2 = \int_0^{2\pi} |\zeta'(\theta)|^2 d\theta,$$

as claimed. \square

Proposition 3.6. *If (M, g) is a Riemannian manifold and $(E, \langle \cdot, \cdot \rangle, \nabla)$ is a normed vector bundle with connection over M , for every compact subset $K \subset M$ there exists $C_K \in \mathbb{R}^+$ with the following property. If $\alpha \in C^\infty(S^1; M)$ is such that $\text{Im } \alpha \subset K$ and $\xi, \zeta \in \Gamma(\alpha; E)$, then*

$$|\langle \nabla_\theta \xi, \zeta \rangle| \leq \|\nabla_\theta \xi\|_2 \|\nabla_\theta \zeta\|_2 + C_K \min(\|\text{d}\alpha\|_1, \|\text{d}\alpha\|_2^2) \|\xi\|_{2,1} \|\zeta\|_2,$$

where $\nabla_\theta \equiv \nabla_{\partial_\theta}^\alpha$ is the covariant derivative with respect to the oriented unit field on S^1 and all the norms are computed with respect to the standard metric on S^1 .

Proof. Identify $E_{\alpha(0)}$ with \mathbb{R}^k (or \mathbb{C}^k), preserving the metric. Denote by $\mathfrak{so}(E_{\alpha(0)}) \approx \mathfrak{so}_k$ (or $\mathfrak{u}(E_{\alpha(0)}) \approx \mathfrak{u}_k$) the Lie algebra of the Lie group $\text{SO}(E_{\alpha(0)}) \approx \text{SO}_k$ (or of $\text{U}(E_{\alpha(0)}) \approx \text{U}_k$). For each $\chi \in \mathfrak{so}(E_{\alpha(0)})$ (or $\chi \in \mathfrak{u}(E_{\alpha(0)})$), let $e^\chi \in \text{SO}(E_{\alpha(0)})$ (or $e^\chi \in \text{U}(E_{\alpha(0)})$) be the exponential of χ . Let

$$\Pi_\theta : E_{\alpha(0)} \rightarrow E_{\alpha(\theta)}$$

be the parallel transport along the curve $t \rightarrow \alpha(t)$ with $t \in [0, \theta]$. By Corollary 3.2, there exists $\chi \in so(E_{\alpha(0)})$ (or $\chi \in u(E_{\alpha(0)})$) such that

$$\Pi_{2\pi} = e^\chi \quad \text{and} \quad |\chi| \leq C_K \min(\|\mathrm{d}\alpha\|_1, \|\mathrm{d}\alpha\|_2^2). \quad (3.11)$$

By the first statement in (3.11),

$$\Psi: S^1 \times E_{\alpha(0)} \rightarrow \alpha^* E, \quad (\theta, v) \rightarrow e^{-\theta\chi/2\pi} \Pi_\theta(v),$$

is a smooth isometry. Let $\Phi_2 = \pi_2 \circ \Psi^{-1}: \alpha^* E \rightarrow E_{\alpha(0)}$ and

$$\bar{\zeta} = \frac{1}{2\pi} \int_0^{2\pi} \{\Phi_2 \zeta\}(\theta) \mathrm{d}\theta \in E_{\alpha(0)}.$$

By Hölder's inequality and Lemma 3.5,

$$\begin{aligned} |\langle \nabla_\theta \xi, \zeta - \Psi \bar{\zeta} \rangle| &\leq \|\nabla_\theta \xi\|_2 \|\zeta - \Psi \bar{\zeta}\|_2 \\ &= \|\nabla_\theta \xi\|_2 \|\Phi_2 \zeta - \bar{\zeta}\|_2 \leq \|\nabla_\theta \xi\|_2 \|\mathrm{d}(\Phi_2 \zeta)\|_2. \end{aligned} \quad (3.12)$$

By the product rule,

$$\begin{aligned} \|\mathrm{d}(\Phi_2 \zeta)\|_2 &\leq \|\mathrm{d}(\Pi^{-1} \zeta)\|_2 + |\chi/2\pi| \|\Pi^{-1} \zeta\|_2 = \|\nabla_\theta \zeta\|_2 + |\chi/2\pi| \|\zeta\|_2 \\ &\leq \|\nabla_\theta \zeta\|_2 + C_K \min(\|\mathrm{d}\alpha\|_1, \|\mathrm{d}\alpha\|_2^2) \|\zeta\|_2. \end{aligned} \quad (3.13)$$

On the other hand, by integration by parts, we obtain

$$\langle \nabla_\theta \xi, \zeta - \Psi \bar{\zeta} \rangle = \langle \nabla_\theta \xi, \zeta \rangle + \langle \xi, \nabla_\theta(\Psi \bar{\zeta}) \rangle. \quad (3.14)$$

Since $\Psi \bar{\zeta}$ is the parallel transport of $e^{\theta\chi/2\pi} \bar{\zeta}$,

$$\begin{aligned} |\langle \xi, \nabla_\theta(\Psi \bar{\zeta}) \rangle| &\leq \|\xi\|_2 \|\nabla_\theta(\Psi \bar{\zeta})\|_2 = \|\xi\|_2 |\chi/2\pi| \|\Psi \bar{\zeta}\|_2 \\ &\leq C_K \min(\|\mathrm{d}\alpha\|_1, \|\mathrm{d}\alpha\|_2^2) \|\xi\|_2 \|\zeta\|_2. \end{aligned} \quad (3.15)$$

The claim follows from equations (3.12)-(3.15). \square

Let $B_{R,r} \subset \mathbb{R}^2$ denote the open annulus with radii $r < R$ centered at the origin.

Corollary 3.7 (of Lemma 3.5). *There exists $C \in C^\infty(\mathbb{R}; \mathbb{R})$ such that for all $R \in \mathbb{R}^+$*

$$r \in (0, R], \quad \zeta \in C^\infty(B_{R,r}; \mathbb{R}^k), \quad \int_{B_{R,r}} \zeta = 0 \quad \implies \quad \|\zeta\|_1 \leq C(R/r) R^2 \|\mathrm{d}\zeta\|_2.$$

Proof. It is sufficient to assume that $k=1$. Define

$$\xi: S^1 \rightarrow \mathbb{R} \quad \text{by} \quad \xi(\theta) = \int_r^R \zeta(\rho, \theta) \rho \mathrm{d}\rho.$$

By Hölder's inequality and Lemma 3.5,

$$\begin{aligned} \left(\int_0^{2\pi} \left| \int_r^R \zeta(\rho, \theta) \rho \mathrm{d}\rho \right| \mathrm{d}\theta \right)^2 &\leq 2\pi \int_0^{2\pi} |\xi(\theta)|^2 \mathrm{d}\theta \leq 2\pi \int_0^{2\pi} |\xi'(\theta)|^2 \mathrm{d}\theta \\ &\leq 2\pi \int_0^{2\pi} \left(\int_r^R |\mathrm{d}_{(\rho,\theta)} \zeta| \rho^2 \mathrm{d}\rho \right)^2 \mathrm{d}\theta \\ &\leq \frac{\pi R^4}{2} \int_0^{2\pi} \int_r^R |\mathrm{d}_{(\rho,\theta)} \zeta|^2 \rho \mathrm{d}\rho \mathrm{d}\theta = \frac{\pi R^4}{2} \|\mathrm{d}\zeta\|_2^2. \end{aligned} \quad (3.16)$$

If the function $\rho \rightarrow \zeta(\rho, \theta)$ does not change sign on (r, R) , then

$$\int_r^R |\zeta(\rho, \theta)| \rho d\rho = \left| \int_r^R \zeta(\rho, \theta) \rho d\rho \right|.$$

On the other hand, if this function vanishes somewhere on (r, R) , then

$$|\zeta(\rho, \theta)| \leq \int_r^R |d_{(t,\theta)} \zeta| dt \quad \forall \rho \quad \implies \quad \int_r^R |\zeta(\rho, \theta)| \rho d\rho \leq \frac{R^2}{2} \int_r^R |d_{(t,\theta)} \zeta| dt.$$

Combining these two cases and using (3.16) and Hölder's inequality, we obtain

$$\begin{aligned} \int_0^{2\pi} \int_r^R |\zeta(\rho, \theta)| \rho d\rho d\theta &\leq \int_0^{2\pi} \left| \int_r^R \zeta(\rho, \theta) \rho d\rho \right| d\theta + \frac{R^2}{2} \int_0^{2\pi} \int_r^R |d_{(\rho,\theta)} \zeta| d\rho d\theta \\ &\leq \frac{\sqrt{\pi} R^2}{\sqrt{2}} \|d\zeta\|_2 + \frac{R^2}{2} \|d\zeta\|_2 \left(\int_0^{2\pi} \int_r^R \rho^{-1} d\rho d\theta \right)^{1/2} \\ &= \sqrt{\frac{\pi}{2}} \left(1 + \sqrt{\ln(R/r)} \right) R^2 \|d\zeta\|_2, \end{aligned} \tag{3.17}$$

as claimed. \square

Remark 3.8. By Corollary 4.7 below, C can in fact be chosen to be a constant function. Corollary 3.7 suffices for gluing J -holomorphic maps in symplectic topology, but Corollary 4.7 leads to a sharper version of Proposition 4.14; see Remark 4.13.

3.3 Exponential-like maps and differentiation

Let (M, g, \exp, ∇) be a smooth Riemannian manifold with an exponential-like map \exp and connection ∇ in TM , which is g -compatible, but not necessarily torsion-free. Let

$$T_{\nabla}(\xi(x), \zeta(x)) \equiv (\nabla_{\xi} \zeta - \nabla_{\zeta} \xi - [\xi, \zeta])|_x \quad \forall x \in M, \xi, \zeta \in \Gamma(M; TM),$$

be the torsion tensor of ∇ . If $\alpha: (-\epsilon, \epsilon) \rightarrow M$ is a smooth curve and $\xi \in \Gamma(\alpha; TM)$, put

$$\Phi_{\alpha(0)} \left(\alpha'(0); \xi(0), \frac{D}{ds} \xi \Big|_{s=0} \right) = \Pi_{\xi(0)}^{-1} \left(\frac{d}{ds} \exp(\xi(s)) \Big|_{s=0} \right) = \Pi_{\xi(0)}^{-1} (\{d_{\xi(0)} \exp\}(\xi'(0))),$$

where $\xi'(0) \in T_{\xi(0)}(TM)$ is the tangent vector to the curve $\xi: (-\epsilon, \epsilon) \rightarrow TM$ at $s=0$.

Lemma 3.9. *If (M, g, \exp, ∇) is a smooth Riemannian manifold with an exponential-like map and a g -compatible connection, there exists $C \in C^{\infty}(TM; \mathbb{R})$ such that*

$$\left| \Phi_x(v; w_0, w_1) - (v + w_1 - T_{\nabla}(v, w_0)) \right| \leq C(w_0) (|v| |w_0|^2 + |w_0| |w_1|)$$

for all $x \in M$ and $v, w_0, w_1 \in T_x M$.

Proof. Let $\alpha: (-\epsilon, \epsilon) \rightarrow M$ be a smooth curve and $\xi \in \Gamma(\alpha; TM)$ such that

$$\alpha(0) = x, \quad \alpha'(0) = v, \quad \xi(0) = w_0, \quad \frac{D}{ds} \xi(s) \Big|_{s=0} = w_1.$$

Put

$$\begin{aligned} F_{v,w_0,w_1}(t) &= \frac{d}{ds} \exp(t\xi(s)) \Big|_{s=0} = \{d_{tw_0} \exp\} (d_{w_0} m_t(\xi'(0))), \\ H_{v,w_0,w_1}(t) &= \Pi_{tw_0}(v + tw_1 - tT_{\nabla}(v, w_0)), \end{aligned}$$

where $m_t: TM \rightarrow TM$ is the scalar multiplication by t . Then,

$$\begin{aligned} F_{v,w_0,w_1}(0) &= \frac{d}{ds} \alpha(s) \Big|_{s=0} = v = H_{v,w_0,w_1}(0), \\ \frac{D}{dt} F_{v,w_0,w_1}(t) \Big|_{t=0} &= \frac{D}{ds} \frac{d}{dt} \exp(t\xi(s)) \Big|_{t=0} \Big|_{s=0} - T_{\nabla}(v, w_0) = w_1 - T_{\nabla}(v, w_0) = \frac{D}{dt} H_{v,w_0,w_1}(t) \Big|_{t=0}; \end{aligned}$$

see Corollary 3.3. Since

$$F_{v,w_0,\cdot}(t) - H_{v,w_0,\cdot}(t) \in \text{Hom}(T_x M \oplus T_x M, T_{\exp(tw_0)} M),$$

combining the last two equations, we obtain

$$|F_{v,w_0,w_1}(t) - H_{v,w_0,w_1}(t)| \leq C(w_0, t)t^2(|v| + |w_1|) \quad \forall v, w_0, w_1 \in T_x M, x \in M, t \in \mathbb{R},$$

where C is a smooth function on $TM \times \mathbb{R}$. Since

$$F_{v,w_0,w_1}(t) - H_{v,w_0,w_1}(t) = F_{v,tw_0,tw_1}(1) - H_{v,tw_0,tw_1}(1),$$

we conclude that there exists $C \in C^\infty(TM)$ such that

$$|F_{v,w_0,w_1}(1) - H_{v,w_0,w_1}(1)| \leq C(w_0)(|w_0|^2|v| + |w_0||w_1|) \quad \forall v, w_0, w_1 \in T_x M, x \in M, \quad (3.18)$$

as claimed. \square

For any $v, w_0, w_1 \in T_x M$, let $\tilde{\Phi}_x(v; w_0, w_1) = \Phi_x(v; w_0, w_1) - (v + w_1 - T_{\nabla}(v, w_0))$.

Corollary 3.10. *If (M, g, \exp, ∇) is a smooth Riemannian manifold with an exponential-like map and a g -compatible connection, there exists $C \in C^\infty(TM \times_M TM; \mathbb{R})$ such that*

$$\begin{aligned} & \left| \tilde{\Phi}_x(v; w_0, w_1) - \tilde{\Phi}_x(v; w'_0, w'_1) \right| \\ & \leq C(w_0, w'_0) \left((|w_0| + |w'_0|)|v| + |w_1| + |w'_1| \right) |w_0 - w'_0| + (|w_0| + |w'_0|)|w_1 - w'_1| \end{aligned}$$

for all $x \in M$ and $v, w_0, w_1, w'_0, w'_1 \in T_x M$.

Proof. By the proof of Lemma 3.9,

$$\tilde{\Phi}(v; w_0, w_1) = \tilde{\Phi}_1(w_0; v) + \tilde{\Phi}_2(w_0; w_1)$$

for some smooth bundle sections $\tilde{\Phi}_1, \tilde{\Phi}_2: TM \rightarrow \pi_{TM}^* \text{Hom}(TM, TM)$ such that

$$|\tilde{\Phi}_1(w_0; \cdot)| \leq C_1(w_0)|w_0|^2, \quad |\tilde{\Phi}_2(w_0; \cdot)| \leq C_2(w_0)|w_0| \quad \forall w_0 \in TM.$$

Thus,

$$\begin{aligned} |\tilde{\Phi}_1(w_0; \cdot) - \tilde{\Phi}_1(w'_0; \cdot)| &\leq C'_1(w_0, w'_0)(|w_0| + |w'_0|)|w_0 - w'_0| \\ |\tilde{\Phi}_2(w_0; \cdot) - \tilde{\Phi}_2(w'_0; \cdot)| &\leq C'_2(w_0, w'_0)|w_0 - w'_0| \end{aligned} \quad \forall w_0, w'_0 \in T_x M.$$

From the linearity of $\tilde{\Phi}_1(w_0; \cdot)$ and $\tilde{\Phi}_2(w_0; \cdot)$ in the second input, we conclude that

$$\begin{aligned} |\tilde{\Phi}_1(w_0; v) - \tilde{\Phi}_1(w'_0; v)| &\leq C'_1(w_0, w'_0)(|w_0| + |w'_0|)|w_0 - w'_0||v|, \\ |\tilde{\Phi}_2(w_0; w_1) - \tilde{\Phi}_2(w'_0; w'_1)| &\leq C'_2(w_0, w'_0)|w_0 - w'_0||w_1| + C_2(w'_0)|w'_0||w_1 - w'_1|. \end{aligned}$$

This establishes the claim. \square

3.4 Expansion of the $\bar{\partial}$ -operator

Let (M, J) and (Σ, j) be almost-complex manifolds. If $u: \Sigma \rightarrow M$ is a smooth map, let

$$\begin{aligned}\Gamma(u) &= \Gamma(\Sigma; u^*TM), & \Gamma_{J,j}^{0,1}(u) &= \Gamma(\Sigma; T^*\Sigma^{0,1} \otimes_{\mathbb{C}} u^*TM), \\ \bar{\partial}_{J,j}u &= \frac{1}{2}(du + J \circ du \circ j) \in \Gamma_{J,j}^{0,1}(u),\end{aligned}$$

as in (2.5). If ∇ is a connection in TM , define

$$D_{J,j;u}^{\nabla}: \Gamma(u) \rightarrow \Gamma_{J,j}^{0,1}(u) \quad \text{by} \quad D_{J,j;u}^{\nabla}\xi = \frac{1}{2}(\nabla^u\xi + J\nabla_j^u\xi) - \frac{1}{2}(T_{\nabla}(du, \xi) + JT_{\nabla}(du \circ j, \xi)).$$

If in addition $\exp: TM \rightarrow M$ is an exponential-like map and $\nabla J = 0$, define

$$\begin{aligned}\exp_u: \Gamma(u) &\rightarrow C^\infty(\Sigma; M), & \bar{\partial}_u, N_{\exp}^{\nabla}: \Gamma(u) &\rightarrow \Gamma_{J,j}^{0,1}(u) \quad \text{by} \\ \{\exp_u(\xi)\}_z &= \exp(\xi(z)) \quad \forall z \in \Sigma, & \{\bar{\partial}_u\xi\}_z(v) &= \Pi_{\xi(z)}^{-1}(\{\bar{\partial}_{J,j}(\exp_u(\xi))\}_z(v)) \quad \forall z \in \Sigma, v \in T_z\Sigma, \\ \bar{\partial}_u\xi &= \bar{\partial}_{J,j}u + D_{J,j;u}^{\nabla}\xi + N_{\exp}^{\nabla}(\xi).\end{aligned}$$

Lemma 3.11. *If (M, J, g, \exp, ∇) is an almost-complex Riemannian manifold with an exponential-like map and a g -compatible connection in (TM, J) , there exists $C \in C^\infty(TM \times_M TM; \mathbb{R})$ with the following property. If (Σ, j) is an almost complex manifold, $u: \Sigma \rightarrow M$ is a smooth map, and $\xi, \xi' \in \Gamma(u)$, then*

$$\begin{aligned}\left| \{N_{\exp}^{\nabla}(\xi)\}_z(v) - \{N_{\exp}^{\nabla}(\xi')\}_z(v) \right| &\leq C(\xi(z), \xi'(z)) \left((|\xi(z)| + |\xi'(z)|)(|\nabla_v(\xi - \xi')| + |\nabla_{jv}(\xi - \xi')|) \right. \\ &\quad \left. + (|\mathrm{d}_z u(v)| + |\mathrm{d}_z u(jv)|)(|\xi(z)| + |\xi'(z)|) + (|\nabla_v\xi| + |\nabla_{jv}\xi| + |\nabla_v\xi'| + |\nabla_{jv}\xi'|)|\xi(z) - \xi'(z)| \right)\end{aligned}$$

for all $z \in \Sigma, v \in T_z\Sigma$. Furthermore, $N_{\exp}^{\nabla}(0) = 0$.

Proof. Since the connection ∇ commutes with J , so does the parallel transport Π . Thus, with notation as in Section 3.3,

$$\{N_{\exp}^{\nabla}(\xi)\}_z(v) = \frac{1}{2} \left(\tilde{\Phi}(\mathrm{d}_z u(v); \xi(z), \nabla_v\xi) + J(u(z))\tilde{\Phi}(\mathrm{d}_z u(jv); \xi(z), \nabla_{jv}\xi) \right).$$

The claim now follows from Corollary 3.10. \square

Definition 3.12. *Let M be a smooth manifold and $(E, \langle \cdot, \cdot \rangle, \nabla)$ a normed vector bundle with connection over M . If $C_0 \in \mathbb{R}^+$, (Σ, j) is an almost complex manifold, and $u: \Sigma \rightarrow M$ is a smooth map, norms $\|\cdot\|_{p,1}$ and $\|\cdot\|_p$ on $\Gamma(u; E)$ and $\Gamma^1(u; E)$, respectively, are C_0 -admissible if for all $\xi \in \Gamma(u; E)$, $\eta \in \Gamma^1(u; E)$, and every continuous function $f: \Sigma \rightarrow \mathbb{R}$,*

$$\|f\eta\|_p \leq \|f\|_{C^0}\|\eta\|_p, \quad \|\eta \circ j\|_p = \|\eta\|_p, \quad \|\nabla^u\xi\|_p \leq \|\xi\|_{p,1}, \quad \|\xi\|_{C^0} \leq C_0\|\xi\|_{p,1}.$$

Proposition 3.13. *If (M, J, g, \exp, ∇) is an almost-complex Riemannian manifold with an exponential-like map and a g -compatible connection in (TM, J) , for every compact subset $K \subset M$ there exists $C_K \in C^\infty(\mathbb{R}; \mathbb{R})$ with the following property. If (Σ, j) is an almost complex manifold, $u: \Sigma \rightarrow K$ is a smooth map, and $\|\cdot\|_{p,1}$ and $\|\cdot\|_p$ are C_0 -admissible norms on $\Gamma(u; TM)$ and $\Gamma^1(u; TM)$, respectively, then*

$$\|N_{\exp}^{\nabla}(\xi) - N_{\exp}^{\nabla}(\xi')\|_p \leq C_K(C_0 + \|du\|_p + \|\xi\|_{p,1} + \|\xi'\|_{p,1})(\|\xi\|_{p,1} + \|\xi'\|_{p,1})\|\xi - \xi'\|_{p,1}$$

for all $\xi, \xi' \in \Gamma(u)$. Furthermore, $N_{\exp}^{\nabla}(0) = 0$. If the g -ball $B_{g;\delta}(u(z))$ of radius δ around $f(z)$ for some $z \in \Sigma$ is isomorphic to an open subset of \mathbb{C}^n and $|\xi(z)| < \delta$, then $\{N_{\exp}^{\nabla}\xi\}_z = 0$.

Proof. The first two statements follow from Lemma 3.11 and Definition 3.12. The last claim is clear from the definition of N_{exp}^{∇} . \square

Remark 3.14. As the notation suggests, one possibility for the norms $\|\cdot\|_{p,1}$ and $\|\cdot\|_p$ is the usual Sobolev L_1^p and L^p -norms with respect to some Riemannian metric on Σ , where $p > \dim_{\mathbb{R}}\Sigma$. Another natural possibility in the $\dim_{\mathbb{R}}\Sigma=2$ case is the modified Sobolev norms introduced in [3, Section 3]; these are particularly suited for gluing pseudo-holomorphic curves. By Proposition 4.10 below, in the $\dim_{\mathbb{R}}\Sigma=2$ case the constant C_0 itself is a function of $\|du\|_p$ only for either of these two choices of norms.

Remark 3.15. By Proposition 3.13, the operator $D_{J,j;u}^{\nabla}$ defined above is a linearization of the $\bar{\partial}$ -operator on the space of smooth maps to M at u . If ∇' is any connection in TM , the connection

$$\nabla: \Gamma(M; TM) \longrightarrow \Gamma(M; T^*M \otimes_{\mathbb{R}} TM), \quad \nabla_v \xi = \frac{1}{2} \left(\nabla'_v \xi - J \nabla'_v (J\xi) \right) \quad \forall v \in TM, \xi \in \Gamma(M; TM),$$

is J -compatible. If in addition ∇' and J are compatible with a Riemannian metric g on M , then so is ∇ . If ∇' is also the Levi-Civita connection of the metric g (i.e. $T_{\nabla'}=0$),

$$T_{\nabla}(v, w) = \frac{1}{2} (J(\nabla'_w J)v - J(\nabla'_v J)w) \quad \forall v, w \in T_x M, x \in M.$$

If the 2-form $\omega(\cdot, \cdot) \equiv g(J\cdot, \cdot)$ is closed as well, then

$$\nabla'_{Jv} J = -J \nabla'_v J \quad \forall v \in TM$$

by [4, (C.7.5)] and thus

$$T_{\nabla}(v, w) = -\frac{1}{4} (J(\nabla'_v J)w - J(\nabla'_w J)v - (\nabla'_{Jv} J)w + (\nabla'_{Jw} J)v) = -A_J(v, w) \quad \forall v, w \in T_x M, x \in M,$$

where A_J is the Nijenhuis tensor of J as in (2.27). The operator $D_{J,j;u}^{\nabla}$ then becomes

$$D_{J,j;u}^{\nabla}: \Gamma(u) \longrightarrow \Gamma_{J,j}^{0,1}(u), \quad D_{J,j;u}^{\nabla} \xi = \bar{\partial}_{\nabla^u} \xi + A_J(\partial_{J,j} u, \xi), \quad (3.19)$$

where

$$\begin{aligned} \bar{\partial}_{\nabla^u} \xi &= \frac{1}{2} (\nabla^u \xi + J \nabla_j^u \xi) \in \Gamma_{J,j}^{0,1}(u), \\ \partial_{J,j} u &= \frac{1}{2} (du - J \circ du \circ j) \in \Gamma(\Sigma; T^* \Sigma^{1,0} \otimes_{\mathbb{C}} u^* TM). \end{aligned}$$

This agrees with [4, (3.1.5)], since the Nijenhuis tensor of J is defined to be $-4A_J$ in [4, p18].

4 Sobolev and elliptic inequalities

This appendix refines, in the $n=2$ case, the proofs of Sobolev Embedding Theorems given in [5] to obtain a C^0 -estimate in Proposition 4.10 and elliptic estimates for the $\bar{\partial}$ -operator in Propositions 4.14 and 4.16. If $R, r \in \mathbb{R}$, let

$$B_R = \{x \in \mathbb{R}^2: |x| < R\}, \quad B_{R,r} = B_R - \bar{B}_r, \quad \tilde{B}_{R,r} = B_R - B_r.$$

4.1 Euclidian case

If ξ is an \mathbb{R}^k -valued function defined on a subset B of \mathbb{R}^2 , let $\text{supp}_{\mathbb{R}^2}(\xi)$ be the closure of $\text{supp}(\xi) \subset B$ in \mathbb{R}^2 . If U is an open subset of \mathbb{R}^2 , $\xi \in C^\infty(U; \mathbb{R}^k)$, and $p \geq 1$, let

$$\|\xi\|_p \equiv \left(\int_U |\xi|^p \right)^{1/p}, \quad \|\xi\|_{p,1} \equiv \|\xi\|_p + \|\text{d}\xi\|_p,$$

be the usual Sobolev norms of ξ .

Lemma 4.1. *For every bounded convex domain $\mathcal{D} \subset \mathbb{R}^2$, $\xi \in C^\infty(\mathcal{D}; \mathbb{R}^k)$, and $x \in \mathcal{D}$,*

$$|\xi_{\mathcal{D}} - \xi(x)| \leq \frac{2r_0^2}{|\mathcal{D}|} \int_{\mathcal{D}} |\text{d}_y \xi| |y-x|^{-1} \text{d}y,$$

where $2r_0$ is the diameter of \mathcal{D} , $|\mathcal{D}|$ is the area of \mathcal{D} , and

$$\xi_{\mathcal{D}} = \frac{1}{|\mathcal{D}|} \left(\int_{\mathcal{D}} \xi(y) \text{d}y \right)$$

is the average value of ξ on \mathcal{D} .

Proof. For any $y \in \mathcal{D}$,

$$\xi(y) - \xi(x) = \int_0^1 \frac{\text{d}}{\text{d}t} \xi(x+t(y-x)) \text{d}t = \int_0^1 \text{d}_{x+t(y-x)} \xi(y-x) \text{d}t.$$

Putting $g(z) = |\text{d}_z \xi|$ if $z \in \mathcal{D}$ and $g(z) = 0$ otherwise, we obtain

$$|\xi_{\mathcal{D}} - \xi(x)| \leq \frac{1}{|\mathcal{D}|} \int_{y \in \mathcal{D}} |\xi(y) - \xi(x)| \text{d}y \leq \frac{1}{|\mathcal{D}|} \int_{y \in \mathcal{D}} \int_0^\infty g(x+t(y-x)) |y-x| \text{d}t \text{d}y.$$

Rewriting the last integral in polar coordinates (r, θ) centered at x , we obtain

$$\begin{aligned} |\xi_{\mathcal{D}} - \xi(x)| &\leq \frac{1}{|\mathcal{D}|} \int_0^{2\pi} \int_0^{2r_0} \int_0^\infty g(tr, \theta) r^2 \text{d}t \text{d}r \text{d}\theta \\ &= \frac{1}{|\mathcal{D}|} \int_0^{2\pi} \int_0^{2r_0} \int_0^\infty g(t, \theta) r \text{d}t \text{d}r \text{d}\theta = \frac{2r_0^2}{|\mathcal{D}|} \int_0^{2\pi} \int_0^\infty g(t, \theta) \text{d}t \text{d}\theta \\ &= \frac{2r_0^2}{|\mathcal{D}|} \int_{\mathcal{D}} |\text{d}_y \xi| |y-x|^{-1} \text{d}y. \end{aligned}$$

This establishes the claim. □

Corollary 4.2. *For every $p > 2$, there exists $C_p > 0$ such that*

$$r \in [0, R/2], \quad \xi \in C^\infty(B_{R,r}; \mathbb{R}^k) \quad \implies \quad |\xi(x) - \xi(y)| \leq C_p R^{\frac{p-2}{p}} \|\text{d}\xi\|_p \quad \forall x, y \in B_{R,r}.$$

Proof. For any $x \in B_{R,r}$, put

$$\mathcal{D}_x = \{y \in B_{R,r} : \langle x, |x|y - rx \rangle > 0\}.$$




Figure 2: A convex region \mathcal{D}_x of the annulus $\mathcal{D}_{R,r}$ containing x

If $x \neq 0$, \mathcal{D}_x is the part of the annulus on the same side of the line $\langle x, y - rx/|x| \rangle = 0$ as x ; see Figure 2. In particular,

$$\text{diam}(\mathcal{D}_x) \leq 2R, \quad |\mathcal{D}_x| \geq \left(\frac{\pi}{3} - \frac{\sqrt{3}}{4}\right)R^2.$$

Thus, by Lemma 4.1 and Hölder's inequality,

$$\begin{aligned} |\xi(x) - \xi_{\mathcal{D}_x}| &\leq 12 \int_{y \in \mathcal{D}_x} |d_y \xi| |y-x|^{-1} dy \\ &\leq 12 \left(\int_{y \in B_{2R}(x)} |y-x|^{-\frac{p}{p-1}} \right)^{\frac{p-1}{p}} \|\text{d}\xi\|_p \leq C_p R^{\frac{p-2}{p}} \|\text{d}\xi\|_p, \end{aligned} \quad (4.1)$$

since $\frac{p}{p-1} < 2$. Let

$$x_{\pm} = (\pm(R-r)/2, 0), \quad y_{\pm} = (0, \pm(R-r)/2).$$

Since each of the convex regions $\mathcal{D}_{x_{\pm}}$ intersects $\mathcal{D}_{y_{\pm}}$ and \mathcal{D}_x intersects at least one (in fact precisely two if $r \neq 0$) of these four convex regions for every $x \in B_{R,r}$,

$$|\xi(x) - \xi(y)| \leq 8C_p R^{\frac{p-2}{p}} \|\text{d}\xi\|_p \quad \forall x, y \in B_{R,r}$$

by (4.1) and triangle inequality. □

Corollary 4.3. *For every $p > 2$, there exists $C_p \in C^\infty(\mathbb{R}^+; \mathbb{R})$ such that*

$$r \in [0, R/2], \quad \xi \in C^\infty(B_{R,r}; \mathbb{R}^k) \implies \|\xi\|_{C^0} \leq C_p(R) \|\xi\|_{p,1}.$$

Proof. By Corollary 4.2 and Hölder's inequality, for every $x \in B_{R,r}$

$$\begin{aligned} |\xi(x)| &\leq |\xi_{B_{R,r}}| + C_p R^{\frac{p-2}{p}} \|\text{d}\xi\|_p \leq \frac{1}{|B_{R,r}|} \|\xi\|_1 + C_p R^{\frac{p-2}{p}} \|\text{d}\xi\|_p \\ &\leq |B_{R,r}|^{-\frac{1}{p}} \|\xi\|_p + C_p R^{\frac{p-2}{p}} \|\text{d}\xi\|_p \leq (1+C_p) R^{-\frac{2}{p}} (\|\xi\|_p + R \|\text{d}\xi\|_p). \end{aligned} \quad (4.2)$$

This implies the claim. □

Lemma 4.4. For all $R > 0$ and $r \in [0, R)$,

$$\zeta \in C^\infty(B_{R,r}; \mathbb{R}^k), \quad \text{supp}_{\mathbb{R}^2}(\zeta) \subset \tilde{B}_{R,r} \implies \|\zeta\|_2 \leq \|d\zeta\|_1.$$

Proof. Such a function ζ can be viewed as a function on the complement of the ball B_r in \mathbb{R}^2 . Since ζ vanishes at infinity, for any $(x, y) \in B_{R,r}$

$$\zeta(x, y) = \begin{cases} \int_{-\infty}^x \zeta_s(s, y) ds, & \text{if } x \leq 0; \\ -\int_x^\infty \zeta_s(s, y) ds, & \text{if } x \geq 0; \end{cases} \quad \zeta(x, y) = \begin{cases} \int_{-\infty}^y \zeta_t(x, t) dt, & \text{if } y \leq 0; \\ -\int_y^\infty \zeta_t(x, t) dt, & \text{if } y \geq 0. \end{cases}$$

Taking the absolute value in these equations, we obtain

$$|\zeta(x, y)| \leq \int_{-\infty}^\infty |d_{(s,y)}\zeta| ds \quad \text{and} \quad |\zeta(x, y)| \leq \int_{-\infty}^\infty |d_{(x,t)}\zeta| dt, \quad (4.3)$$

where we formally set ζ and $d\zeta$ to be zero on the smaller disk. Multiplying the two inequalities in (4.3) and integrating with respect to x and y , we conclude

$$\int_{-\infty}^\infty \int_{-\infty}^\infty |\zeta(x, y)|^2 dx dy \leq \left(\int_{-\infty}^\infty \int_{-\infty}^\infty |d_{(x,y)}\zeta| dx dy \right)^2,$$

as claimed. \square

Corollary 4.5. For all $p, q \geq 1$ with $1 - 2/p \geq -2/q$, there exists $C_{p,q} \in \mathbb{R}^+$ such that

$$r \in [0, R), \quad \xi \in C^\infty(B_{R,r}; \mathbb{R}^k), \quad \text{supp}_{\mathbb{R}^2}(\xi) \subset \tilde{B}_{R,r} \implies \|\xi\|_q \leq C_{p,q} R^{1 - \frac{2}{p} + \frac{2}{q}} \|d\xi\|_p.$$

Proof. We can assume that $k = 1$. For $\epsilon > 0$, let $\zeta_\epsilon = (\xi^2 + \epsilon)^{\frac{q}{4}} - \epsilon^{\frac{q}{4}}$. By Lemma 4.4 and Hölder's inequality,

$$\begin{aligned} \|\xi\|_q^q &\leq \|\zeta_\epsilon + \epsilon^{\frac{q}{4}}\|_2^2 \leq 2\|d\zeta_\epsilon\|_1^2 + 2\epsilon^{\frac{q}{2}}\pi R^2 = 2\left\|\frac{q}{2}(\xi^2 + \epsilon)^{\frac{q}{4}-1}\xi d\xi\right\|_1^2 + 2\epsilon^{\frac{q}{2}}\pi R^2 \\ &\leq q^2\|(\xi^2 + \epsilon)^{\frac{q}{4}-\frac{1}{2}}d\xi\|_1^2 + 2\epsilon^{\frac{q}{2}}\pi R^2 \leq q^2\|d\xi\|_p^2\|(\xi^2 + \epsilon)^{\frac{q-2}{4}}\|_{\frac{p}{p-1}}^2 + 2\epsilon^{\frac{q}{2}}\pi R^2. \end{aligned} \quad (4.4)$$

Note that

$$1 - \frac{2}{p} = -\frac{2}{q} \implies \frac{q-2}{4} \frac{p}{p-1} = \frac{q-2}{4} \frac{2q}{q-2} = \frac{q}{2}.$$

Thus, letting ϵ go to zero in (4.4), we obtain

$$\|\xi\|_q^q \leq q^2\|d\xi\|_p^2\|\xi\|_q^{q-2} \implies \|\xi\|_q \leq q\|d\xi\|_p.$$

The case $1 - \frac{2}{p} > -\frac{2}{q}$ follows by Hölder's inequality. \square

Remark 4.6. By Hölder's inequality, the constant $C_{p,q}$ can be taken to be

$$C_{p,q} = \max(2, q)\pi^{\frac{1}{2}}\left(1 - \frac{2}{p} + \frac{2}{q}\right).$$

Corollary 4.7 (of Lemmas 4.1, 4.4). There exists $C > 0$ such that for all $R \in \mathbb{R}^+$

$$r \in [0, R], \quad \zeta \in C^\infty(B_{R,r}; \mathbb{R}^k), \quad \int_{B_{R,r}} \zeta = 0 \implies \|\zeta\|_1 \leq CR^2\|d\zeta\|_2.$$

Proof. (1) If $\zeta \in C^\infty(B_{R,r}; \mathbb{R}^k)$ integrates to 0 over its domain, then so does the function

$$\tilde{\zeta} \in C^\infty(B_{1,r/R}; \mathbb{R}^k), \quad \tilde{\zeta}(z) = \zeta(Rz).$$

Furthermore, $\|\tilde{\zeta}\|_1 = \|\zeta\|_1/R^2$ and $\|d\tilde{\zeta}\|_2 = \|d\zeta\|_2$. Thus, it is sufficient to prove the claim for $R=1$.

(2) If $r=0$, for some open half-disk $\mathcal{D} \subset B_{1,0}$

$$\int_{\mathcal{D}} \zeta = 0, \quad \|\zeta|_{\mathcal{D}}\|_1 \geq \frac{1}{2} \|\zeta\|_1. \quad (4.5)$$

By the first condition, Lemma 4.1, and Hölder's inequality

$$\|\zeta|_{\mathcal{D}}\|_1 \leq \frac{4}{\pi} \int_{\mathcal{D}} \int_{\mathcal{D}} |d_y \zeta| |y-x|^{-1} dy dx \leq 16 \int_{\mathcal{D}} |d_y \zeta| dy \leq 8\sqrt{2\pi} \|d\zeta\|_2.$$

Along with the second assumption in (4.5), this implies the claim for $r=0$ with $C=16\sqrt{2\pi}$.

(3) Let $\beta: \mathbb{R} \rightarrow [0, 1]$ be a smooth function such that

$$\beta(t) = \begin{cases} 1, & \text{if } t \leq 1/2; \\ 0, & \text{if } t \geq 1. \end{cases}$$

It remains to prove the claim for all $r > 0$ and $R=1$. By (3.17), we can assume that

$$r \leq \frac{1}{48\sqrt{3\pi}\|\beta'\|_{C^0}} < \frac{1}{96\sqrt{3\pi}}. \quad (4.6)$$

We first consider the case

$$\|\zeta|_{B_{2r,r}}\|_1 \geq \frac{1}{25} \|\zeta\|_1. \quad (4.7)$$

Using polar coordinates, define $\tilde{\zeta} \in C^\infty(B_{1,r}; \mathbb{R}^k)$ by

$$\tilde{\zeta}(\rho, \theta) = \beta(\rho)\zeta(\rho, \theta).$$

By Hölder's inequality and Lemma 4.4,

$$\|\zeta|_{B_{2r,r}}\|_1 \leq \sqrt{3\pi r} \|\tilde{\zeta}\|_2 \leq \sqrt{3\pi r} \|d\tilde{\zeta}\|_1 \leq \sqrt{3\pi r} (\|d\zeta\|_1 + \|\beta'\|_{C^0} \|\zeta|_{B_{1,1/2}}\|_1).$$

Along with the assumptions (4.6) and (4.7), this implies the bound with

$$C = 25 \frac{\sqrt{3\pi r}}{1 - 24\sqrt{3\pi}\|\beta'\|_{C^0} r} \leq \frac{25}{48}.$$

Finally, suppose

$$\|\zeta|_{B_{2r,r}}\|_1 \leq \frac{1}{25} \|\zeta\|_1. \quad (4.8)$$

Split the annulus $B_{1,r}$ into 3 wedges of equal area; split each wedge into a large convex outer portion and a small inner portion by drawing the line segment tangent to the circle of radius r and with the end points on the sides of the wedges $2r$ from the center as in Figure 3. By (4.8),

$$A \equiv \|\zeta|_{\mathcal{D}_+}\|_1 \geq \frac{8}{25} \|\zeta\|_1 \quad (4.9)$$




Figure 3: A large convex region \mathcal{D}_+ of an annulus \mathcal{D}

for the outer piece \mathcal{D}_+ of some wedge \mathcal{D} . If

$$\left| \int_{\mathcal{D}_+} \zeta \right| \leq \frac{3}{10}A,$$

then by Lemma 4.1, (4.6), and Hölder's inequality,

$$\begin{aligned} A &\leq \frac{3}{10}A + \frac{2 \left(\frac{\sqrt{3}}{2}\right)^2}{\frac{\pi}{3} \left(1 - \left(\frac{1}{96\sqrt{3}\pi}\right)^2\right)} \int_{\mathcal{D}_+} \int_{\mathcal{D}_+} |\mathrm{d}_y \zeta| |y-x|^{-1} \mathrm{d}y \mathrm{d}x \\ &\leq \frac{3}{10}A + \frac{9}{2\pi} \cdot \frac{7\sqrt{2}}{9} \cdot 2\pi\sqrt{3} \int_{\mathcal{D}} |\mathrm{d}_y \zeta| \mathrm{d}y \leq \frac{3}{10}A + 7\sqrt{2\pi} \|\mathrm{d}\zeta\|_2. \end{aligned}$$

Along with the assumption (4.9), this implies the bound with $C = 125\sqrt{2\pi}/4$. If

$$\left| \int_{\mathcal{D}_+} \zeta \right| \geq \frac{3}{10}A,$$

then by (4.8), (4.9), and (3.16),

$$\begin{aligned} A &\leq \|\xi|_{\mathcal{D}}\|_1 \leq \|\zeta|_{\mathcal{D}}\|_1 - \left| \int_{\mathcal{D}} \zeta \right| + \int_0^{2\pi} \left| \int_r^1 \zeta(\rho, \theta) \rho \mathrm{d}\rho \right| \mathrm{d}\theta \\ &\leq \left(A + \frac{1}{8}A\right) - \left(\frac{3}{10}A - \frac{1}{8}A\right) + \sqrt{\frac{\pi}{2}} \|\mathrm{d}\zeta\|_2 = \frac{19}{20}A + \sqrt{\frac{\pi}{2}} \|\mathrm{d}\zeta\|_2. \end{aligned}$$

Along with the assumption (4.9), this implies the bound with $C = 125\sqrt{2\pi}/4$. Since β can be chosen so that $\|\beta'\|_{C^0} < 3$ (actually arbitrarily close to 2), comparing with (3.17) for $R/r = 144\sqrt{3\pi}$ we conclude that the claim holds with $C = 125\sqrt{2\pi}/4$ for all r . \square

4.2 Bundle sections along smooth maps

Let (M, g) be a Riemannian manifold and $(E, \langle \cdot, \cdot \rangle, \nabla)$ a normed vector bundle with connection over M . If $u \in C^\infty(\tilde{B}_{R,r}; M)$, $\xi \in \Gamma(u; E)$, and $p \geq 1$, let

$$\|\xi\|_p \equiv \left(\int_{\tilde{B}_{R,r}} |\xi|^p \right)^{1/p}, \quad \|\xi\|_{p,1} \equiv \|\xi\|_p + \|\nabla^u \xi\|_p.$$

Lemma 4.8. *If (M, g) is a Riemannian manifold, $(E, \langle \cdot, \cdot \rangle, \nabla)$ is a normed vector bundle with connection over M , and $p, q \geq 1$ are such that $1 - 2/p \geq -2/q$, for every compact subset $K \subset M$ there exists $C_{K;p,q} \in \mathbb{R}^+$ with the following property. If $R \in \mathbb{R}^+$, $r \in [0, R)$, $u \in C^\infty(\tilde{B}_{R,r}; M)$ is such that $\text{Im } u \subset K$, and $\xi \in \Gamma_c(u; E)$, then*

$$\|\xi\|_q \leq C_{K;p,q} R^{1-\frac{2}{p}+\frac{2}{q}} (\|\nabla^u \xi\|_p + \|\xi \otimes du\|_p).$$

Proof. Let $\exp: TM \rightarrow M$ be an exponential-like map and $\{U_i: i \in [N]\}$ a finite open cover of K such that the g -diameter of each set U_i is at most $r_{\text{exp}}^g(K)/2$. Let $\{W_i: i \in [N]\}$ be an open cover of K such that $\overline{W_i} \subset U_i$. Choose smooth functions $\eta_i: M \rightarrow [0, 1]$ such that $\eta_i = 1$ on W_i and $\eta_i = 0$ outside of U_i . For each $i \in [N]$, pick $x_i \in W_i$. For each $z \in u^{-1}(U_i) \subset \tilde{B}_{R,r}$, define $\tilde{u}_i(z) \in T_{x_i} M$ and $\xi_i(z) \in E_{x_i}$ by

$$\exp_{x_i} \tilde{u}_i(z) = u(z), \quad |\tilde{u}_i(z)| < r_{\text{exp}}(x_i); \quad \Pi_{\tilde{u}_i(z)} \xi_i(z) = \xi(z).$$

For any $z \in B_{R,r}$, put $\tilde{\xi}_i(z) = \eta_i(u(z)) \xi_i(z)$. Since $\tilde{\xi}_i \in C_c^\infty(\tilde{B}_{R,r}; E_{x_i})$, by Corollary 4.5 there exists $C_{i;p,q} > 0$ such that

$$\|\xi|_{u^{-1}(W_i)}\|_q = \|\tilde{\xi}_i|_{u^{-1}(W_i)}\|_q \leq \|\tilde{\xi}_i\|_q \leq C_{i;p,q} R^{1-\frac{2}{p}+\frac{2}{q}} \|\text{d}\tilde{\xi}_i\|_p. \quad (4.10)$$

Since $\text{d}\tilde{\xi}_i = (\text{d}\eta_i \circ du)\xi_i + (\eta_i \circ u)\text{d}\xi_i$ on $u^{-1}(U_i)$ and vanishes outside of $u^{-1}(U_i)$,

$$\|\text{d}\tilde{\xi}_i\|_p \leq \|\text{d}\xi_i|_{u^{-1}(U_i)}\|_p + C_i \|\xi_i \otimes du\|_p. \quad (4.11)$$

On the other hand, by Corollary 3.3, if $u(z) \in U_i$

$$\left| \nabla^u \xi|_z - \Pi_{\tilde{u}_i(z)} \circ \text{d}_z \xi_i \right| \leq C_K |\text{d}_z u| |\xi(z)|. \quad (4.12)$$

Combining equations (4.10)-(4.12), we obtain

$$\|\xi|_{u^{-1}(W_i)}\|_q \leq \tilde{C}_{i;p,q} R^{1-\frac{2}{p}+\frac{2}{q}} (\|\xi\|_{p,1} + \|\xi \otimes du\|_p).$$

The claim follows by summing the last inequality over all i . □

Lemma 4.9. *If (M, g) is a Riemannian manifold, $(E, \langle \cdot, \cdot \rangle, \nabla)$ is a normed vector bundle with connection over M , and $p > 2$, for every compact subset $K \subset M$ there exists $C_{K;p} \in C^\infty(\mathbb{R}^+; \mathbb{R})$ with the following property. If $R \in \mathbb{R}^+$, $r \in [0, R/2]$, $u \in C^\infty(B_{R,r}; M)$ is such that $\text{Im } u \subset K$, and $\xi \in \Gamma(u; E)$, then*

$$\|\xi\|_{C^0} \leq C_{K;p}(R) (\|\xi\|_{p,1} + \|\xi \otimes du\|_p).$$

Proof. We continue with the setup in the proof of Lemma 4.8. By Corollary 4.3,

$$\|\xi|_{u^{-1}(W_i)}\|_{C^0} \leq \|\tilde{\xi}_i\|_{C^0} \leq C_{i;p}(R) \|\tilde{\xi}_i\|_{p,1} \leq C_{i;p}(R) (\|\xi|_{u^{-1}(U_i)}\|_p + \|\text{d}\tilde{\xi}_i\|_p).$$

As above, we obtain

$$\|\text{d}\tilde{\xi}_i\|_p \leq C_i (\|\nabla^u \xi\|_p + \|\xi \otimes du\|_p),$$

and the claim follows. □

Proposition 4.10. *If (M, g) is a Riemannian manifold, $(E, \langle \cdot, \cdot \rangle, \nabla)$ is a normed vector bundle with connection over M , and $p > 2$, for every compact subset $K \subset M$ there exists $C_{K;p} \in C^\infty(\mathbb{R}^+ \times \mathbb{R}; \mathbb{R})$ with the following property. If $R \in \mathbb{R}^+$, $r \in [0, R/2]$, $u \in C^\infty(B_{R,r}; M)$ is such that $\text{Im } u \subset K$, and $\xi \in \Gamma_c(u; E)$, then*

$$\|\xi\|_{C^0} \leq C_{K;p}(R, \|du\|_p) \|\xi\|_{p,1}.$$

The same statement holds if $B_{R,r}$ is replaced by a fixed compact Riemann surface (Σ, g_Σ) .

Proof. By Lemma 4.9 applied with $\tilde{p} = (p+2)/2$ and Hölder's inequality,

$$\|\xi\|_{C^0} \leq C_{K;\tilde{p}}(R) (\|\xi\|_{\tilde{p},1} + \|\xi \otimes du\|_{\tilde{p}}) \leq \tilde{C}_{K;\tilde{p}}(R) (\|\xi\|_{p,1} + \|du\|_p \|\xi\|_{q_1}), \quad (4.13)$$

where $q_1 = p(p+2)/(p-2)$. If $q_1 \leq p$, then the proof is complete. Otherwise, apply Lemma 4.8 with $p_1 = 2q_1/(q_1+2)$ and Hölder's inequality:

$$\|\xi\|_{q_1} \leq C_{K;p_1,q_1}(R) (\|\xi\|_{p_1,1} + \|\xi \otimes du\|_{p_1}) \leq C_{K;1}(R) (\|\xi\|_{p,1} + \|du\|_p \|\xi\|_{q_2}), \quad (4.14)$$

where $q_2 = pp_1/(p-p_1)$. If $q_2 \leq p$, then the claim follows from equations (4.13) and (4.14). Otherwise, we can continue and construct sequences $\{p_i\}, \{q_i\}, \{C_{K;i}\}$ such that

$$p_i = \frac{2q_i}{q_i + 2}, \quad q_{i+1} = \frac{pp_i}{p - p_i}; \quad (4.15)$$

$$\|\xi\|_{q_i} \leq C_{K;i}(R) (\|\xi\|_{p,1} + \|du\|_p \|\xi\|_{q_{i+1}}). \quad (4.16)$$

The recursion (4.15) implies that

$$q_{i+1} = \frac{2p}{2p + (p-2)q_i} q_i \implies \text{if } q_i > 0, \text{ then } 0 < q_{i+1} < q_i.$$

Thus, if $q_i > 2$ for all i , then the sequence $\{q_i\}$ must have a limit $q \geq 2$ with

$$q = \frac{2p}{2p + (p-2)q} q \implies (p-2)q = 0 \implies q = 0,$$

since $p > 2$ by assumption. Thus, $q_N \leq p$ for N sufficiently large and the first claim follows from (4.13) and the equations (4.16) with i running from 1 to N , where N is the smallest integer such that $q_{N+1} \leq p$. The second claim follows immediately from the first. \square

4.3 Elliptic estimates

If $A_1 = B_{R_1, r_1}$ and $A_2 = \bar{B}_{R_2, r_2}$ are two annuli in \mathbb{R}^2 , we write $A_2 \Subset_\delta A_1$ if $R_1 - R_2 > \delta$ and $r_2 - r_1 \geq \delta$.

Lemma 4.11. *For any $\delta > 0$, $p \geq 1$, and open annulus A_1 , there exists $C_{\delta,p}(A_1) > 0$ such that for any annulus $A_2 \Subset_\delta A_1$ and $\xi \in C^\infty(A_1; \mathbb{C}^k)$,*

$$\|\xi|_{A_2}\|_{p,1} \leq C_{\delta,p}(A_1) (\|\bar{\partial}\xi\|_p + \|d\xi\|_2 + \|\xi\|_1),$$

where the norms are taken with respect to the standard metric on \mathbb{R}^2 .

Proof. We can assume that A_2 is the maximal annulus such that $A_2 \Subset_\delta A_1$. Let $\eta: A_1 \rightarrow [0, 1]$ be a compactly supported smooth function such that $\eta|_{A_2} = 1$. By the fundamental elliptic inequality for the $\bar{\partial}$ -operator on S^2 [4, Lemma C.2.1],

$$\begin{aligned} \|\xi|_{A_2}\|_{p,1} &\leq \|\eta\xi\|_{p,1} \leq C_p(A_1)(\|\bar{\partial}(\eta\xi)\|_p + \|\eta\xi\|_p) \\ &\leq C_p(A_1)(\|\bar{\partial}\xi\|_p + \|(\mathrm{d}\eta)\xi\|_p + \|\eta\xi\|_p). \end{aligned} \quad (4.17)$$

By Corollary 4.5 with $(p, q) = (2, p)$ and $(p, q) = (1, 2)$ and Hölder's inequality,

$$\begin{aligned} \|\eta\xi\|_p &\leq C_p(A_1)\|\mathrm{d}(\eta\xi)\|_2 \leq C_p(A_1)(\|\mathrm{d}\xi\|_2 + \|(\mathrm{d}\eta)\xi\|_2) \\ &\leq \tilde{C}_p(A_1)(\|\mathrm{d}\xi\|_2 + \|\mathrm{d}((\mathrm{d}\eta)\xi)\|_1) \leq \tilde{C}_{p,\delta}(A_1)(\|\mathrm{d}\xi\|_2 + \|\mathrm{d}\xi\|_1 + \|\xi\|_1) \\ &\leq C_{\delta,p}(A_1)(\|\mathrm{d}\xi\|_2 + \|\xi\|_1). \end{aligned} \quad (4.18)$$

Similarly,

$$\|(\mathrm{d}\eta)\xi\|_p \leq C_{\delta,p}(A_1)(\|\mathrm{d}\xi\|_2 + \|\xi\|_1). \quad (4.19)$$

The claim follows by plugging (4.18) and (4.19) into (4.17). \square

Corollary 4.12. *For any $\delta > 0$, $p \geq 1$, and open annulus A_1 , there exists $C_{\delta,p}(A_1) > 0$ such that for any annulus $A_2 \Subset_\delta A_1$, and $\xi \in C^\infty(A_1; \mathbb{C}^n)$,*

$$\|\mathrm{d}\xi|_{A_2}\|_p \leq C_{\delta,p}(A_1)(\|\bar{\partial}\xi\|_p + \|\mathrm{d}\xi\|_2).$$

Proof. With $|A_1|$ denoting the area of A_1 , let

$$\bar{\xi} = \frac{1}{|A_1|} \int_{A_1} \xi$$

be the average value of ξ . By Lemma 4.11,

$$\begin{aligned} \|\mathrm{d}\xi|_{A_2}\|_p &= \|\mathrm{d}(\xi - \bar{\xi})|_{A_2}\|_p \leq C_{\delta,p}(A_1)(\|\bar{\partial}(\xi - \bar{\xi})\|_p + \|\mathrm{d}(\xi - \bar{\xi})\|_2 + \|\xi - \bar{\xi}\|_1) \\ &= C_{\delta,p}(A_1)(\|\bar{\partial}\xi\|_p + \|\mathrm{d}\xi\|_2 + \|\xi - \bar{\xi}\|_1). \end{aligned} \quad (4.20)$$

The claim follows by applying Corollary 4.7 with $\zeta = \xi - \bar{\xi}$. \square

Remark 4.13. The case $r_1 > 0$ (which is the case needed for gluing pseudo-holomorphic maps in symplectic topology) follows from Corollary 3.7; Corollary 4.7 can be used to obtain a sharper statement in this case (that $C_{\delta,p}(A_1)$ does not depend on r_1). The $r_1 = 0$ case requires only the first two steps in the proof of Corollary 4.7.

A smooth generalized CR-operator in a smooth complex vector bundle (E, ∇) with connection over an almost complex manifold (M, J) is an operator of the form

$$D = \bar{\partial}_\nabla + A: \Gamma(M; E) \rightarrow \Gamma(M; T^*M^{0,1} \otimes_{\mathbb{C}} E),$$

where

$$\bar{\partial}_\nabla \xi = \frac{1}{2}(\nabla \xi + \mathrm{i}\nabla_J \xi) \quad \forall \xi \in \Gamma(M; TM), \quad A \in \Gamma(M; \mathrm{Hom}(E; T^*M^{0,1} \otimes_{\mathbb{C}} E)).$$

If in addition $u: \Sigma \rightarrow M$ is a smooth map from an almost complex manifold (Σ, \mathfrak{j}) , the pull-back CR-operator is given by

$$D_u = \bar{\partial}_{\nabla^u} + A \circ \partial u: \Gamma(u; E) \rightarrow \Gamma^{0,1}(u; E).$$

Proposition 4.14. *If (M, g) is a Riemannian manifold with an almost complex structure J , $(E, \langle \cdot, \cdot \rangle, \nabla)$ is a normed complex vector bundle with connection over M and a smooth generalized CR-operator D , and $p \geq 1$, then for every compact subset $K \subset M$, $\delta > 0$, and open annulus $A_1 \subset \mathbb{R}^2$, there exists $C_{K; \delta, p}(A_1) \in \mathbb{R}^+$ with the following property. If $u \in C^\infty(A_1; M)$ is such that $\text{Im } u \subset K$, $\xi \in \Gamma(u; E)$, and $A_2 \Subset_\delta A_1$ is an annulus, then*

$$\|\nabla^u \xi|_{A_2}\|_p \leq C_{K; \delta, p}(A_1) (\|D_u \xi\|_p + \|\nabla^u \xi\|_2 + \|\xi \otimes du\|_p),$$

where the norms are taken with respect to the standard metric on \mathbb{R}^2 .

Proof. We continue with the setup in the proof of Lemma 4.8. By Corollary 4.12,

$$\begin{aligned} \|\text{d}\tilde{\xi}_i|_{A_2}\|_p &\leq C_{i; \delta, p}(A_1) (\|\bar{\partial}\tilde{\xi}_i\|_p + \|\text{d}\tilde{\xi}_i\|_2) \\ &\leq C'_{i; \delta, p}(A_1) (\|\bar{\partial}\tilde{\xi}_i|_{u^{-1}(U_i)}\|_p + \|\text{d}\tilde{\xi}_i|_{u^{-1}(U_i)}\|_2 + \|\xi \otimes du\|_p). \end{aligned} \quad (4.21)$$

Since ∇ commutes with the complex structure in E and $\tilde{\xi}_i = \xi_i$ on $u^{-1}(W_i)$, it follows from (4.12) and (4.21) that

$$\begin{aligned} \|\nabla^u \xi|_{A_2 \cap u^{-1}(W_i)}\|_p &\leq \|\text{d}\tilde{\xi}_i|_{A_2}\|_p + C_K \|\xi \otimes du\|_p \\ &\leq \tilde{C}_{i; \delta, p}(A_1) (\|\bar{\partial}\nabla^u \xi\|_p + \|\nabla^u \xi\|_2 + \|\xi \otimes du\|_p) \\ &\leq \tilde{C}'_{i; \delta, p}(A_1) (\|D_u \xi\|_p + \|\nabla^u \xi\|_2 + \|\xi \otimes du\|_p). \end{aligned} \quad (4.22)$$

The claim is obtained by summing the last equation over all i . □

Lemma 4.15. *If (M, g) is a Riemannian manifold with an almost complex structure J , $(E, \langle \cdot, \cdot \rangle, \nabla)$ is a normed complex vector bundle with connection over M and a smooth generalized CR-operator D , and $p > 2$, then for every compact subset $K \subset M$ and open ball $B \subset \mathbb{R}^2$, there exists $C_{K; B, p} \in C^\infty(\mathbb{R}; \mathbb{R})$ with the following property. If $u \in C^\infty(B; M)$ is such that $\text{Im } u \subset K$ and $\xi \in \Gamma_c(u; E)$, then*

$$\|\xi\|_{p,1} \leq C_{K; B, p} (\|du\|_p) (\|D_u \xi\|_p + \|\xi\|_p),$$

where the norms are taken with respect to the standard metric on \mathbb{R}^2 .

Proof. By an argument nearly identical to the proof of Proposition 4.14,

$$\|\xi\|_{p',1} \leq C_{K; p'}(B) (\|D_u \xi\|_{p'} + \|\xi\|_{p'} + \|\xi \otimes du\|_{p'})$$

for any $p' \geq 1$. On the other hand, by Proposition 4.10,

$$\|\xi\|_{C^0} \leq C_{K; B, \tilde{p}} (\|du\|_{\tilde{p}}) \|\xi\|_{\tilde{p},1},$$

where $\tilde{p} = (p+2)/2$. Proceeding as in the proof of Proposition 4.10, we then obtain

$$\begin{aligned} \|\xi\|_{p,1} &\leq C_{K; B, p} (\|du\|_{\tilde{p}}) (\|D_u \xi\|_p + \|\xi\|_p + \|du\|_p \|\xi\|_{\tilde{p},1}), \\ \|\xi\|_{\tilde{p},1} &\leq C_{K; \tilde{p}}(B) (\|D_u \xi\|_p + \|\xi\|_p + \|du\|_p \|\xi\|_{q_1}), \\ \|\xi\|_{q_i} &\leq C_{K; p_i, q_i}(B) (\|\xi\|_{p_i,1} + \|\xi \otimes du\|_{p_i}) \\ &\leq C_{K; B, i} (\|du\|_p) (\|D_u \xi\|_p + \|\xi\|_p + \|du\|_p \|\xi\|_{q_{i+1}}); \end{aligned}$$

we stop the recursion at the same value of $i = N$ as in the proof of Proposition 4.10. □

Proposition 4.16. *If (M, g) is a Riemannian manifold with an almost complex structure J , $(E, \langle \cdot, \cdot \rangle, \nabla)$ is a normed complex vector bundle with connection over M and a smooth generalized CR-operator D , and $p > 2$, then for every compact subset $K \subset M$ and compact Riemann surface (Σ, g_Σ) , there exists $C_{K; \Sigma, p} \in C^\infty(\mathbb{R}; \mathbb{R})$ with the following property. If $u \in C^\infty(\Sigma; M)$ is such that $\text{Im } u \subset K$ and $\xi \in \Gamma(u; E)$, then*

$$\|\xi\|_{p,1} \leq C_{K; \Sigma, p} (\|du\|_p) (\|D_u \xi\|_p + \|\xi\|_p).$$

Proof. This statement is immediate from Lemma 4.15. □

Department of Mathematics, SUNY Stony Brook, NY 11794-3651
azinge@math.sunysb.edu

References

- [1] I. Chavel, *Riemannian Geometry: A Modern Introduction*, Cambridge University Press, 1996.
- [2] A. Floer, *The unregularized gradient flow of the symplectic action*, Comm. Pure Appl. Math XLI (1988), 775-813.
- [3] J. Li and G. Tian, *Virtual moduli cycles and Gromov-Witten invariants of general symplectic manifolds*, *Topics in Symplectic 4-Manifolds*, Internat. Press, 1998.
- [4] D. McDuff and D. Salamon, *J-Holomorphic Curves and Symplectic Topology*, AMS Colloquium Publ. 52, 2004.
- [5] T. Mrowka, *18.966 Lecture Notes*, Spring 1998.
- [6] F. Warner, *Foundations of Differentiable Manifolds and Lie Groups*, Springer 1983.