

PROBLEM SET 6 - Product measures

1. Suppose $E \subset \mathbb{R}^2$ is such that $E_y = \{x : (x, y) \in E\} \subset \mathbb{R}$ has measure zero for almost every y . Show that E has measure zero.
2. If f and g are measurable on \mathbb{R} , show that $h(x, y) = f(x)g(y)$ is measurable on \mathbb{R}^2 .
3. Show

$$\int_{\mathbb{R}^n} e^{-|x|^2} dm = \pi^{n/2}.$$

Hint: For $n = 1$ use

$$\left(\int e^{-x^2} dx\right)^2 = \int \int e^{-x^2-y^2} dx dy,$$

and polar coordinates. For $n > 1$, use $|x|^2 = x_1^2 + \dots + x_n^2$ and Fubini's theorem to reduce to the $n = 1$ case.

4. The convolution $f * g$ of two functions $f, g \in L^1(\mathbb{R})$ is defined as the function

$$f * g(y) = \int_{\mathbb{R}} f(y-x)g(x)dx.$$

Show that $f * g$ is in $L^1(\mathbb{R})$ and

$$\int f * g(y)dy = \left(\int_{\mathbb{R}} f(x)dx\right)\left(\int_{\mathbb{R}} g(x)dx\right).$$

5. If $f \in L^1(\mathbb{R})$ define its Fourier transform as

$$\hat{f}(t) = \int_{\mathbb{R}} f(x)e^{-ixt}dx.$$

Show that

$$\widehat{f * g}(t) = \hat{f}(t)\hat{g}(t).$$