

ERRATUM: SYMPLECTIC INVOLUTIONS OF $K3^{[n]}$ TYPE AND KUMMER N TYPE MANIFOLDS

LJUDMILA KAMENOVA, GIOVANNI MONGARDI, AND ALEXEI OBLOMKOV

ABSTRACT. In this note we present a corrected formula for the enumeration of connected components of the locus fixed by a symplectic involution inside hyperkähler manifolds of types $K3^{[n]}$ and generalized Kummer. We also provide further precisions concerning the involutions considered in the Kummer case.

In the [3] the enumerative part of Theorems 1.2 and 1.3 is incorrect. The source of error is an erroneous interpretation of the results of [1] in the proof of Lemma 3.13. The error does not affect the enumeration of the top dimensional components as in Theorem 3.7 and 3.9. Moreover, the result of Theorem 1.3 does not apply to all involutions with a non trivial action on H^3 , and we detail here the correct type of involutions considered.

The lower-dimensional component counts are special cases of Theorems 3.0.3 and 3.0.7 from [4]. To state the results let introduce the standard theta function:

$$\vartheta(\eta; q) = \sum_{n \in \mathbb{Z}} q^{n^2} \eta^n.$$

Theorem 1. *Let S be a $K3$ surface and let ι be a symplectic involution of S . Let $S^{[n]}$ be the Hilbert scheme of n points on S and let us consider the induced action of ι on $S^{[n]}$. Then all the irreducible components of the locus stabilized by ι are deformation equivalent to Hilbert schemes of points on a $K3$ surface or isolated points, and their number for each dimension $2k$ is*

$$N_k = \Theta(n - 2k),$$

where $\Theta(j)$ is the j -th coefficient of the Theta series

$$\Theta(q) = \sum_{i \geq 0} \Theta(i) q^i = \vartheta(q; q^2)^8.$$

Remark 1. All involutions on generalized Kummer manifolds act trivially on the second cohomology, and they can be subdivided in three different geometrical categories:

- Involutions obtained by a translation of a two torsion point.
- Involutions obtained by a sign change composed with a translation by a two torsion point.
- Involutions obtained by a composition of a translation by a point of order four with a sign change.

The first kind of involutions only appears when n is odd, while the third kind only when $n \equiv 3$ modulo 4.

In this paper we deal with the second kind of involutions, and we call them *regular involutions*. The fixed locus of the first kind of involutions was analyzed in [5, Lemma 3.5], while the third kind was analyzed only in the case of Kummer sixfolds in [2, Lemma 2.4]. It would be interesting to compute the fixed locus in general for an involution of the third kind.

Theorem 2. *Let $X = K_n(A)$ be a n -Kummer hyperkähler manifold and let $\iota \in \text{Aut}(A)$ be a regular symplectic involution of the abelian surface A . Then all irreducible components of X^ι are of $K3^{[k]}$ type, and their number is*

$$N_k = \Theta(n - 2k; 1),$$

where $\Theta(j; 1)$ is the j -th coefficient of the Theta series

$$\Theta(q) = \sum_{i \geq 0, \gamma \in A, \gamma^2 = 1} \Theta(i; \gamma) q^i \gamma = \prod_{\gamma \in A, \gamma^2 = 1} \vartheta(\gamma \cdot q; q^2).$$

Remark 2. All three kinds of involutions of Remark 1 can be recognized by their action on higher cohomology: involutions obtained by pure translations have a trivial action on H^3 , and involutions of the third kind, those obtained by composing an order four translation with -1 , are recognizable by their action on H^6 . Indeed, if $n \equiv 3$ modulo 4, in H^6 there are 256 distinguished classes of codimension three subvarieties of a generalized Kummer. These subvarieties are in correspondance with points in $A[4]$, each of them is the locus of subschemes having support of multiplicity at least four on a given four torsion point of A . Regular involutions fix 16 of these subvarieties and permute the rest, while involutions of the third kind freely permute these subvarieties.

ACKNOWLEDGEMENTS

We are grateful to Mirko Mauri for pointing out the inaccuracy in Theorem 2. L.K. is partially supported by a grant from the Simons Foundation/SFARI (522730, LK). G.M. is supported by PRIN2020 research grant "2020KKWT53" and by PRIN2022 research grant "2022PEKYBJ", and is a member of INdAM GNSAGA. A.O. is partially supported by NSF grant DMS-2200798 and NSF FGR grant DMS-1760373.

REFERENCES

- [1] W. Crawley-Boevey, Geometry of the moment map for representations of quivers. *Compositio Mathematica*, 126(3):257–293, 2001.
- [2] S. Floccari, Sixfolds of generalized Kummer type and K3 surfaces. arXiv:2210.02948 [math.AG] (2022).
- [3] L. Kamenova, G. Mongardi, A. Oblomkov, Symplectic involutions of $K3^{[n]}$ type and Kummer n type manifolds. *Bull. of London Math. Soc.* 54 (2022) 894–909.
- [4] L. Kamenova, G. Mongardi, A. Oblomkov, Fixed loci of symplectic automorphisms of $K3^{[n]}$ and n -Kummer type manifolds, arXiv:2308.14692 [math.AG] (2023).
- [5] K. Oguiso, No cohomologically trivial non-trivial automorphism of generalized Kummer manifolds *Nagoya Math. J.* 1-13. doi:10.1017/nmj.2018.29.

DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, MATH TOWER 3-115, STONY BROOK, NY 11794-3651, USA

Email address: kamenova@math.stonybrook.edu

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI BOLOGNA, PIAZZA DI PORTA SAN DONATO 5, BOLOGNA, ITALIA 40126

Email address: giovanni.mongardi2@unibo.it

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MASSACHUSETTS, LEDERLE GRADUATE TOWER, AMHERST, MA 01003, USA

Email address: oblomkov@math.umass.edu