

On the Bartnik extension problem for the static vacuum Einstein equations

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2013 Class. Quantum Grav. 30 125005

(<http://iopscience.iop.org/0264-9381/30/12/125005>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 24.46.189.47

The article was downloaded on 19/05/2013 at 07:15

Please note that [terms and conditions apply](#).

On the Bartnik extension problem for the static vacuum Einstein equations

Michael T Anderson and Marcus A Khuri

Department of Mathematics, Stony Brook University, Stony Brook, NY 11790, USA

E-mail: anderson@math.sunysb.edu and khuri@math.sunysb.edu

Received 6 December 2012, in final form 14 March 2013

Published 14 May 2013

Online at stacks.iop.org/CQG/30/125005

Abstract

We develop a framework for understanding the existence of asymptotically flat solutions to the static vacuum Einstein equations on $M = \mathbb{R}^3 \setminus B$ with geometric boundary conditions on $\partial M \simeq S^2$. A partial existence result is obtained, giving a partial resolution of a conjecture of Bartnik on such static vacuum extensions. The existence and uniqueness of such extensions is closely related to Bartnik's definition of quasi-local mass.

PACS numbers: 04.20.-q, 04:20.Cv, 02.40.Vh, 02.30.Jr

1. Introduction

This paper is concerned with a conjecture of Bartnik [B3, B4] on the existence and uniqueness of static solutions to the vacuum Einstein equations with certain prescribed boundary data. On the physical side, this is closely related to the issue of local mass in general relativity while, on the mathematical side, to the issue of global existence and uniqueness for a rather complicated geometric nonlinear system of elliptic boundary value problems.

Let M be a 3-manifold diffeomorphic to $\mathbb{R}^3 \setminus B$ where B is a 3-ball, so that $\partial M \simeq S^2$. The static vacuum Einstein equations are the equations for a pair (g, u) consisting of a smooth Riemannian metric g on M and a positive potential function $u : M \rightarrow \mathbb{R}^+$ given by

$$u \operatorname{Ric}_g = D^2 u, \quad \Delta u = 0, \quad (1.1)$$

where the Hessian D^2 and Laplacian $\Delta = \operatorname{tr} D^2$ are taken with respect to g . The equations (1.1) are equivalent to the statement that the four-dimensional metric

$$g_{\mathcal{M}} = \pm u^2 dt^2 + g, \quad (1.2)$$

on the 4-manifold $\mathcal{M} = \mathbb{R} \times M$ is Ricci-flat, i.e.

$$\operatorname{Ric}_{g_{\mathcal{M}}} = 0. \quad (1.3)$$

This holds for either choice of sign in (1.2) and since most of the analysis of the paper concerns the Riemannian data (g, u) in (1.1), we will assume $g_{\mathcal{M}}$ is Riemannian, and moreover identify

t in (1.2) periodically, to obtain a metric on $\mathcal{M} = S^1 \times M$ with t replaced by the angular variable θ .

Given (M, g, u) as above, let γ be the Riemannian metric induced on $S^2 = \partial M$ and let H be the mean curvature of $\partial M \subset (M, g)$, (with respect to the inward unit normal into M). Then (one version of) the Bartnik conjecture [B4] states that, given an arbitrary such pair in C^∞ ,

$$(\gamma, H) \in \text{Met}^\infty(S^2) \times C_+^\infty(S^2), \quad H > 0, \tag{1.4}$$

there exists a unique asymptotically flat solution (g, u) to the static vacuum Einstein equations (1.1) inducing the boundary data (γ, H) on ∂M .

This conjecture is a natural outgrowth of Bartnik’s concept of quasi-local mass $m_B(\Omega)$, [B2, B3], defined as follows. Let (Ω, g) be a smooth compact 3-manifold with smooth boundary of non-negative scalar curvature, and define an *admissible extension* of (Ω, g) to be a complete, asymptotically flat 3-manifold (\tilde{M}, g) of non-negative scalar curvature in which (Ω, g) embeds isometrically and is not enclosed by any compact minimal surfaces (horizons). Then

$$m_B(\Omega) = \inf\{m_{\text{ADM}}(\tilde{M}) : (\tilde{M}, g) \text{ is an admissible extension of } (\Omega, g)\}, \tag{1.5}$$

where $m_{\text{ADM}}(\tilde{M})$ is the ADM mass of (\tilde{M}, g) , cf [B1]. In [HI] Huisken and Ilmanen have proved a number of basic properties of $m_B(\Omega)$, in particular that $m_B(\Omega) > 0$ unless (Ω, g) is locally isometric to Euclidean space. In [Br] Bray discusses a similar definition, where the boundary $\partial\Omega$ is required to be outer-minimizing in (\tilde{M}, g) . As will be seen below, the outer-minimizing property plays an important role in this paper, although for somewhat different reasons than in [Br].

Conjecturally, an extension (\tilde{M}, g) realizing the infimum in (1.5) is a solution to the static vacuum Einstein equations (1.1) on $M = \tilde{M} \setminus \Omega$ which is Lipschitz, (but not smooth), across the junction $\partial\Omega$ and for which the induced metric and mean curvature at the boundary of the interior and exterior regions agree:

$$g|_{\partial M} = g|_{\partial\Omega}, \quad H_{\partial M} = H_{\partial\Omega},$$

leading to the boundary data (1.4). Observe that the boundary data (γ, H) have the character of a mixed Dirichlet–Neumann type boundary value problem for the static equations (1.1), but the potential function u is absent from the boundary data. We point out that more standard Dirichlet or Neumann boundary data are not suitable for the (static) Einstein equations, cf [A3].

In this paper, we develop a general framework for the Bartnik conjecture and make partial progress on its resolution. To describe the setting, let $\mathcal{E}_S = \mathcal{E}_S^{m,\alpha}$ be the moduli space of AF static vacuum solutions (M, g, u) on a given 3-manifold M which are $C^{m,\alpha}$ up to ∂M , $m \geq 3$. The exact definition is given in section 2, but basically \mathcal{E}_S is the space of all AF static vacuum metrics on M modulo the action of the group Diff_1 of diffeomorphisms on M equal to the identity on ∂M . Next, let $\text{Met}^{m,\alpha}(\partial M)$ be the space of $C^{m,\alpha}$ metrics on $\partial M \simeq S^2$ and $C^{m-1,\alpha}(\partial M)$ be the space of $C^{m-1,\alpha}$ functions on ∂M . One thus has a natural map, mapping a static vacuum solution to its Bartnik boundary data:

$$\begin{aligned} \Pi_B : \mathcal{E}_S^{m,\alpha} &\rightarrow \text{Met}^{m,\alpha}(\partial M) \times C^{m-1,\alpha}(\partial M), \\ \Pi_B(g) &= (\gamma, H). \end{aligned} \tag{1.6}$$

Theorem 1.1. *The space $\mathcal{E}_S^{m,\alpha}$ is a smooth (infinite dimensional) Banach manifold, and the map Π_B is C^∞ smooth and Fredholm, of Fredholm index 0.*

Theorem 1.1 essentially amounts to the statement that the static vacuum Einstein equations (1.1) with boundary conditions (1.4) form an elliptic boundary value problem, modulo gauge transformations, i.e. diffeomorphisms, and that one has a well-behaved local

existence theory for this problem. We note that this boundary value problem also has a variational characterization, cf proposition 3.7.

Let \mathcal{E}^+ be the open submanifold of \mathcal{E}_S for which the mean curvature H is positive, i.e.

$$\mathcal{E}^+ = (\Pi_B)^{-1}(\text{Met}^{m,\alpha}(\partial M) \times C_+^{m-1,\alpha}(\partial M)).$$

The Bartnik conjecture above may thus be rephrased to state that the smooth map Π_B , restricted to the open submanifold \mathcal{E}^+ ,

$$\Pi_B : \mathcal{E}^+ \rightarrow \text{Met}^{m,\alpha}(\partial M) \times C_+^{m-1,\alpha}(\partial M), \tag{1.7}$$

is surjective and injective, and hence, via the inverse function theorem, is a smooth diffeomorphism.

However, this most optimistic version of the conjecture does not hold, in that Π_B in (1.7) cannot be a diffeomorphism. To illustrate the problem consider (for example) the flat solution g_{flat} with $u = 1$; there are boundaries $\partial M = S^2 \subset \mathbb{R}^3$ given by embedded spheres (S^2, γ_i, H_i) with H_i uniformly positive, which converge smoothly in \mathbb{R}^3 to a limit which is an immersed but not embedded sphere in \mathbb{R}^3 . Such a limit is then at the boundary $\partial \mathcal{E}^+$, but the limit boundary data $(\gamma, H) \in \text{Met}^{m,\alpha}(\partial M) \times C_+^{m-1,\alpha}(\partial M)$. In other words, the condition that the boundary data (γ, H) is uniformly controlled in the target space is not sufficient to ensure that one stays within the class of manifolds-with-boundary. (In particular, there cannot be a smooth inverse map to Π_B .) As a concrete example, let T^2 be a torus of revolution embedded in \mathbb{R}^3 with $H > 0$. One may remove a (small) essential annulus from T^2 and smoothly attach two embedded discs to obtain a 2-sphere S^2 with $H > 0$, cf figure 1. This surface may be deformed to obtain a curve $(S^2)_t, t \in [0, 1]$, of positive mean curvature spheres which for $t < \frac{1}{2}$ are embedded and for $t \geq \frac{1}{2}$ are immersed, with a single self-intersection point of the discs at $t = \frac{1}{2}$. (The same situation holds with any background static vacuum solution and varying boundary ∂M within M .) This passage from embedded to immersed behavior also shows that the boundary map Π_B on \mathcal{E}^+ is not proper.




Figure 1. An illustration of the one-parameter family of spheres $(S^2)_t, t \in [0, 1]$, in the process of passing from embedding to immersion.

The basic issue is in fact that of finding domains within \mathcal{E}^+ on which Π_B is proper. Recall that a map between two topological spaces is proper, if the preimage of any compact set is itself compact. In the current setting, Π_B is proper on a domain $\mathcal{U} \subset \mathcal{E}^+$ if whenever (γ_i, H_i) is a sequence of boundary data converging to limit data (γ, H) and $(M, g_i, u_i) \in \mathcal{U}$ are any solutions with $\Pi_B(g_i, u_i) = (\gamma_i, H_i)$, then (M, g_i, u_i) converges, in a subsequence, to a limit solution $(M, g, u) \in \mathcal{U}$. Here convergence is in the topology of the target and domain spaces respectively. In other words, control of the boundary data $(\gamma, H) = \Pi_B(g, u)$ implies

global control of the solution (M, g, u) within \mathcal{U} . Equivalently, one needs *a priori* estimates controlling the behavior of the full solution (M, g, u) in terms of the boundary data (γ, H) .

Now let $\mathcal{E}^o \subset \mathcal{E}^+$ be the domain for which the boundary ∂M is strictly *outer-minimizing*, i.e. for which

$$\text{area}(\Sigma) > \text{area}(\partial M), \tag{1.8}$$

for any surface $\Sigma \subset M$ homologous to ∂M with $\Sigma \neq \partial M$ (we point out that the examples in figure 1 are *not* strictly outer-minimizing for t sufficiently close to $\frac{1}{2}$). Clearly \mathcal{E}^o is an open submanifold of \mathcal{E}^+ . It is not known (although likely to be true) that \mathcal{E}^o is connected. Throughout the following, we thus assume that \mathcal{E}^o is taken to be the connected component containing the standard flat exterior solution where M is the exterior of the standard unit ball in \mathbb{R}^3 , with ∂M the round S^2 of radius 1. One then has a natural boundary map

$$\Pi^o : \mathcal{E}^o \rightarrow \text{Met}^{m,\alpha}(\partial M) \times C_+^{m-1,\alpha}(\partial M). \tag{1.9}$$

The second main result of this paper is the following.

Theorem 1.2. *The boundary map Π^o in (1.9) is ‘almost’ proper, in the following sense. If (γ_i, H_i) is a sequence of boundary data converging to limit data (γ, H) in $\text{Met}^{m,\alpha}(\partial M) \times C_+^{m-1,\alpha}(\partial M)$, and $(M, g_i, u_i) \in \mathcal{E}^o$ are any solutions with $\Pi^o(g_i, u_i) = (\gamma_i, H_i)$, then (M, g_i, u_i) converges in $\mathcal{E}_S^{m,\alpha}$, in a subsequence, to a limit solution $(M, g, u) \in \mathcal{E}_S^{m,\alpha}$ which is (at least) weakly outer-minimizing, i.e.*

$$\text{area}(\Sigma) \geq \text{area}(\partial M), \tag{1.10}$$

for Σ as in (1.8).

Roughly speaking, theorem 1.3 thus shows that static vacuum solutions (M, g, u) with outer-minimizing boundary are controlled by their boundary data (γ, H) . The issue remains however of how to determine from the boundary data (γ, H) whether the boundary is outer-minimizing in (M, g, u) . We point out that the full global property (1.8) is not actually necessary; theorem 1.2 remains valid if (1.8) holds only in an arbitrarily small neighborhood of ∂M , (depending on (M, g, u)), cf remark 4.4.

A smooth proper Fredholm map $F : B_1 \rightarrow B_2$ between connected Banach manifolds has a \mathbb{Z}_2 -valued degree $\text{deg}_{\mathbb{Z}_2} F$, the Smale degree, cf [Sm]. When the index of F is zero, the degree is given by the number of preimages of a regular value modulo 2. If the spaces or map have a suitable orientation, this can be extended to a \mathbb{Z} -valued degree $\text{deg}_{\mathbb{Z}} F$, cf [ET] for instance. Essentially an immediate consequence of its definition and the Sard–Smale theorem [Sm] is that if such a degree is non-zero,

$$\text{deg}_{\mathbb{Z}_2} F \neq 0,$$

then F is surjective.

The definition of degree above may be extended to maps which are almost proper in the sense above, cf [BFP] for instance. Thus, let $\partial \mathcal{E}^o$ be the boundary of \mathcal{E}^o within the space \mathcal{E}_S of static vacuum solutions. This is the space of solutions in \mathcal{E}_S satisfying (1.10) but not (1.8). Let $Z = \Pi_B(\partial \mathcal{E}^o) \subset \text{Met}^{m,\alpha}(\partial M) \times C^{m-1,\alpha}(\partial M)$ be the image of $\partial \mathcal{E}^o$ under the boundary map Π_B and let $\mathcal{E}^P = (\Pi^o)^{-1}([\text{Met}^{m,\alpha}(\partial M) \times C^{m-1,\alpha}(\partial M)] \setminus Z)$ be the corresponding inverse image. Then, as discussed in section 5, the induced boundary map (restriction of Π^o to \mathcal{E}^P)

$$\Pi^P : \mathcal{E}^P \rightarrow [\text{Met}^{m,\alpha}(\partial M) \times C^{m-1,\alpha}(\partial M)] \setminus Z,$$

is proper. In particular, \mathcal{E}^P has a finite number of connected components \mathcal{E}^{P_i} and the induced boundary map Π^{P_i} on \mathcal{E}^{P_i} has a well-defined \mathbb{Z}_2 -valued degree (with respect to a component of the target space). We also expect that Π^{P_i} has a well-defined \mathbb{Z} -valued degree.

Let \mathcal{E}^{P_0} be the component of \mathcal{E}^P containing the standard round exterior flat solution as following (1.8), and let \mathcal{T}_0 be the component of $[\text{Met}^{m,\alpha}(\partial M) \times C_+^{m-1,\alpha}(\partial M)] \setminus Z$ containing the corresponding standard boundary data $(\gamma_{+1}, 2)$. One thus has the boundary map

$$\Pi^{P_0} : \mathcal{E}^{P_0} \rightarrow \mathcal{T}_0. \tag{1.11}$$

A further main result of this paper is the computation of this degree.

Theorem 1.3. *For the map Π^{P_0} in (1.11), one has*

$$\text{deg}_{\mathbb{Z}_2} \Pi^{P_0} = 1. \tag{1.12}$$

The proof of theorem 1.3 is based on the well-known black hole uniqueness theorem for the Schwarzschild metric, cf [I, R, BM].

It follows that the boundary map Π^{P_0} maps \mathcal{E}^{P_0} surjectively onto the component \mathcal{T}_0 of the target $[\text{Met}^{m,\alpha}(\partial M) \times C_+^{m-1,\alpha}(\partial M)] \setminus Z$. In particular, the image of Π^{P_0} and so the image of Π_B has non-empty interior in the target space $\text{Met}^{m,\alpha}(\partial M) \times C_+^{m-1,\alpha}(\partial M)$; this has not been previously known, cf remark 5.4 for further discussion.

Of course, the main issue at this point is what can be said about the structure of the set Z ? It is of first category (so non-generic) but its more detailed structure awaits future study. An alternate approach bypassing the issue of the boundary $\partial \mathcal{E}^o$ would be to find conditions on the boundary data (γ, H) which imply ∂M is outer-minimizing as in (1.8) in any static vacuum extension (M, g, u) of (γ, H) . For example, in \mathbb{R}^3 (where $u = 1$) convexity suffices, which is expressed in terms of boundary data as $K_\gamma > 0$ where K is the Gauss curvature. It is an open problem whether this condition also suffices for general static vacuum solutions.

The contents of the paper are briefly as follows. In section 2, we present background information on the structure of static vacuum solutions and choices of gauge. Section 3 discusses elliptic boundary value problems for the Einstein equations and proves the basic structure theorems on the moduli space \mathcal{E}_S and the boundary map Π_B , including theorem 1.1. In section 4, we then prove the requisite *a priori* estimates and establish the almost properness of Π^o on \mathcal{E}^o , proving theorem 1.2. Finally, section 5 contains the computation of the degree of Π^{P_0} and closes with several related remarks.

2. Background discussion

Let M be a 3-manifold with compact boundary ∂M , and with a single open end E . (All of the results of this section and of section 3 hold in all dimensions, but for simplicity, we restrict to dimension 3). *A priori*, ∂M need not be connected, although this will be assumed later on. As following (1.2) and (1.3), we let $\mathcal{M} = S^1 \times M$. Almost all of the discussion and computation in sections 2 and 3 is carried out on the 4-manifold \mathcal{M} and $g_{\mathcal{M}}$ will often be denoted g for notational simplicity.

Let $\text{Met}_S^{m,\alpha}(\mathcal{M}) = \text{Met}_S^{m,\alpha}(\mathcal{M})$ be the space of complete (up to the boundary) $C^{m,\alpha}$ static metrics on \mathcal{M} , i.e. metrics of the form (1.2), $m \geq 2$. One has

$$\text{Met}_S^{m,\alpha}(\mathcal{M}) \simeq \text{Met}^{m,\alpha}(M) \times C_+^{m,\alpha}(M), \tag{2.1}$$

where $C_+^{m,\alpha}(M)$ is the space of positive $C^{m,\alpha}$ functions on M . The space $\mathbb{E}_S = \mathbb{E}_S(\mathcal{M})$ of static Einstein (Ricci-flat) metrics on \mathcal{M} is equivalent to the space of pairs $g_{\mathcal{M}} = (g_M, u) \in \text{Met}(M) \times C_+(M)$ satisfying (1.1) or (1.3) (the smoothness indices will be occasionally dropped when unimportant). It is well-known [M] that away from the boundary, solutions of the static vacuum equations are analytic in appropriate coordinates.

Recall that a complete metric $g \in \text{Met}^{m,\alpha}(E)$ on an end E is asymptotically flat if E is diffeomorphic to $\mathbb{R}^3 \setminus B$, where B is a 3-ball, and there exists a diffeomorphism $F : \mathbb{R}^3 \setminus B \rightarrow E$ such that, in the chart F ,

$$g_{ij} = \delta_{ij} + O(r^{-1}), \quad \partial_k g_{ij} = O(r^{-2}), \quad \partial_k \partial_\ell g_{ij} = O(r^{-3}), \tag{2.2}$$

in the standard Euclidean coordinates on \mathbb{R}^3 . The static vacuum equations (1.1) are invariant under multiplication of the potential u by constants. Throughout the paper, we assume that u is normalized so that $u \rightarrow 1$ at infinity, and that u is asymptotically constant in the sense that

$$u = 1 + O(r^{-1}), \quad \partial_k u = O(r^{-2}), \quad \partial_k \partial_\ell u = O(r^{-3}). \tag{2.3}$$

Thus the 4-metric $g_{\mathcal{M}}$ is asymptotic to the product $S^1 \times \mathbb{R}^3$ at infinity.

It is proved in [A2] that ends of static vacuum solutions (M, g, u) are either asymptotically flat or parabolic, where parabolic is understood in the sense of potential theory; equivalently, parabolic ends have small volume growth in that the area of geodesic spheres grows slower than $r^{1+\varepsilon}$, for any $\varepsilon > 0$. Moreover, asymptotically flat ends are strongly asymptotically flat in that the metric and potential have asymptotic expansions of the form

$$g_{ij} = \left(1 + \frac{2m}{r}\right) \delta_{ij} + \dots, \quad u = 1 - \frac{2m}{r} + \dots, \tag{2.4}$$

where the mass m may a priori be any value $m \in \mathbb{R}$, cf also [KO]. These two behaviors, asymptotically flat and parabolic, are radically different and there is no curve of asymptotic structures for static vacuum solutions which joins them. The finer behavior of asymptotically flat ends is described by the mass parameter m and higher multipole moments, cf [BS].

Let \tilde{g} be a fixed asymptotically flat (background) metric in \mathbb{E}_S ; henceforth \mathbb{E}_S will denote the space of asymptotically flat static vacuum Einstein solutions. The static Einstein equations are not elliptic, due to their invariance under diffeomorphisms, and for several reasons one needs to choose an elliptic gauge. To begin, we consider the Bianchi gauge, and define

$$\begin{aligned} \Phi_{\tilde{g}} : \text{Met}_S^{m,\alpha}(\mathcal{M}) &\rightarrow S^{m-2,\alpha}(\mathcal{M}), \\ \Phi_{\tilde{g}}(g) &= \text{Ric}_g + \delta_g^* \beta_{\tilde{g}}(g), \end{aligned} \tag{2.5}$$

where $\beta_{\tilde{g}}$ is the Bianchi operator, $\beta_{\tilde{g}}(g) = \delta_{\tilde{g}}(g) + \frac{1}{2} \text{dtr}_{\tilde{g}}(g)$. Also, $(\delta^* X)(A, B) = \frac{1}{2} (\langle \nabla_A X, B \rangle + \langle \nabla_B X, A \rangle)$ and $\delta X = -\text{tr}(\delta^* X)$ is the divergence. The operator $\Phi_{\tilde{g}}$ is a C^∞ smooth map into the space $S^{m-2,\alpha}(\mathcal{M})$ of static symmetric bilinear forms on \mathcal{M} ; note that here a symmetric bilinear form is referred to as static if its components do not depend on time and the mixed time/space components vanish.

Using standard formulas for the linearization of the Ricci and scalar curvatures, cf [Be, p 63] for instance, the linearization of Φ at $g = \tilde{g} \in \mathbb{E}_S$ is given by

$$L(h) = 2(D\Phi_{\tilde{g}})(h) = D^* Dh - 2R(h), \tag{2.6}$$

where $R(h)(X, Y) = \langle R(e_i, X)Y, h(e_i) \rangle$ with e_i an orthonormal basis. Clearly, L is elliptic and formally self-adjoint. In section 3 we will discuss boundary value problems for Φ and L .

Next, the asymptotic behavior in the asymptotically flat end E requires the introduction of suitable weighted function spaces. We will use the standard weighted Hölder spaces, although one could equally well use weighted Sobolev spaces. Thus, define $\text{Met}_\delta^{m,\alpha}(\mathcal{M}) \subset \text{Met}_S^{m,\alpha}(\mathcal{M})$ to be the subspace of metrics which decay to Euclidean data at a rate $r^{-\delta}$ at infinity; more precisely, the component functions g_{ij} and u of $g_{\mathcal{M}}$ should satisfy

$$g_{ij} - \delta_{ij} \in C_\delta^{m,\alpha}(\mathbb{R}^3 \setminus B), \quad u - 1 \in C_\delta^{m,\alpha}(\mathbb{R}^3 \setminus B).$$

Here C_δ^m consists of functions v such that

$$\|v\|_{C_\delta^m} = \sum_{k=0}^m \sup r^{k+\delta} |\nabla^k v| < \infty,$$

while $C_\delta^{m,\alpha}$ consists of functions such that

$$\|v\|_{C_\delta^{m,\alpha}} = \|v\|_{C_\delta^m} + \sup_{x,y} \left[\min(r(x), r(y))^{m+\alpha+\delta} \frac{|\nabla^m v(x) - \nabla^m v(y)|}{|x-y|^\alpha} \right] < \infty,$$

cf [B1, LP]. Throughout the following, we assume the decay rate δ is fixed, and chosen to satisfy

$$\frac{1}{2} < \delta < 1. \tag{2.7}$$

It is well-known, cf [B1, LP], that the Laplacian on functions, and Laplace-type operators on tensors, as in (2.6), are Fredholm when acting on these weighted Hölder spaces.

The map Φ in (2.5) clearly induces a smooth map

$$\Phi : \text{Met}_\delta^{m,\alpha}(\mathcal{M}) \rightarrow S_\delta^{m-2,\alpha}(\mathcal{M}). \tag{2.8}$$

Observe that g is Einstein if $\Phi_{\tilde{g}}(g) = 0$ and $\beta_{\tilde{g}}(g) = 0$, so that g is in Bianchi gauge with respect to \tilde{g} . (Although $\Phi_{\tilde{g}}$ is defined for all $g \in \text{Met}_\delta^{m,\alpha}(\mathcal{M})$, we will only consider it acting on g near \tilde{g} .)

Given $\tilde{g} \in \mathbb{E}_S$, let $\text{Met}_C^{m,\alpha}(\mathcal{M}) \subset \text{Met}_\delta^{m,\alpha}(\mathcal{M})$ be the space of $C^{m,\alpha}$ smooth static AF Riemannian metrics on \mathcal{M} , satisfying the Bianchi gauge constraint

$$\beta_{\tilde{g}}(g) = 0 \text{ at } \partial\mathcal{M}. \tag{2.9}$$

As above,

$$\Phi : \text{Met}_C^{m,\alpha}(\mathcal{M}) \rightarrow S_\delta^{m-2,\alpha}(\mathcal{M}).$$

Similarly let $Z_C^{m,\alpha}$ be the space of metrics $g \in \text{Met}_C^{m,\alpha}(\mathcal{M})$ satisfying $\Phi_{\tilde{g}}(g) = 0$, and let

$$\mathbb{E}_C \subset Z_C \tag{2.10}$$

be the subset of static Einstein metrics $g = g_N$, $\text{Ric}_g = 0$ in Z_C . The following result justifies the use of the operator Φ to study \mathbb{E}_C .

Proposition 2.1. *Any static metric $g = g_M \in Z_C$ sufficiently close to \tilde{g} is necessarily Einstein, $g \in \mathbb{E}_C$. Moreover, this also holds infinitesimally in the following sense. Let κ be an infinitesimal deformation of $g \in Z_C$, i.e. $\kappa \in \text{Ker } D\Phi$. If $\beta_{\tilde{g}}(g) = 0$, (for example $\tilde{g} = g$), then*

$$\beta_{\tilde{g}}(\kappa) = 0, \tag{2.11}$$

and κ is an infinitesimal Einstein deformation, i.e. the variation of g in the direction κ preserves (1.3) to first order.

Proof. Since $g \in Z_C$, one has $\Phi(g) = 0$, i.e.

$$\text{Ric}_g + \delta_g^* \beta_{\tilde{g}}(g) = 0.$$

Applying the Bianchi operator β_g and using the Bianchi identity $\beta_g(\text{Ric}_g) = 0$ gives

$$\beta_g(\delta_g^*(\beta_{\tilde{g}}(g))) = 0. \tag{2.12}$$

Set $V = \beta_{\tilde{g}}(g)$, and notice that a simple computation produces the Weitzenböck formula $2\beta_g \delta_g^*(V) = D^*DV - \text{Ric}(V)$. Also, since $g, \tilde{g} \in \text{Met}_\delta^{m,\alpha}(\mathcal{M})$, $V \in \chi_{1+\delta}^{m-1,\alpha}(M)$, where $\chi_{1+\delta}^{m-1,\alpha}(M)$ is the space of vector fields whose components are in $C_{1+\delta}^{m-1,\alpha}(M)$. When acting on vector fields V with $V = 0$ on ∂M , as in (2.9), the operator D^*D is positive, with trivial kernel. Namely, if $W \in C_{1+\delta}^{m-1,\alpha}$ is in the kernel of D^*D , then integrating by parts gives

$$\int_{B(r)} |DW|^2 + \int_{S(r)} \langle W, \nabla_N W \rangle = 0,$$

where $B(r) = \{x \in M : \text{dist}(x, \partial M) \leq r\}$ and N is the outward unit normal. (Since $W = 0$ on ∂M , there is no boundary term at ∂M). Letting $r \rightarrow \infty$, the boundary integral tends to 0 and so $DW = 0$, which in turn implies $W = 0$.

Since D^*D is self-adjoint and Fredholm, it has a smallest positive eigenvalue bounded away from 0. For g sufficiently close to \tilde{g} , $|\text{Ric}| \sim 0$ pointwise and $\text{Ric}(V) \in C_{3+\delta}^{m-3,\alpha}(M)$, so we may assume that $2\beta_g \delta_g^*$ is a positive operator on V . Hence, again since $V = 0$ on ∂M , the only solution of (2.12) is $V = 0$, which implies $g \in \mathbb{E}_C$.

To prove the second statement, let $g_t = g + t\kappa$. Applying the Bianchi operator β_{g_t} to $\Phi(g_t)$ gives

$$\beta_{g_t} \Phi(g_t) = \beta_{g_t} \delta_{g_t}^* (\beta_{\tilde{g}}(g_t)). \tag{2.13}$$

Taking the derivative with respect to t at $t = 0$, one has $(\beta_{g_t} \Phi(g_t))' = \beta' \Phi + \beta \Phi'$. Both terms here vanish since $g \in Z_C$ and κ is formally tangent to Z_C . Hence the variation of the right-hand side of (2.13) vanishes. Since $\beta_{\tilde{g}}(g) = 0$, this gives $\beta_g \delta_g^* (\beta_{\tilde{g}}(\kappa)) = 0$. The equation (2.11) then follows exactly as following (2.12), with $V = \beta_{\tilde{g}}(\kappa)$. \square

Let $\mathcal{D}_1^{m+1,\alpha}$ denote the space of $C_\delta^{m+1,\alpha}$ static diffeomorphisms of \mathcal{M} which equal the identity on $\partial \mathcal{M}$. These are diffeomorphisms which decay to the identity at the rate $r^{-\delta}$ and are independent of the t or θ -variable in (1.2). The group \mathcal{D}_1 acts freely and continuously on $\text{Met}(\mathcal{M})$ and \mathbb{E}_S by pullback and one has the following local slice theorem for this action; we refer to [A3] for the proof.

Lemma 2.2. *Given any $\tilde{g} \in \mathbb{E}_S^{m,\alpha}$ and $g \in \text{Met}_\delta^{m,\alpha}(\mathcal{M})$ near \tilde{g} , there exists a unique diffeomorphism $\phi \in \mathcal{D}_1^{m+1,\alpha}$ close to the identity, such that*

$$\beta_{\tilde{g}}(\phi^* g) = 0. \tag{2.14}$$

In particular, $\phi^ g \in \text{Met}_C^{m,\alpha}(\mathcal{M})$.*

Lemma 2.2 implies that if $g \in \mathbb{E}_S^{m,\alpha}$ is a static Einstein metric near \tilde{g} , then g is isometric, by a diffeomorphism in $\mathcal{D}_1^{m+1,\alpha}$, to an Einstein metric in $\mathbb{E}_C^{m,\alpha}$, so that $\mathbb{E}_C^{m,\alpha}$ is a slice for $\mathbb{E}_S^{m,\alpha}$ under the action of $\mathcal{D}_1^{m+1,\alpha}$.

To prove that the moduli space \mathcal{E} is a smooth Banach manifold, (cf theorem 3.6), it is important to have a gauge with choice of boundary data in which the Einstein equations form a self-adjoint elliptic boundary value problem. This is not the case for the operator Φ and we are not aware of geometrically natural self-adjoint boundary conditions for Φ . For this reason, we will also consider another natural gauge, namely the divergence-free gauge.

To do this, instead of Φ , consider

$$\hat{\Phi}(g) = \hat{\Phi}_{\tilde{g}}(g) = \text{Ric}_g - \frac{s}{2}g + \delta_g^* \delta_{\tilde{g}} g, \tag{2.15}$$

where s is the scalar curvature of $g = g_{\mathcal{M}}$. The linearization of $\hat{\Phi}$ at $g = \tilde{g} \in \mathbb{E}_S$ is given by

$$\hat{L}(h) = 2(D\hat{\Phi}_{\tilde{g}})_g(h) = D^*Dh - 2R(h) - (D^2 \text{tr} h + \delta \delta h g) + \Delta \text{tr} h g. \tag{2.16}$$

In analogy to (2.9), define

$$\text{Met}_D^{m,\alpha}(M) = \{g \in \text{Met}_\delta^{m,\alpha}(\mathcal{M}) : \delta_{\tilde{g}} g = 0 \text{ at } \partial \mathcal{M}\}. \tag{2.17}$$

Similarly, let $Z_D^{m,\alpha} = \hat{\Phi}^{-1}(0) \cap \text{Met}_D^{m,\alpha}(M)$ and $\mathbb{E}_D \subset Z_D$ be the space of Einstein metrics in divergence-free gauge with respect to $\tilde{g} \in \mathbb{E}_S$.

It is easy to see that proposition 2.1 and lemma 2.2 hold in this divergence-free gauge in place of the previous Bianchi gauge, with essentially the same proof. Thus $\mathbb{E}_D = Z_D$ and for $g \in \mathbb{E}_D$,

$$\delta_{\tilde{g}} g = 0 \text{ on } \mathcal{M}. \tag{2.18}$$

Moreover, the diffeomorphism group \mathcal{D}_1 transforms one gauge choice uniquely to the other. For instance, suppose $\beta_{\tilde{g}}(g) = 0$. Then we claim there is a unique $\phi \in \mathcal{D}_1^{m+1,\alpha}$ such that

$$\delta_{\tilde{g}}(\phi^*g) = 0. \tag{2.19}$$

At the linearized level, with $g = \tilde{g}$, this amounts to finding a vector field V with $V = 0$ on $\partial\mathcal{M}$ such that if $\beta_{\tilde{g}}h = 0$ then $\delta_{\tilde{g}}(h + \delta^*V) = 0$. This equation is equivalent to the equation $\delta\delta^*V = \frac{1}{2}d \operatorname{tr} h$, which is uniquely solvable for V with $V = 0$ on $\partial\mathcal{M}$. The local result in (2.19) then follows from the inverse function theorem.

3. The moduli space

In this section, we study boundary value problems for the elliptic operators Φ and $\hat{\Phi}$, and use this to prove that the moduli space \mathcal{E}_S of static vacuum solutions is a smooth Banach manifold for which the boundary map Π_B is Fredholm, of Fredholm index 0, cf theorem 3.6.

We begin with the Bianchi-gauged Einstein operator Φ in (2.5), i.e.

$$\Phi_{\tilde{g}}(g) = \operatorname{Ric}_g + \delta_g^* \beta_{\tilde{g}}(g).$$

Let A denote the second fundamental form of ∂M in M , $A(X, Y) = \langle \nabla_X N, Y \rangle$, where N is the unit inward normal into M , X, Y tangent to ∂M . Similarly, let $H_M = \operatorname{tr} A$ denote the mean curvature of ∂M in M . Throughout the paper W^T will denote the restriction or the orthogonal projection of a tensor W to $T(\partial N)$ or $T(\partial M)$.

Proposition 3.1. *Near any given background solution $\tilde{g} \in \mathbb{E}_S^{m,\alpha}$, the operator $\Phi = \Phi_{\tilde{g}}$ in (2.5) with boundary conditions:*

$$\beta_{\tilde{g}}(g) = 0, \quad g|_{T(\partial M)} = \gamma_M, \quad H_M = h_M \text{ at } \partial\mathcal{M}, \tag{3.1}$$

is an elliptic boundary value problem of Fredholm index 0.

Here the induced metric γ_M is in $\operatorname{Met}^{m,\alpha}(\partial M)$ while the mean curvature h_M of ∂M in (M, g_M) is in $C^{m-1,\alpha}(\partial M)$. Note that the potential u does not enter this boundary data and so is formally undetermined at ∂M . Also the static property implies that $\beta_{\tilde{g}}(g)$ vanishes in the vertical direction, $\beta_{\tilde{g}}(g)(\partial_\theta) = 0$.

Proof. It suffices to prove that the leading order part of the linearized operators forms an elliptic system. Recall from (2.6) that the linearization of Φ at $\tilde{g} = g$ is given by

$$L(h) = 2(D\Phi_g)(h) = D^*Dh - 2R(h).$$

The leading order symbol of $L = 2D\Phi$ at ξ' is

$$\sigma(L) = -|\xi'|^2 I, \tag{3.2}$$

where I is the $Q \times Q$ identity matrix, with $Q = (n(n+1)/2) + 1$; Q is the sum of the dimension of the space of symmetric bilinear forms on \mathbb{R}^n , together with the extra vertical S^1 direction. Here $n = 3$ but we give the proof for general dimensions. For static metrics, all components of the metric are locally functions on \mathbb{R}^n , and all derivatives in the vertical S^1 direction are trivial. In the following, the subscript 0 represents the direction normal to ∂M in M , (or $\partial\mathcal{M}$ in \mathcal{M}), subscript 1 denotes the vertical direction, tangent to S^1 , while indices 2 through n represent the directions tangent to ∂M . Note that one has $h_{1\alpha} = 0$, for all $\alpha \neq 1$. The positive roots of (3.2) are $i|\xi|$, where $\xi' = (\xi_0, \xi)$, with multiplicity Q at $\xi \in T^*(\mathbb{R}^n)$.

Writing $\xi' = (z, \xi_i)$, $i = 2, \dots, n$, (as above $\xi_1 = 0$), the symbols of the leading order terms in the boundary operators are given by:

$$-2izh_{0k} - 2i \sum_{j \geq 2} \xi_j h_{jk} + i\xi_k \text{tr } h = 0, \quad k \geq 2,$$

$$-2izh_{00} - 2i \sum_{k \geq 2} \xi_k h_{0k} + iz \text{tr } h = 0,$$

$$h^T = (\gamma')^T, \text{ and } (H_M)'_h = \omega.$$

This gives $n + \frac{n(n-1)}{2} + 1 = Q$ boundary equations, as required. Ellipticity requires that the operator defined by the boundary symbols above has trivial kernel when z is set to the root $i|\xi|$. Carrying this out then gives the system

$$2|\xi| h_{0k} - 2i \sum_{j \geq 2} \xi_j h_{jk} + i\xi_k \text{tr } h = 0, \quad k \geq 2, \tag{3.3}$$

$$2|\xi| h_{00} - 2i \sum_{k \geq 2} \xi_k h_{0k} - |\xi| \text{tr } h = 0, \tag{3.4}$$

$$h_{11} = \phi, \quad h^T = 0, \quad (H_M)'_h = 0, \tag{3.5}$$

where ϕ is an undetermined function.

Multiplying (3.3) by $i\xi_k$ and summing gives, via (3.5),

$$2|\xi| i \sum_{k \geq 2} \xi_k h_{0k} = |\xi|^2 \text{tr } h.$$

Substituting (3.4) on the term on the left above then gives

$$2|\xi|^2 h_{00} - 2|\xi|^2 \text{tr } h = 0.$$

Since $\text{tr } h = h_{00} + \phi$, it follows that $\phi = 0$.

Next, to compute H'_M , we first observe that in general

$$2A'_h = \nabla_N h + 2A \circ h - 2\delta^*(h(N)^T) - \delta^*(h_{00}N). \tag{3.6}$$

This follows by differentiating the defining formula $2A = \mathcal{L}_N g$, and using the identities $2N'_h = -2h(N)^T - h_{00}N$, $\mathcal{L}_N h = \nabla_N h + 2A \circ h$. Since $H_M = \text{tr}_M A$, $H'_h = \text{tr}_M A'_h - \text{tr}_M A \circ h$ and so

$$2(H_M)'_h = \text{tr}_M(\nabla_N h - 2\delta^*(h(N)^T) - \delta^*(h_{00}N)). \tag{3.7}$$

Hence the symbol of $2(H_M)'_h$ is given by $\sum_{k \geq 2} (izh_{kk} - 2i\xi_k h_{0k})$. Setting this to 0 at the root $z = i|\xi|$ gives

$$\sum_{k \geq 2} (|\xi| h_{kk} + 2i\xi_k h_{0k}) = 0. \tag{3.8}$$

Via (3.5), this gives $-2i \sum_{k \geq 2} \xi_k h_{0k} = 0$, and substituting this in (3.4) and using the fact that $\phi = 0$ gives

$$2|\xi| h_{00} - |\xi| h_{00} = 0,$$

so that $h_{00} = 0$. It follows from (3.3) that $h_{0k} = 0$ and hence $h = 0$. This proves ellipticity of the boundary value problem (3.1) and the Fredholm property follows from the fact that the Laplace-type operator L is Fredholm on $\text{Met}_\delta^{m,\alpha}$, cf [LP].

Finally, it is straightforward to verify that the boundary data (3.1) may be continuously deformed through elliptic boundary data to elliptic boundary data for which L is self-adjoint and so of index 0. This is proved in [A3] in a slightly different setting and the proof carries

over here with only minor change, and so we refer to [A3] for further details. The homotopy invariance of the index then completes the proof. \square

As noted in section 2, we are not aware of a geometrically natural self-adjoint elliptic boundary value problem for Φ . In particular, the boundary conditions (3.1) are not self-adjoint. This property is important for the proof of theorem 3.6, and for this reason, we turn to the operator $\hat{\Phi}$ in (2.15) with linearization at $\tilde{g} = g$ given by \hat{L} in (2.16).

Regarding boundary conditions for \hat{L} , for $h \in S_\delta^{m-2,\alpha}(\mathcal{M})$, let $h^T = h|_{\partial\mathcal{M}}$ and $[h^T]_0$ be the projection of h^T onto the space of forms trace-free with respect to $\gamma = \gamma_{\mathcal{M}}$. Similarly, H'_h denotes here the linearization of the mean curvature $H = H_{\mathcal{M}}$ of $\partial\mathcal{M} \subset \mathcal{M}$.

We then have:

Lemma 3.2. *The operator \hat{L} with boundary conditions*

$$\delta h = 0, \quad [h^T]_0 = 0, \quad H'_h = 0, \tag{3.9}$$

is a self-adjoint elliptic operator. Moreover, under the first two conditions $\delta h = 0$ and $[h^T]_0 = 0$, the operator \hat{L} is self-adjoint exactly for the boundary condition $H'_h = 0$.

Proof. It is a rather long (and uninteresting) calculation to prove that the operator \hat{L} with boundary data (3.9) forms an elliptic system; this has been verified by computer computation using Maple. More conceptually, instead we will make use of proposition 3.1 to simplify the proof. First, recall, [ADN, T], that ellipticity of a boundary value problem is equivalent to the existence of a uniform estimate

$$\|h\|_{C^{m,\alpha}} \leq C(\|\hat{L}(h)\|_{C^{m-2,\alpha}} + \|B_j(h)\|_{C^{m-j,\alpha}} + \|h\|_{C^0}), \tag{3.10}$$

where B_j is the part of the boundary operator of order j , together with such an estimate for the adjoint operator. As seen below, the boundary value problem is self-adjoint, so it suffices to establish (3.10).

First, it is simple to prove (3.10) for L in place of \hat{L} via a slight modification of the proof of proposition 3.1. Namely, for the boundary condition $[h^T]_0 = 0$, we have $h^T = \phi\gamma$ on $\partial\mathcal{M}$, (in place of (3.5)). Note also that (3.3) and (3.4) hold, but without the $\text{tr } h$ terms. The analogue of (3.3) then gives

$$|\xi|h_{0k} = i\xi_k\phi,$$

and hence, via the analogue of (3.4),

$$|\xi|^2 h_{00} = -|\xi|^2\phi,$$

so that $h_{00} + \phi = 0$. Next, via the condition $H'_h = 0$, the analogue of (3.8) becomes

$$\sum_{k \geq 1} (|\xi|h_{kk} + 2i\xi_k h_{0k}) = 0,$$

which gives

$$n|\xi|\phi = -2i \sum \xi_k h_{0k} = 2|\xi|\phi.$$

Since $n \geq 3$, this implies $\phi = 0$, and so $h_{00} = 0$, hence $h_{0k} = 0$. It follows that $h = 0$, which proves ellipticity of L with the boundary conditions (3.9). Thus, (3.10) holds with L in place of \hat{L} .

Next, one has

$$\hat{L} = L - (D^2 \text{tr } h - \Delta \text{tr } h g) - \delta\delta h g. \tag{3.11}$$

Thus to prove (3.10), it suffices to prove

$$\|\delta h\|_{C^{m-1,\alpha}} \leq C(\|\hat{L}(h)\|_{C^{m-2,\alpha}} + \|B_j(h)\|_{C^{m-j,\alpha}} + \|h\|_{C^0}), \tag{3.12}$$

$$\|D^2 \operatorname{tr} h\|_{C^{m-2,\alpha}} \leq C(\|\hat{L}(h)\|_{C^{m-2,\alpha}} + \|B_j(h)\|_{C^{m-j,\alpha}} + \|h\|_{C^0}). \tag{3.13}$$

From (2.15) and (2.16) and the Bianchi identity, (as in (2.13)), one has $\delta \hat{L}(h) = 2\delta \delta^*(\delta h)$ and the operator $\delta \delta^*$ is elliptic with respect to Dirichlet boundary conditions. Since the boundary data δh in (3.9) is included in the boundary operators B_j , this proves (3.12).

Using this and taking the trace of (3.11) shows that

$$\|D^2 \operatorname{tr} h\|_{C^{m-2,\alpha}} \leq C(\|\hat{L}(h)\|_{C^{m-2,\alpha}} + \|B_j(h)\|_{C^{m-j,\alpha}} + \|NN(\operatorname{tr} h)\|_{C^{m-2,\alpha}} + \|h\|_{C^0}),$$

so that it suffices to prove that the boundary conditions B cover $NN(\operatorname{tr} h)$. For this, a simple computation using (3.7), (cf also (3.19) below), gives

$$N(\operatorname{tr} h) = 2H'_h - \delta((h(N))^T) - (\delta h)(N) + O(h), \tag{3.14}$$

where $O(h)$ is of differential order 0 in h . Using the standard interpolation $\|h\|_{C^{m-1,\alpha}} \leq \varepsilon \|h\|_{C^{m,\alpha}} + \varepsilon^{-1} \|h\|_{C^0}$, where $\varepsilon > 0$ is an arbitrary constant, shows that it suffices here and below only to consider terms with the leading number of derivatives of h .

Now the Gauss equations at $\partial \mathcal{M}$ are $|A|^2 - H^2 + s_{\gamma_{\mathcal{M}}} = s_{g_{\mathcal{M}}} - 2\operatorname{Ric}(N, N)$ and hence,

$$(|A|^2 - H^2 + s_{\gamma_{\mathcal{M}}})'_h = -2\hat{L}(h)(N, N) + 2\delta^* \delta(h)(N, N) + O(h).$$

One has $s'_{\gamma_{\mathcal{M}}}(h^T) = -\Delta \operatorname{tr} h^T + \delta \delta(h^T) + O(h^T)$ and A'_h, H'_h only involve first order derivatives in h . Writing then $h^T = B_0(h) + \frac{1}{n} \operatorname{tr}_{\partial \mathcal{M}} h \gamma_{\mathcal{M}}$, it follows that $\operatorname{tr}_{\partial \mathcal{M}} h$ at $\partial \mathcal{M}$ is controlled by $\hat{L}(h), B_j(h)$, in that

$$\|\operatorname{tr}_{\partial \mathcal{M}} h\|_{C^{m,\alpha}} \leq C(\|h\|_{C^{m-1,\alpha}} + \|\hat{L}(h)\|_{C^{m-2,\alpha}} + \|B_j(h)\|_{C^{m-j,\alpha}}),$$

and hence

$$\|h^T\|_{C^{m,\alpha}} \leq C(\|h\|_{C^{m-1,\alpha}} + \|\hat{L}(h)\|_{C^{m-2,\alpha}} + \|B_j(h)\|_{C^{m-j,\alpha}}),$$

i.e. h^T is controlled at $\partial \mathcal{M}$ by $\hat{L}(h)$ and $B_j(h)$. Next, at $\partial \mathcal{M}$, one has $-(\delta h)(T) = \nabla_N h(N, T) + \nabla_{e_i} h(e_i, T)$, which then gives control as above on $(\nabla_N h)(N, T)$, and so control on $\nabla_N(h(N)^T)$. In turn, this gives then control on $\delta_{\partial \mathcal{M}}(\nabla_N(h(N)^T))$, which modulo lower order (curvature) terms, equals $N(\delta(h(N)^T))$. The N -derivative of (3.14) also holds and shows that control of $N(\delta(h(N)^T))$ implies control of $NN(\operatorname{tr} h)$, so that (3.13) holds, provided $N(H'_h)$ is controlled. But the Riccati equation gives $N(H) = -|A|^2 - \operatorname{Ric}(N, N)$; taking the linearization of this in the direction h shows that $N(H'_h)$ is indeed controlled by $\hat{L}(h)$ and the boundary conditions B_j . This completes the proof of ellipticity.

Next, we prove the operator \hat{L} with boundary conditions (3.9) is self-adjoint. To begin, integrating the terms in the expression (2.16) for \hat{L} by parts over \mathcal{M} gives

$$\begin{aligned} \int_{\mathcal{M}} \langle D^* D(h), k \rangle + \int_{\partial \mathcal{M}} \langle \nabla_N h, k \rangle &= \int_{\mathcal{M}} \langle D^* D(k), h \rangle + \int_{\partial \mathcal{M}} \langle \nabla_N k, h \rangle, \\ \int_{\mathcal{M}} \delta \delta h \operatorname{tr} k + \int_{\partial \mathcal{M}} (\delta h)(N) \operatorname{tr} k &= \int_{\mathcal{M}} \langle h, D^2(\operatorname{tr} k) \rangle - \int_{\partial \mathcal{M}} h(N, \operatorname{dtr} k), \end{aligned}$$

and

$$\int_{\mathcal{M}} (\Delta \operatorname{tr} h) \operatorname{tr} k - \int_{\partial \mathcal{M}} N(\operatorname{tr} h) \operatorname{tr} k = \int_{\mathcal{M}} (\Delta \operatorname{tr} k) \operatorname{tr} h - \int_{\partial \mathcal{M}} N(\operatorname{tr} k) \operatorname{tr} h.$$

Here the boundary terms on $S(r)$ all tend to 0 as $r \rightarrow \infty$, since the components of h and k are in $C_\delta^{m,\alpha}$ and $\delta > \frac{1}{2}$. It follows that

$$\int_{\mathcal{M}} \langle \hat{L}(h), k \rangle + \int_{\partial \mathcal{M}} \langle B(h), k \rangle = \int_{\mathcal{M}} \langle \hat{L}(k), h \rangle + \int_{\partial \mathcal{M}} \langle B(k), h \rangle, \tag{3.15}$$

where

$$\langle B(k), h \rangle = \langle \nabla_N k, h \rangle + h(N, \text{dtr } k) - (\delta k)(N) \text{tr } h - \text{tr } h N(\text{tr } k). \quad (3.16)$$

Setting $Z(k, h) = \langle B(k), h \rangle - \langle B(h), k \rangle$, we thus need to show that

$$\int_{\partial \mathcal{M}} Z(h, k) = 0, \quad (3.17)$$

when h, k satisfy the boundary conditions (3.9).

Thus suppose h and k both satisfy (3.9). A simple calculation shows that $(\delta k)(T) = 0$ is equivalent to

$$(\nabla_N k)(N)^T = \delta_{\partial \mathcal{M}}(k^T) - \alpha(k(N)), \quad (3.18)$$

where $\alpha(k(N)) = [A(k(N)) + Hk(N)^T]$, (all taken on $\partial \mathcal{M}$), while $(\delta k)(N) = 0$ is equivalent to

$$N(k_{00}) = \delta_{\partial \mathcal{M}}(k(N)^T) + \langle A, k \rangle - k_{00}H. \quad (3.19)$$

The same equations hold for h , and one also has

$$h^T = \phi_h \gamma, \quad \text{and} \quad k^T = \phi_k \gamma. \quad (3.20)$$

We thus need to calculate

$$B(k, h) = \langle \nabla_N k, h \rangle + h(N, \text{dtr } k) - \text{tr } h N(\text{tr } k),$$

and skew-symmetrize. To begin, write $\langle \nabla_N k, h \rangle = \langle (\nabla_N k)(N), h(N) \rangle + \langle (\nabla_N k)(e_i), h(e_i) \rangle$, so that $\langle \nabla_N k, h \rangle = N(k_{00})h_{00} + \phi_h[N(\text{tr } k) - N(k_{00})] + 2\langle (\nabla_N k)(N), h(N)^T \rangle$, where we have used the relation $\text{tr}_\gamma \nabla_N k = \text{tr}_N \nabla_N k - N(k_{00})$. Thus, $B(k, h)$ equals

$$\begin{aligned} & N(k_{00})h_{00} + \phi_h[N(\text{tr } k) - N(k_{00})] + 2\langle (\nabla_N k)(N), h(N)^T \rangle \\ & - N(\text{tr } k)[\text{tr } h - h_{00}] + \langle h(N)^T, \text{dtr } k \rangle. \end{aligned} \quad (3.21)$$

By (3.18) and (3.20),

$$2\langle (\nabla_N k)(N), h(N)^T \rangle = -2\langle \text{d}\phi_k, h(N)^T \rangle - 2\alpha(k, h) = -2\phi_k \delta_{\partial \mathcal{M}}(h(N)^T) - 2\alpha(k, h) + \omega_1$$

where ω_1 is a divergence term and $\alpha(k, h) = \langle \alpha(k(N)), h(N)^T \rangle$. Similarly, by (3.19) and (3.20),

$$N(k_{00}) = \delta_{\partial \mathcal{M}}(k(N)^T) + (\phi_k - k_{00})H,$$

where here and in the following $\delta = \delta_{\partial \mathcal{M}}$. Note also that $\langle h(N)^T, \text{dtr } k \rangle = \text{tr } k \delta_{\partial \mathcal{M}}(h(N)^T) + \omega_2$, where ω_2 is another divergence term. Since (3.17) involves integration over $\partial \mathcal{M}$, in the following we ignore the divergence terms. Substituting these computations in (3.21) gives

$$\begin{aligned} & \delta(k(N)^T)[h_{00} - \phi_h] + \delta(h(N)^T)[\text{tr } k - 2\phi_k] - (n-1)\phi_h N(\text{tr } k) \\ & + H(\phi_k - k_{00})(h_{00} - \phi_h) - 2\alpha(h, k). \end{aligned}$$

When skew-symmetrizing, the last two terms $H(\phi_k - k_{00})(h_{00} - \phi_h) - 2\alpha(h, k)$ cancel, while the first three terms combine to give

$$-(n-1)[\phi_h \delta(k(N)^T) - \phi_k \delta(h(N)^T)] - (n-1)[N(\text{tr } k)\phi_h - N(\text{tr } h)\phi_k],$$

or equivalently, (after dividing by $n-1$),

$$-\phi_h[N(\text{tr } k) + \delta(k(N)^T)] + \phi_k[N(\text{tr } h) + \delta(h(N)^T)]. \quad (3.22)$$

On the other hand, by (3.6) or (3.7),

$$\begin{aligned} 2(H')_k &= \text{tr}[\nabla_N k - 2\delta^*(k(N)^T) - \delta^*(k_{00}N)] \\ &= N(\text{tr } k) + 2\delta(k(N)^T) - k_{00}H - N(k_{00}). \end{aligned}$$

Substituting (3.19) gives

$$2(H')_k = N(\text{tr } k) + \delta(k(N)^T) - H\phi_k,$$

so (3.22) becomes

$$-\phi_h[2(H')_k + H\phi_k] + \phi_k[2(H')_h + H\phi_h] = -2\phi_h(H')_k + 2\phi_k(H')_h.$$

This vanishes exactly when H'_k and H'_h vanish. This completes the proof. \square

The main step in the proof of the manifold theorem, (theorem 3.6), is the following result.

Theorem 3.3. *Suppose $\pi_1(M, \partial M) = 0$ and $m \geq 3$. Then at any $\tilde{g} \in \mathbb{E}_S$, the map $\hat{\Phi}$ is a submersion, i.e. the derivative*

$$(D\hat{\Phi})_{\tilde{g}} : T_{\tilde{g}}\text{Met}_D^{m,\alpha}(\mathcal{M}) \rightarrow T_{\hat{\Phi}(\tilde{g})}S_\delta^{m-2,\alpha}(\mathcal{M}) \tag{3.23}$$

is surjective and its kernel splits in $T_{\tilde{g}}\text{Met}_D^{m,\alpha}(\mathcal{M})$.

Proof. The operator $\hat{L} = 2D\hat{\Phi}_{\tilde{g}}$ is elliptic in the interior, and the boundary data in lemma 3.2 give a self-adjoint elliptic boundary value problem. Let $S_B^{m,\alpha}(\mathcal{M})$ be the space of $C^{m,\alpha}$ symmetric bilinear forms on \mathcal{M} satisfying the boundary condition $B(h) = 0$ from lemma 3.2, i.e.

$$B(h) = \{\delta h, [h^T]_0, (H')_h\} = (0, 0, 0).$$

Clearly, $S_B^{m,\alpha}(\mathcal{M}) \subset S_D^{m,\alpha}(\mathcal{M})$, where $S_D^{m,\alpha}(\mathcal{M}) = T_{\tilde{g}}(\text{Met}_D^{m,\alpha}(\mathcal{M}))$. Throughout the following, we set $\tilde{g} = g$. The operator \hat{L} , mapping

$$S_B^{m,\alpha}(\mathcal{M}) \rightarrow S_\delta^{m-2,\alpha}(\mathcal{M}), \quad \hat{L}(h) = f, \quad B(h) = 0 \text{ at } \partial\mathcal{M},$$

is then Fredholm, of Fredholm index 0. On $S_B^{m,\alpha}(\mathcal{M})$, the image $\text{Im}(\hat{L})$ is a closed subspace of the range $S_\delta^{m-2,\alpha}(\mathcal{M})$, of finite codimension, and with codimension equal to dimension of the kernel K .

If $K = 0$, then \hat{L} maps $S_B^{m,\alpha}(\mathcal{M})$ onto $S_\delta^{m-2,\alpha}(\mathcal{M})$, which proves the result. Thus suppose $K \neq 0$. Then as in (3.15), by the self-adjointness, one has for any $h \in S_B^{m,\alpha}(\mathcal{M})$ and $k \in K$,

$$\int_{\mathcal{M}} \langle \hat{L}(h), k \rangle = \int_{\mathcal{M}} \langle h, \hat{L}(k) \rangle = 0,$$

since the boundary terms vanish and $\hat{L}(k) = 0$. Thus $\text{Im}(\hat{L}|_{S_B^{m,\alpha}(\mathcal{M})}) = K^\perp$, (where K^\perp is taken with respect to the L^2 inner product). To prove surjectivity on $S_D^{m,\alpha}(\mathcal{M})$, it thus suffices to prove that for any $k \in K$, there exists $h \in S_D^{m,\alpha}(\mathcal{M})$ such that

$$\int_{\mathcal{M}} \langle \hat{L}(h), k \rangle \neq 0. \tag{3.24}$$

Suppose then (3.24) does not hold, so that

$$\int_{\mathcal{M}} \langle \hat{L}(h), k \rangle = 0, \tag{3.25}$$

for all $h \in S_D^{m,\alpha}(\mathcal{M})$, i.e. for which $\delta h = 0$ on $\partial\mathcal{M}$. Integrating by parts, it follows that

$$\int_{\mathcal{M}} \langle h, \hat{L}(k) \rangle + \int_{\partial\mathcal{M}} Z(h, k) = 0, \tag{3.26}$$

for $Z(h, k)$ as following (3.16). As before, the boundary terms at infinity vanish, since $\delta > \frac{1}{2}$.

Choosing $h \in S_D^{m,\alpha}(\mathcal{M})$ arbitrary of compact support in \mathcal{M} , it follows from (3.26) that

$$\hat{L}(k) = 0, \tag{3.27}$$

i.e. k is formally tangent to $\hat{Z} = \hat{\Phi}^{-1}(0)$. Of course this is already known, since $k \in K$. Moreover, one also has

$$\delta k = 0 \text{ on } \mathcal{M}. \tag{3.28}$$

To see this, let $h = \delta^*V$, with V any vector field vanishing on $\partial\mathcal{M}$. Since g is Einstein and so $(\text{Ric} - \frac{s}{2}g)'_{\delta^*V} = 0$, it follows from (3.5) and (3.6) that $\hat{L}(h) = \delta^*Y$, where $Y = 2\delta\delta^*V$. As in lemma 2.2, the operator $\delta\delta^*$ is surjective, (in fact an isomorphism), on vector fields vanishing at $\partial\mathcal{M}$, so that Y may be arbitrarily prescribed. Moreover, $h \in S_D^{m,\alpha}(\mathcal{M})$ if and only if $Y = 0$ at $\partial\mathcal{M}$. Then (3.25) gives

$$0 = \int_{\mathcal{M}} \langle \hat{L}(\delta^*V), k \rangle = \int_{\mathcal{M}} \langle \delta^*Y, k \rangle = \int_{\mathcal{M}} \langle Y, \delta k \rangle + \int_{\partial\mathcal{M}} k(Y, N) = \int_{\mathcal{M}} \langle Y, \delta k \rangle,$$

since $Y = 0$ on $\partial\mathcal{M}$. Here we have used again the fact that the boundary term at infinity vanishes, since $|k| = O(r^{-\delta})$ and $|Y| = O(r^{-1-\delta})$. Since Y is otherwise arbitrary, this gives (3.28).

Returning now to (3.26), (3.27) gives

$$\int_{\partial\mathcal{M}} Z(h, k) = 0, \tag{3.29}$$

for all h with $\delta h = 0$ on $\partial\mathcal{M}$. Next, we choose certain test forms $h \in S_D(\mathcal{M})$ in (3.29). First, choose h such that $h = 0$ on $\partial\mathcal{M}$. Then $\nabla_N h$ is freely specifiable, subject to the divergence constraint $\delta h = 0$; all computations here and below are at $\partial\mathcal{M}$. Since $h = 0$, this constraint gives $(\nabla_N h)(N) = 0$, which is equivalent to the tangential and normal constraints:

$$(\nabla_N h)(N, T) = 0, \tag{3.30}$$

$$N(h_{00}) = 0, \tag{3.31}$$

for any T tangent to $\partial\mathcal{M}$. Choosing h and $\nabla_N h$ satisfying $h = 0$ and (3.30) and (3.31) at $\partial\mathcal{M}$, the terms $(\nabla_N h)(T_1, T_2)$ are freely specifiable on $\partial\mathcal{M}$, where T_1, T_2 are any vectors tangent to $\partial\mathcal{M}$. Substituting such h in (3.29) and using (3.28), it follows that

$$\int_{\partial\mathcal{M}} \langle \nabla_N h, k \rangle + (k_{00} - \text{tr} k)N(\text{tr} h) = 0. \tag{3.32}$$

Now choose $\nabla_N h = fg^T$, where $g^T = g|_{T(\partial\mathcal{M})}$. This choice satisfies the constraints (3.30) and (3.31). The integrand in (3.32) then becomes $f \text{tr}^T k - N(\text{tr} h) \text{tr}^T k$. Since $N(\text{tr} h) = \langle \nabla_N h, g \rangle = nf$, and since f is arbitrary, it follows that $\text{tr}^T k = 0$. In turn, since the tangential part of $\nabla_N h$ is arbitrary, (3.32) implies

$$k^T = 0, \text{ on } \partial\mathcal{M}. \tag{3.33}$$

Lemma 3.4. *At $\partial\mathcal{M}$, one has*

$$(A'_k)^T = 0, \tag{3.34}$$

i.e. $(\nabla_N k)^T = 2[\delta^*(k(N)^T)]^T + k_{00}A$, since $k^T = 0$, cf (3.6).

Proof. The proof is a straightforward, but rather long computation. To begin, as preceding (3.21) and using (3.33), one has $\langle \nabla_N h, k \rangle = 2\langle (\nabla_N h)(N), k(N)^T \rangle + N(h_{00})k_{00}$. By (3.18), $(\nabla_N h)(N)^T = \delta_{\partial\mathcal{M}}(h^T) - \alpha(h(N))$, so that

$$\begin{aligned} \int_{\partial\mathcal{M}} \langle \nabla_N h, k \rangle &= \int_{\partial\mathcal{M}} 2\langle \delta_{\partial\mathcal{M}}(h^T), k(N)^T \rangle + N(h_{00})k_{00} - 2\alpha(h, k) \\ &= \int_{\partial\mathcal{M}} 2\langle h^T, (\delta_{\partial\mathcal{M}})^*(k(N)^T) \rangle + N(h_{00})k_{00} - 2\alpha(h, k). \end{aligned} \tag{3.35}$$

Further, for Z tangent to $\partial\mathcal{M}$, one has $(\delta_{\partial\mathcal{M}})^*(k(N)^T)(Z, Z) = \langle \nabla_Z^T k(N)^T, Z \rangle = \langle \nabla_Z k(N)^T, Z \rangle = \delta^*(k(N)^T)(Z, Z)$, where now δ^* is taken with respect to the ambient metric $g_{\mathcal{M}}$, (not the boundary metric $\gamma_{\mathcal{M}}$). So this gives

$$\int_{\partial\mathcal{M}} \langle \nabla_N h, k \rangle = \int_{\partial\mathcal{M}} \langle h^T, 2\delta^*(k(N)^T) \rangle + N(h_{00})k_{00} - 2\alpha(h, k). \quad (3.36)$$

On the other hand, one computes $\langle \nabla_N k, h \rangle = \langle (\nabla_N k)^T, h^T \rangle + \langle \nabla_N k(N), h(N) \rangle = \langle (\nabla_N k)^T, h^T \rangle - \langle \alpha(k(N)), h(N)^T \rangle + N(k_{00})h_{00}$, again by (3.18) and (3.33). Taking the difference of this with (3.36) and noting that α is symmetric, gives

$$\int_{\partial\mathcal{M}} \langle h^T, (\nabla_N k)^T - 2\delta^*(k(N)^T) \rangle + N(k_{00})h_{00} - N(h_{00})k_{00} = E, \quad (3.37)$$

where via (3.16) and (3.17), E is given by

$$E = \int_{\partial\mathcal{M}} [k(N, d \operatorname{tr} h) - h(N, d \operatorname{tr} k)] - [N(\operatorname{tr} h)\operatorname{tr} k - \operatorname{tr} hN(\operatorname{tr} k)].$$

Computing this term-by-term gives: $k_{00}N(\operatorname{tr} h) + \langle k(N)^T, d^T \operatorname{tr} h \rangle - h_{00}N(\operatorname{tr} k) - \langle h(N)^T, d^T \operatorname{tr} k \rangle - N(\operatorname{tr} h)\operatorname{tr} k + \operatorname{tr} hN(\operatorname{tr} k)$. Since $\operatorname{tr} k = k_{00}$, the first and second-to-last terms cancel. Integrating over $\partial\mathcal{M}$ and using the divergence theorem shows that

$$E = \int_{\partial\mathcal{M}} \operatorname{tr} h \delta_{\partial\mathcal{M}}(k(N)^T) - k_{00} \delta^T(h(N)^T) - h_{00}N(\operatorname{tr} k) + \operatorname{tr} hN(\operatorname{tr} k). \quad (3.38)$$

Next we claim that

$$\delta_{\partial\mathcal{M}}(h(N)^T) = N(h_{00}) + Hh_{00} - \langle A, h \rangle, \quad (3.39)$$

and similarly for k . This follows from the following computation: $\delta_{\partial\mathcal{M}}(h(N)^T) = \delta_{\partial\mathcal{M}}(h(N)) - \delta^T(h_{00}N) = \delta_{\partial\mathcal{M}}(h(N)) + Hh_{00}$, while $\delta_{\partial\mathcal{M}}(h(N)) = \delta(h(N)) + N(h_{00})$. Since $\delta(h(N)) = (\delta h)(N) - \langle A, h \rangle$, this gives the claim. Substituting (3.39) into (3.38), and using $\langle A, k \rangle = 0$ implies that

$$E = \int_{\partial\mathcal{M}} \operatorname{tr} h(N(k_{00}) + Hk_{00}) - k_{00}(N(h_{00}) + Hh_{00} - \langle A, h \rangle) - h_{00}N(\operatorname{tr} k) + \operatorname{tr} hN(\operatorname{tr} k),$$

and rearranging terms gives

$$E = \int_{\partial\mathcal{M}} \langle A, h \rangle k_{00} + N(\operatorname{tr} k)[\operatorname{tr} h - h_{00}] + \operatorname{tr} hN(k_{00}) - k_{00}N(h_{00}) + H(\operatorname{tr} hk_{00} - \operatorname{tr} kh_{00}). \quad (3.40)$$

Now substitute (3.40) into (3.37): the $k_{00}N(h_{00})$ term cancels to give

$$\begin{aligned} & \int_{\partial\mathcal{M}} \langle h^T, (\nabla_N k)^T - 2\delta^*(k(N)^T) \rangle - \langle A, h \rangle k_{00} \\ &= - \int_{\partial\mathcal{M}} N(k_{00})h_{00} - N(\operatorname{tr} k)[\operatorname{tr} h - h_{00}] - \operatorname{tr} hN(k_{00}) - H(\operatorname{tr} hk_{00} - \operatorname{tr} kh_{00}). \end{aligned} \quad (3.41)$$

The integrand on the right combines to: $-N(k_{00})(h_{00} - \operatorname{tr} h) - N(\operatorname{tr} k)[h_{00} - \operatorname{tr} h] - H \operatorname{tr} k(h_{00} - \operatorname{tr} h) = -[N(k_{00}) + N(\operatorname{tr} k) + H \operatorname{tr} k](h_{00} - \operatorname{tr} h)$. Since $h_{00} - \operatorname{tr} h = -\operatorname{tr}^T h = -\langle h^T, g^T \rangle$, and since h^T may be chosen arbitrarily, (the constraint $\delta h = 0$ imposes no constraint on h^T), it follows that

$$(\nabla_N k)^T = 2[\delta^*(k(N)^T)]^T + k_{00}A + [N(k_{00}) + N(\operatorname{tr} k) + H \operatorname{tr} k]g^T. \quad (3.42)$$

To complete the proof of (3.34), we thus need to show that

$$N(k_{00}) + N(\operatorname{tr} k) + H \operatorname{tr} k = 0. \quad (3.43)$$

To obtain (3.43), take the g^T -trace of (3.42). One has $\langle \nabla_N k, g^T \rangle = N(\text{tr } k) - N(k_{00})$, while $\langle \delta^*(k(N)^T), g^T \rangle = \langle \nabla_{e_i} k(N)^T, e_i \rangle = \langle \nabla_{e_i} k(N), e_i \rangle - k_{00}H = \langle (\nabla_{e_i} k)(N), e_i \rangle - k(\nabla_{e_i} N, e_i) - k_{00}H = -N(k_{00}) - k_{00}H$, the last equality using (3.33) and (3.28). This gives

$$N(\text{tr } k) - N(k_{00}) = -2N(k_{00}) - 2k_{00}H + k_{00}H - n[N(k_{00}) + N(\text{tr } k) + H\text{tr } k],$$

which implies (3.43). This completes the proof of the lemma. □

To complete the proof of theorem 3.3, (3.33) and (3.34) show that

$$k^T = (A'_k)^T = 0,$$

at $\partial\mathcal{M}$. One also has $\hat{L}(k) = \delta k = 0$ on \mathcal{M} , so that k is an infinitesimal Einstein deformation on \mathcal{M} . By the local unique continuation result of [AH], together with the global hypothesis $\pi_1(M, \partial M) = 0$, it follows that $k = 0$. This shows that \hat{L} is surjective. The fact that its kernel splits is standard, cf [A3]. This completes the proof. □

Via the implicit function theorem, one obtains:

Corollary 3.5. *Suppose $\pi_1(M, \partial M) = 0$ and $m \geq 3$. Then the local spaces $\mathbb{E}_D^{m,\alpha}$ are infinite dimensional C^∞ Banach manifolds, with*

$$T_{\tilde{g}}\mathbb{E}_D = \text{Ker}(D\hat{\Phi}_{\tilde{g}}). \tag{3.44}$$

Proof. This is an immediate consequence of theorem 3.3, the fact from proposition 2.1 that $\mathbb{E}_D = Z_D$, (cf (2.18)), and the implicit function theorem, (or regular value theorem), in Banach spaces. □

This leads to the main result of this section.

Theorem 3.6. *Suppose $\pi_1(M, \partial M) = 0$ and $m \geq 3$. Then the moduli space $\mathcal{E}_S = \mathcal{E}_S^{m,\alpha}$ is a C^∞ smooth infinite dimensional Banach manifold for which the boundary map*

$$\Pi_B : \mathcal{E}_S \rightarrow \text{Met}^{m,\alpha}(\partial M) \times C^{m-1,\alpha}(\partial M), \tag{3.45}$$

is a C^∞ smooth Fredholm map, of Fredholm index 0.

Proof. Recall from section 1 that the moduli space \mathcal{E}_S of static vacuum Einstein metrics is defined to be the quotient $\mathbb{E}_S^{m,\alpha} / \mathcal{D}_1^{m+1,\alpha}$. The local spaces \mathbb{E}_D are smooth Banach manifolds and depend smoothly on the background metric \tilde{g} , since the divergence-free gauge condition (2.18) varies smoothly with \tilde{g} . As noted preceding lemma 2.2, the action of \mathcal{D}_1 on \mathbb{E} is free and the local spaces \mathbb{E}_D are smooth local slices for the action of \mathcal{D}_1 on \mathbb{E}_S . Hence the global space \mathbb{E}_S is a smooth Banach manifold, as is the quotient \mathcal{E}_S . The local slices \mathbb{E}_D represent local coordinate patches for \mathcal{E}_S .

Proposition 3.1 implies that the boundary map $\Pi_B : \mathbb{E}_S^{m,\alpha} \rightarrow \text{Met}^{m,\alpha}(\partial M) \times C^{m-1,\alpha}(\partial M)$ is smooth and Fredholm, of Fredholm index 0. Moreover, Π_B is invariant under the action of $\mathcal{D}_1^{m+1,\alpha}(M)$ on $\mathcal{E}_S^{m,\alpha}$ and so it descends to a smooth Fredholm map as in (3.45), still of index 0. □

The boundary conditions (γ, H) for the operator $\hat{\Phi}$ are also self-adjoint; in fact they arise naturally from a variational principle (Lagrangian) on a space of static (non-vacuum) metrics.

To describe this, let $S(h) = -\Delta \text{tr } h + \delta\delta(h) - \langle \text{Ric}, h \rangle$ be the linearization of the scalar curvature s , with adjoint S^* given by

$$S^*u = D^2u - \Delta u g - u \text{Ric}.$$

It is well-known that the static vacuum equations are given by $S^*u = 0$ and $s = 0$ on M .

Proposition 3.7. For (M, g, u) as above, the Bartnik boundary conditions (γ, H) give a well-defined variational problem for the Lagrangian

$$\mathcal{L}(g, u) = \int_M usdV_g - 16\pi m_{\text{ADM}} : \text{Met}_\delta^{m,\alpha}(M) \times C_\delta^{m,\alpha}(M) \rightarrow \mathbb{R}, \quad (3.46)$$

where m_{ADM} is the ADM mass of (M, g, u) . The gradient $\nabla\mathcal{L}$ of \mathcal{L} at (g, u) is given by

$$\nabla\mathcal{L} = \left(S^*u + \frac{1}{2}usg, s, uA - N(u)\gamma, 2u \right), \quad (3.47)$$

in the following sense: if (h, u') is a variation of (g, u) inducing the variation (h^T, H'_h) of the boundary data, then

$$d\mathcal{L}(h, u', h^T, H'_h) = \int_M \left[\left\langle S^*u + \frac{1}{2}usg, h \right\rangle + su' \right] + \int_{\partial M} [uA - N(u)\gamma, h^T] + 2uH'_h. \quad (3.48)$$

In particular, the static vacuum equations are critical points for \mathcal{L} with data (γ, H) fixed.

Proof. Suppose that D is compact domain in M , with N the outward unit normal from D . Varying (g, u) in the direction (h, u') then gives

$$D\mathcal{L}_0(h, u') = \int_D (us' + u's + us(dV)') = \int_D \langle u, s(h) \rangle + \frac{1}{2}us\langle g, h \rangle + su', \quad (3.49)$$

where $\mathcal{L}_0 = \int_D usdV_g$. A straightforward integration by parts gives

$$\int_D \langle S(h), u \rangle = \int_D \langle h, S^*u \rangle + \int_{\partial D} -uN(\text{tr } h) - (\delta h)(N)u - \langle h(N), du \rangle + \text{tr } hN(u). \quad (3.50)$$

The equations (3.49) and (3.50) imply immediately the bulk Euler-Lagrange equations—the first two terms in (3.47). If the bulk term (over D) vanishes, then since u' is arbitrary $s = 0$, so this gives

$$S^*u = 0,$$

with $s = 0$, which are the static vacuum equations.

For the boundary terms, from (3.7) one has

$$2H'_h = N(\text{tr } h) + 2\delta(h(N)^T) - h_{00}H - N(h_{00}).$$

Also by a simple calculation

$$(\delta h)(N) = \delta(h(N)^T) + \langle A, h \rangle - h_{00}H - N(h_{00}),$$

so that,

$$2H'_h - (\delta h)(N) = N(\text{tr } h) + \delta(h(N)^T) - \langle A, h \rangle.$$

This gives

$$\int_{\partial D} u(-N(\text{tr } h) - (\delta h)(N)) = \int_{\partial D} -2uH'_h + \langle du, h(N)^T \rangle - u\langle A, h \rangle.$$

It follows that the boundary term in (3.50) is given by

$$\int_{\partial D} -2uH'_h - u\langle A, h \rangle - N(u)h_{00} + N(u)\text{tr } h = \int_{\partial D} -2uH'_h - u\langle A, h^T \rangle + N(u)\langle h^T, \gamma \rangle. \quad (3.51)$$

Now let the outer boundary of D equal $S(r)$ and consider the limit $r \rightarrow \infty$. Then $u \rightarrow 1$ and $\int_{S(r)} N(u)\langle h^T, \gamma \rangle \rightarrow 0$. It follows that

$$\lim_{r \rightarrow \infty} \int_{S(r)} -2uH'_h - u\langle A, h^T \rangle = \lim_{r \rightarrow \infty} \int_{S(r)} (-N(\text{tr } h) - (\delta h)(N)) = 16\pi(m_{\text{ADM}})',$$

where the second equality follows from standard formulas for the ADM mass and its variation, cf [RT, B3]. Here the variation $(m_{\text{ADM}})'$ is taken in the direction h . The formula (3.50) and (3.51) is also valid at the inner boundary ∂M , with respect to the outward normal. Changing to the inner normal changes the sign of each term, and (3.47) and (3.48) then follow immediately. \square

On-shell, i.e. on the space of solutions \mathcal{E} , the Lagrangian

$$\mathcal{L} = -16\pi m_{\text{ADM}} : \mathcal{E} \rightarrow \mathbb{R}$$

is a smooth function whose derivative is given by the boundary term in (3.48), a result basically due to Bartnik [B3]. After writing this work, we learned that a special case of proposition 3.7 has been noted, without proof, in a paper of Miao, cf [M2].

4. Curvature estimates and properness of Π^o

In contrast to the previous section, where most of the computations were done on the 4-manifold $(\mathcal{M}, g_{\mathcal{M}})$, in this section we focus mostly on the three-dimensional data (M, g, u) . Let $\text{inj}_{\partial M}$ denote the injectivity radius of the normal exponential map from ∂M in M and let R denote the (full) curvature tensor of g . The main result of this section is the following collection of *a priori* estimates.

Theorem 4.1. *For $(M, g, u) \in \mathcal{E}^o = (\mathcal{E}^{m,\alpha})^o$ with $m \geq 2$, one has a global pointwise estimate*

$$|R| \leq \Lambda, \tag{4.1}$$

on M , where Λ depends only on bounds for the boundary data (γ, H) . Moreover, at ∂M , one has the bounds

$$|A| \leq \Lambda, \quad \text{inj}_{\partial M} \geq \Lambda^{-1}. \tag{4.2}$$

The estimates (4.1), (4.2) also hold for higher derivatives of R and A , up to order $m - 2$, $m - 1$ respectively.

Proof. For points in the interior of M , of bounded distance away from ∂M , this follows directly from the *a priori* interior estimates in [A1] which state

$$|R|(x) \leq \frac{K}{t^2(x)}, \quad |\text{d log } u|(x) \leq \frac{K}{t(x)}, \tag{4.3}$$

where $t(x) = \text{dist}(x, \partial M)$, where K is an absolute constant. Similar (scale-invariant) estimates hold for all higher derivatives of R and $\log u$. So one only needs to consider the behavior near ∂M . At ∂M , the Gauss and Gauss–Codazzi (constraint) equations are given by:

$$|A|^2 - H^2 + s_\gamma = -2R_{NN}, \tag{4.4}$$

$$\delta(A - H\gamma) = -u^{-1}D^2u(N, \cdot). \tag{4.5}$$

Also, $-R_{NN} = -\text{Ric}(N, N) = -u^{-1}NN(u) = u^{-1}(\Delta_{\partial M}u + HN(u))$, so that

$$u(|A|^2 - H^2 + s_\gamma) = 2(\Delta_{\partial M}u + HN(u)). \tag{4.6}$$

From (4.4), a bound on $|R|$ at ∂M gives immediately a bound on $|A|$ at ∂M , given control of (γ, H) . Similarly, a bound on $|R|$ on M gives a lower bound on the distance d_{con} to the conjugacy locus of the normal exponential map $\exp_{\partial M}$.

Now, again under a bound on $|R|$, the outer-minimizing property (1.8) implies a lower bound on the distance $\delta_{\partial M}$ to the cut locus of $\exp_{\partial M}$. To see this, suppose that $\delta_{\partial M} \ll 1$ but Λ in (4.1) is bounded, $\Lambda \sim 1$. Then since d_{con} is bounded below, there is a geodesic ζ of length

$2\delta_{\partial M}$ in M meeting ∂M orthogonally at points p_1, p_2 . Let T be the boundary of the tubular neighborhood of ζ of radius r . Then T intersects ∂M in the boundary of two discs D_1, D_2 of radius approximately r , (for r small). If $\delta_{\partial M} \ll r$, then $\text{area } T < \text{area}(D_1 \cup D_2)$. Further, $T \subset M$ is homologous to $D_1 \cup D_2$ in M . Removing then $D_1 \cup D_2$ from ∂M and attaching T shows that ∂M is not outer-minimizing in M , giving a contradiction.

Thus it suffices to obtain a curvature bound at or arbitrarily near ∂M . The higher derivative estimates may then be obtained by standard elliptic regularity methods. The proof of (4.1) is by a blow-up argument. If the curvature bound in (4.1) is false, then there is a sequence $(M, g_i, u_i, x_i) \in \mathcal{E}^o$ with bounded Bartnik boundary data such that

$$|R_{g_i}|(x_i) \rightarrow \infty.$$

Without loss of generality, assume that the curvature of g_i is maximal at x_i . We then rescale the metrics g_i to g'_i so that $|R|$ is bounded, and equals 1 at x_i ,

$$|R_{g'_i}|(x_i) = 1, \quad |R_{g'_i}|(y_i) \leq 1, \tag{4.7}$$

for any $y_i \in (M, g'_i)$. Thus define $g'_i = \lambda_i^2 g_i$ where $\lambda_i = |R_{g_i}|(x_i)$. This gives (4.7) as well as $H_{g'_i} = \lambda_i^{-1} H_{g_i}$, $s_{g'_i} = \lambda_i^{-2} s_{g_i}$ and $t'_i = \lambda_i t_i$. Note that (4.3) implies that $t'_i(x_i) \leq \sqrt{K}$ so that x_i remains within a uniformly bounded distance to the boundary ∂M with respect to g'_i .

One may also need to rescale the potential u . For reasons that will be clearer below, choose points $y_i \in M$ such that $\text{dist}_{g'_i}(y_i, \partial M) = 1$ and $\text{dist}_{g'_i}(y_i, x_i) \leq \sqrt{K}$, and rescale u_i so

$$u'_i(y_i) = 1. \tag{4.8}$$

The sequence (M, g'_i, u'_i) has uniformly bounded curvature and uniform control of the boundary geometry, (boundary metric, second fundamental form and normal exponential map). By (4.8) and the Harnack inequality for positive harmonic functions, the potential u'_i is also uniformly bounded in compact sets. It follows from the convergence theorem in [AT] for manifolds-with-boundary that a subsequence converges weakly, (i.e. in $C^{1,\alpha}$), to a $C^{1,\alpha}$ static limit (X, g, u, x) with boundary $(\partial X, \gamma, u)$. The convergence is uniform on compact subsets. More precisely, given any smooth compact domain Ω in the manifold with boundary X , there is a subsequence, also denoted $\{i\}$ and embeddings $F_i : \Omega \rightarrow (M_i, g'_i, u'_i)$ such that $F_i^*(g'_i, u'_i) \rightarrow (\Omega, g, u)$ in the $C^{1,\alpha}$ topology. Moreover, for $\Omega_1 \subset \Omega_2$, $g_2|_{\Omega_1} = g_1$. For the proof, we refer to [AT, theorem 3.1], and more precisely to the local version of this result in [AT, theorem 3.1.1].

By the normalization in (4.7), the limit (X, g) is complete (without singularities) up to the boundary ∂X . Since ∂M is outer-minimizing in (M, g_i) , the C^0 convergence to the limit implies that ∂X is weakly outer-minimizing in X : if D is any compact smooth domain in ∂X and D' is a surface in X with $\partial D' = \partial D$, then

$$\text{area } D' \geq \text{area } D. \tag{4.9}$$

One has $\partial X = \mathbb{R}^2$, the boundary metric γ is flat, $H = 0$, so ∂X is a minimal surface in X . One has $u > 0$ in the interior of X , (by the maximum principle), but may have $u = 0$ somewhere or everywhere on ∂X . The bound (4.7) and the static equations imply that u_i is uniformly bounded up to ∂X , within bounded distance to x_i and the limit potential u extends at least $C^{1,\alpha}$ up to ∂X .

We will prove below that the convergence to the limit is smooth, so that in particular

$$|R|(x) = 1, \tag{4.10}$$

where $x = \lim x_i$ and $R = R_X$.

On the blow-up limit (X, g) , (4.6) holds and becomes

$$\frac{1}{2}u|A|^2 = \Delta_{\partial M}u,$$

on ∂X . This equation holds weakly on ∂X with $u \in C^{1,\alpha}(\partial X)$; elliptic regularity then implies it holds strongly, and $u \in C^{3,\alpha}(\partial X)$. Since u is harmonic, u is thus $C^{3,\alpha}$ up to ∂X . Also, by the Riccati equation $N(H) = -|A|^2 - R_{NN} = -|A|^2 + \frac{1}{2}(|A|^2 - H^2 + s_\gamma)$, so that

$$N(H) = -\frac{1}{2}(|A|^2 + H^2 - s_\gamma). \tag{4.11}$$

This holds pointwise on the blow-up sequence (M, g'_i, u_i) and since $s_\gamma \rightarrow 0$ and $H \rightarrow 0$ for g'_i , it follows that $N(H)$ is defined pointwise on the limit ∂X and on ∂X ,

$$N(H) \leq 0, \tag{4.12}$$

with equality on any domain only when $A = 0$.

Since ∂X is minimal, (4.12) and the outer-minimizing property (4.9) imply that

$$N(H) = 0, \tag{4.13}$$

on ∂X . In more detail, (4.9) and the fact that $H = 0$ on ∂X implies the second order stability of ∂X , in that the second variation of the area of ∂X is non-negative. Thus, for all f of compact support on ∂X , one has

$$\int_{\partial X} (|df|^2 + f^2 N(H)) \geq 0. \tag{4.14}$$

Choose $f = f_{R,S}(r)$ such that $f = 1$ on $D(R) \subset \partial X = \mathbb{R}^2$ and, for $r \geq R$, $f = (\log r - \log S)/(\log R - \log S)$, for $S \gg R \gg 1$. One may choose R and S sufficiently large such that $\int_{\partial X} |df|^2 < \varepsilon$, for any given $\varepsilon > 0$. This together with (4.12) implies (4.13).

It follows that $A = 0$ and hence by the Liouville theorem on \mathbb{R}^2 , $u = \text{const}$ on ∂X . Using the divergence constraint (4.5), we also now have $0 = \delta(A - H\gamma) = -u^{-1}D^2u(N, \cdot)$, and so $0 = D^2u(N, \cdot) = dN(u) - A(du) = dN(u)$, so that $N(u) = \text{const}$.

Thus, the full Cauchy data $(\gamma, u, A, N(u))$ for the static vacuum equations is fixed and trivial: γ is the flat metric, $A = 0$ and $u, N(u)$ are constant. Observe that this data is realized by the family of flat metrics on $(\mathbb{R}^3)^+$ with either $u = \text{const}$ or u equal to an affine function on $(\mathbb{R}^3)^+$.

Suppose first

$$u = \text{const} > 0 \text{ on } \partial X. \tag{4.15}$$

The static vacuum equations (1.1) are then non-degenerate up to ∂X . The unique continuation property for Einstein metrics with boundary, cf [AH], implies that the Cauchy data uniquely determine the solution locally. Alternately, since the static vacuum equations are non-degenerate up to ∂X and since the boundary data $(\partial X, \gamma, H)$ are real-analytic, elliptic regularity implies that the solution (M, g, u) is real-analytic up to ∂M . Such solutions are uniquely determined (locally) by their Cauchy data. Hence, the limit (X, g, u) is flat in this case.

Moreover, the convergence to the limit is smooth everywhere. This again follows from non-degeneracy and ellipticity. Briefly, the potential equation $\Delta u = 0$ gives a boost on the regularity of u , (given background regularity on g). One substitutes this into the main static vacuum equation $u\text{Ric} = D^2u$, giving thus a boost to the regularity of Ric , inducing then a boost to the regularity of g . This in turn further boosts the regularity of u via the potential equation. Bootstrapping gives $C^{m,\alpha}$ convergence, up to the boundary, in regions where $u > 0$, (given that u is $C^{m,\alpha}$ at the boundary). In sum, one has a contradiction to (4.10).

Thus, suppose instead

$$u = 0 \text{ on } \partial X. \tag{4.16}$$

This situation is more complicated. It is also more difficult to prove smooth convergence in this situation (one may have x in (4.10) at ∂X). Moreover, there are in fact non-flat static vacuum

solutions with flat Cauchy data as above with $u = 0$ on ∂X , (so-called toroidal black holes, cf [P] and [Th]). Thus the unique continuation results used above are false in this degenerate situation where the boundary becomes characteristic.

We observe first that the solution (X, g) is still real-analytic up to ∂X in this case. This follows since the 4-metric $g^4 = u^2 d\theta^2 + g_X$ is Einstein ($\text{Ric} = 0$) and is $C^{1,\alpha}$ up to the horizon or vanishing locus $\partial X = \{u = 0\}$. Elliptic regularity for the Einstein equations then implies that g^4 is real-analytic, and hence so are u, g_X up to ∂X .

To see this in more detail, let U be a chart neighborhood of ∂X in X , so that U is diffeomorphic to a half-ball in \mathbb{R}^3 with boundary a disc $D^2 \subset \mathbb{R}^2$. Over each $p \in U$, one has a circle of length $2\pi u(p)$, with $u \rightarrow 0$ as $p \rightarrow \partial X$. From the work above, u is $C^{1,\alpha}$ up to ∂X with $N(u) = \text{const}$ at ∂X . Note that $N(u) \neq 0$ at ∂X . For if $N(u) = 0$ at ∂X , since also $u = 0$ at ∂X and u is harmonic ($\Delta u = 0$), the unique continuation property for harmonic functions implies that $u = 0$ in X , giving a contradiction. By rescaling u if necessary, one may thus assume that $N(u) = 1$ at ∂X . This implies that the 4-metric g^4 , defined on $B^4 \setminus D^2$ extends to a $C^{1,\alpha}$ metric on the 4-ball B^4 . (The coordinate θ is an angular variable in \mathbb{R}^2 in polar coordinates, shrinking down to the origin on approach to ∂X). It is well-known, cf [Be], that any $C^{1,\alpha}$ weak solution to the Einstein equations is real-analytic (in harmonic or geodesic normal coordinates), which gives the claim above.

To prove the limit is in fact flat, and that one has strong convergence, we need to use the outer-minimizing property again. Thus, first note that (4.11) holds everywhere on the limit (X, g) near ∂X , not just at ∂X ; here A is the second fundamental form of the level sets $S(t)$ of $t = \text{dist}(\partial X, \cdot)$, etc. We have already established $N(H) = 0$ at ∂X , via the outer-minimizing property and the corresponding stability of the second variation operator (4.14). Taking then the derivative of (4.11) in the normal direction gives,

$$NN(H) = -\langle A', A \rangle - \langle A^2, A \rangle - H'H + \frac{1}{2}s'_\gamma, \tag{4.17}$$

where $A' = \nabla_N A$. At ∂X , the first three terms vanish while $(s'_\gamma)_k = -\Delta(\text{tr} k) + \delta\delta k - \langle \text{Ric}_\gamma, k \rangle = 0$, since $k = 2A = 0$. Thus

$$NN(H) = 0, \tag{4.18}$$

at ∂X and it follows that the third variation of the area of ∂X in the unit normal direction vanishes.

Now choose $f = f_{R,S}$ as following (4.14) with R, S large. Let $S_{tf} = \exp_{\partial X}(tf(x))$, where $\exp_{\partial X}$ is the normal exponential map of ∂X into X . Letting $v(t) = \text{area} S_{tf}$, one has

$$v(t) = v(0) + \frac{1}{2}v''(0)t^2 + \frac{1}{6}v'''(0)t^3 + \frac{1}{24}v''''(0)t^4 + O(t^5). \tag{4.19}$$

The expansion (4.19) is valid for all t sufficiently small, $|t| \leq \delta_0$, with δ_0 independent of R, S , since the area and its derivatives are integrals of local expressions, and the local geometry of X is uniformly bounded in a tubular neighborhood of radius 1 about ∂X . By the second variational formula (4.14) and (4.13), for any given $\varepsilon > 0$, one has

$$v''(0) \leq \varepsilon,$$

for R, S sufficiently large. For the same reasons via (4.18),

$$v'''(0) \leq \varepsilon.$$

It follows then from the outer-minimizing property (4.9) and (4.19) that for R, S sufficiently large, one must have

$$v''''(0) \geq -\varepsilon, \tag{4.20}$$

again for any $\varepsilon = \varepsilon(R, S) > 0$. Using the vanishing of the lower order terms, one computes that (4.20) gives

$$\int_{\partial X} f^4 NNN(H) - 6f^2 \langle df \cdot df, A' \rangle \geq -\varepsilon. \tag{4.21}$$

On the other hand, taking the normal derivative of (4.17) gives

$$NNN(H) = -\langle A', A' \rangle - (H')^2 + \frac{1}{2}s''_\gamma. \tag{4.22}$$

We have $H' = N(H) = 0$ and $(s'_\gamma)_{2A} = 2\Delta H + 2\delta\delta A - \langle \text{Ric}_\gamma, 2A \rangle$. For s''_γ , one has $(\Delta H)' = \Delta'H + \Delta H' = 0$ and $\langle \text{Ric}_\gamma, A' \rangle = \langle (\text{Ric}_\gamma)', A \rangle + \langle (\text{Ric}_\gamma), A' \rangle = 0$. So at ∂X , $\frac{1}{2}s''_\gamma = \delta\delta A'$. It follows then from (4.21) and (4.22) that for $f = f_{R,S}$ as above

$$\int_{\partial X} -f^4 |A'|^2 + f^4 \delta\delta A' - 6f^2 \langle df \cdot df, A' \rangle \geq -\varepsilon. \tag{4.23}$$

Integrating the second term by parts gives $\int \langle D^2 f^4, A' \rangle = \int \langle 4f^3 D^2 f + 12f^2 \langle df \cdot df, A' \rangle$. Using the Cauchy–Schwarz and Young inequalities, (4.23) then implies, for any μ small,

$$\int_{\partial X} f^4 |A'|^2 \leq \mu \int_{\partial X} f^4 |A'|^2 + C\mu^{-1} \int_{\partial X} f^2 |D^2 f|^2 + C\mu^{-1} \int_{\partial X} |df|^4 + \varepsilon.$$

Choosing μ small, the first term on the right may be absorbed into the left, while simple computation shows that the last two terms become arbitrarily small for R and S sufficiently large. It follows that

$$A' = 0 \tag{4.24}$$

and so $NNN(H) = 0$ at ∂X . The Riccati equation

$$A' = \nabla_N A = -A^2 - R_N, \tag{4.25}$$

where $R_N(V, W) = \langle R_g(V, N)N, W \rangle$, thus gives $R_N = 0$ at ∂X , and so via the Gauss and Gauss–Codazzi equations $R_g = 0$ at ∂X . Thus the full ambient curvature vanishes at ∂X .

One can now continue inductively in the same way to see that A and R vanish to infinite order at ∂X . A simpler method proceeds as follows. The Riccati equation (4.25) holds along the level sets $S(t)$ of $t = \text{dist}(\partial X, \cdot)$. Since $s_g = 0$ and $\dim X = 3$, $R_N = - * \text{Ric}$, i.e. $R_N(v, v) = -\text{Ric}(w, w)$, where (N, v, w) are an orthonormal basis. Via the static vacuum equations, this gives $\nabla_N A = -A^2 + u^{-1} * (D^2 u)$. Rescale u if necessary so that $N(u) = 1$ at ∂X and set $v = u - t$. Since $A = D^2 t$, one then obtains

$$\nabla_N A = -A^2 + u^{-1}(*A) + u^{-1}(*D^2 v). \tag{4.26}$$

This is a system of ODEs for A , singular at $\partial X = \mathbb{R}^2$, but with indicial root 1. From the work above, we have $v = O(t^2)$ and $A = O(t^2)$. Writing $A = t^2 B$ and substituting in (4.26) shows that $A = O(t^3)$. Also, by the computation following (4.22), $s_\gamma = O(t^3)$ on $S(t)$, and hence using the scalar constraint (4.4) and the relation $R_{NN} = u^{-1} NN(u)$, this in turn implies $v = O(t^3)$, and so on. It follows that (g, u) agree with a flat solution to infinite order at ∂X . Since the solution (X, g, u) is analytic up to ∂X , it follows that (X, g, u) is flat, as claimed.

Next we claim that one has strong convergence to the limit, so that (4.7) is preserved in the limit, i.e. (4.10) holds, contradicting the fact that the limit is flat. Note first that if x in (4.10) is in the interior of X , then strong (C^∞) convergence is immediate, by the interior estimates (4.3), i.e. their higher derivative analogues. Thus we may assume that $x \in \partial X$.

From the work above, we know that $|R|$ is uniformly bounded everywhere on (M, g'_i) and $|R| \rightarrow 0$ everywhere away from $\partial M \rightarrow \partial X$, so $|R|$ jumps quickly from 1 to 0 near x_i . The main point is to prove that

$$|R|(x) \rightarrow 0, \tag{4.27}$$

for all $x \in \partial M \rightarrow \partial X$; it is then easy to prove that $|R| \rightarrow 0$ on (M, g'_i) , cf (4.38) below. To prove (4.27), note that the estimates (4.11)–(4.24) above at ∂X also hold on the blow-up sequence at ∂M , (since $H \rightarrow 0$ and $s_{\gamma'} \rightarrow 0$ on ∂M). It follows then by these arguments that for any $R < \infty$ and $D(R)$ the R -ball about any base point $y_i \in (\partial M, \gamma'_i)$ converging to $y \in \partial X$,

$$\int_{D(R)} |A|^2 + |\nabla_N A|^2 \rightarrow 0. \tag{4.28}$$

To proceed further, consider the divergence constraint $\delta A = -dH - \text{Ric}(N, \cdot)$ on $(\partial M, \gamma'_i)$, as in (4.5). The equations $\delta A = \chi_1$, $d \text{tr} A = \chi_2$ form an elliptic first order system in 2-dimensions, and so one has elliptic estimates. Since $H \rightarrow 0 \in C^{m-1, \alpha}$ on $(\partial M, \gamma'_i)$, $dH \rightarrow 0$ in $C^{m-2, \alpha}$. Also, by assumption (4.7), $\text{Ric}(N, \cdot)$ is bounded in L^∞ . It follows then from elliptic regularity that A is bounded in $L^{1,p}$. By the scalar constraint (4.4), R_{NN} is then also bounded in $L^{1,p}$ and since $\text{tr} R_N = R_{NN}$, it follows from (4.28) and (4.25) that

$$A \rightarrow 0 \text{ and } R_{NN} \rightarrow 0 \text{ in } C^\alpha_{\text{loc}}(\partial M). \tag{4.29}$$

Next the Einstein equation $\text{Ric}_{g_{\mathcal{M}}} = 0$ on $(\mathcal{M}, g_{\mathcal{M}})$ implies that $\delta_{\mathcal{M}} R_{g_{\mathcal{M}}} = 0$. Hence

$$0 = (\delta_{g_{\mathcal{M}}} R)(N, N, \cdot) = \delta_{g_{\mathcal{M}}}(R(\cdot, N)N, \cdot) + 2R(e_\alpha, \nabla_{e_\alpha} N)N = \delta_{g_{\mathcal{M}}}(R(\cdot, N)N, \cdot), \tag{4.30}$$

since one may choose a basis in which $\nabla_{e_\alpha} N = A(e_\alpha) = \lambda_\alpha e_\alpha$. Let V be the unit vertical vector, and note that $\nabla_V V = -dv$, where $v = \log u$. Then $\nabla_V(R(V, N)N, \cdot) = R(dv, N)N$, while $\nabla_N(R(N, N)N, \cdot) = 0$. Hence, for R_N as in (4.25), these computations on ∂M give

$$\delta R_N = -R_N(dv),$$

where the divergence δ and R_N are taken on $(\partial M, \gamma'_i)$. Since R is bounded in L^∞ and $\text{tr} R_N = R_{NN}$ is bounded in $L^{1,p}$, elliptic regularity gives

$$\|R_N\|_{L^{1,p}} \leq C \|dv\|_{L^p}, \tag{4.31}$$

again on compact domains in ∂M converging to a compact domain in ∂X .

To control dv in (4.31), recall that by (4.7) the ambient curvature R is bounded, and so R restricted to ∂M is also bounded. Via the static vacuum equations (1.1), this implies that

$$N(v)A + u^{-1}D^2u \tag{4.32}$$

is bounded on $(\partial M, \gamma'_i)$, where D^2u is the Hessian of $u|_{\partial M} : \partial M \rightarrow \mathbb{R}^+$. Now we claim that each term in (4.32) is bounded, i.e. there exists K such that

$$|u^{-1}D^2u| \leq K, \quad |N(v)A| \leq K, \tag{4.33}$$

pointwise, on domains converging to a bounded domain in ∂X . To prove (4.33), suppose instead that $|u^{-1}D^2u| \rightarrow \infty$ at some sequence of base points $y_i \rightarrow y \in \partial X$. Without loss of generality, we may assume the points y_i realize the maximum of $|u^{-1}D^2u|$ on $D_{y_i}(10) \subset (\partial M, \gamma'_i)$ (possibly up to a factor of 2), where $D_{y_i}(10)$ is the geodesic disc of radius 10 in ∂M centered at y_i . As before, one may then rescale the metrics g'_i further to g''_i so that $|u^{-1}D^2u|(y_i) = 1$ and hence the full curvature $R \rightarrow 0$ in this scale. Also, renormalize u if necessary so that $u(y_i) = 1$. Note that $|du|(y_i)$ must be bounded. For if $|du|(y_i)$ were too large, it follows, (e.g. by a still further rescaling), that u would be close to an affine function on \mathbb{R}^2 and hence u would assume negative values in bounded distance to y_i . Since $u > 0$ everywhere, this is impossible. Thus, by integration along paths, u is bounded in $L^{1,\infty}$ in the scale g''_i , within bounded distance to y_i .

Now working in the scale and normalization above, from divergence constraint (4.5), one has $-u\delta(A - H\gamma) = dN(u) - A(du)$. Since A is bounded in $L^{1,p}$ and u and du are bounded in L^∞ , it follows that $dN(u)$ is bounded in L^p . Thus, $N(u) = c + \phi$, where ϕ is bounded in $L^{1,p}$. Here c is a constant which may, and in fact does, go to $\pm\infty$, in the u -normalization $u(y_i) = 1$ above.

The trace equation (4.6) in this scale and normalization gives

$$\Delta u + H(c + \phi) = uf, \tag{4.34}$$

where f is bounded in $L^{1,p}$. Also, Hc is bounded, since $N(u)A$ is bounded via (4.32), and so $Hc \rightarrow c'$, for some constant c' on ∂X . Since ϕ is also bounded in $L^{1,p}$, it follows that u is bounded in $L^{3,p}$, and hence (in a subsequence), u converges in $C^{2,\alpha}$ to its limit on $\partial X = \mathbb{R}^2$. Hence D^2u converges in C^α to its limit on \mathbb{R}^2 . On the limit, since $H \rightarrow 0$, the trace equation (4.34) becomes

$$\Delta u + c' = 0. \tag{4.35}$$

If $c' = 0$ then since $u > 0$, $u = \text{const}$ and hence $D^2u = 0$, giving a contradiction. If $c' \neq 0$, then again since $u > 0$, one must have $c' < 0$ and u is a quadratic polynomial on \mathbb{R}^2 . Since u is harmonic on the blow-up sequence and $c' < 0$ implies $N(u) \rightarrow -\infty$, this is also easily seen to be inconsistent with the requirement $u > 0$ everywhere. This proves (4.33) holds.

Returning to (4.31), we may work in the normalization above that $u(y_i) = 1$ and then (4.33) implies that $dv = u^{-1}du$ is bounded in L^∞ in bounded domains about y_i . Hence by (4.31), R_N is bounded in $L^{1,p}$ and so bounded in C^α . Since $R_N \rightarrow 0$ in L^2 locally on ∂M , one has

$$R_N \rightarrow 0 \text{ in } C^\alpha_{\text{loc}}(\partial M). \tag{4.36}$$

This is the main part of the estimate (4.27).

Next, one needs the same result for the $\text{Ric}(N, T)$ term. To do this, take the normal derivative of the scalar constraint (4.6), to obtain

$$\Delta N(u) + \Delta' u + N(H)N(u) + HNN(u) = \frac{1}{2}N(u)[|A|^2 - H^2 + s_\gamma] + \frac{1}{2}uN[|A|^2 - H^2 + s_\gamma]. \tag{4.37}$$

Since $\gamma' = 2A$, a standard formula for the variation of the Laplacian, (cf [Be]), gives $\frac{1}{2}\Delta' u = -\langle D^2u, A \rangle + \langle du, \beta(A) \rangle$ which is bounded in L^∞ . Also the terms $N(H)$, H and $NN(u) = -\frac{u}{2}[|A|^2 - H^2 + s_\gamma]$ all go to 0 in L^∞ . For the right side of (4.37), the coefficient of the first term goes to 0 in L^∞ while via (4.36) above, $uN[|A|^2 - H^2 + s_\gamma]$ is bounded in L^p provided $N(s_\gamma)$ is, and this in turn follows from an L^p bound on $\delta\delta A$.

To obtain this, return to (4.30) but with (N, X) in place of (N, N) , with X tangent to the boundary. Arguing as above, it follows that $\delta(R(\cdot, N)X, \cdot)$ is bounded in L^p . Via the Gauss–Codazzi equations $dA(X, Y, Z) = \langle R(N, X)Y, Z \rangle$, it follows that δdA is bounded in L^p . Similarly, in the normalization above, $\delta A = D^2u(N, \cdot) = \nabla_N du$ modulo lower order terms. Taking the exterior derivative d of this, one easily obtains that $d\delta A$ is bounded in L^p . This shows that ΔA is bounded in L^p , which of course gives the same for $\delta\delta A$ by elliptic regularity.

Finally, as in the proof of (4.33), $N(u) = c + \phi$ with ϕ bounded in $L^{1,p}$. Thus it follows from the divergence constraint as preceding (4.34) and elliptic regularity that $dN(u)$ is bounded in $L^{1,p}$ and so converges in C^α on ∂M . Since

$$-u \text{Ric}(N, \cdot) = dN(u) - A(du),$$

it now follows that $\text{Ric}(N, \cdot)$ converges to its limit, necessarily 0, in C^α on ∂M . Lastly, $\delta A = -dH - \text{Ric}(N, \cdot) \rightarrow 0$ in C^α and hence by elliptic regularity, $A \rightarrow 0$ in $C^{1,\alpha}$ so that again via the Gauss–Codazzi equations $dA(X, Y, Z) = \langle R(N, X)Y, Z \rangle \rightarrow 0$ in C^α . Combining the computations above now proves the estimate (4.27).

To complete the proof, we have $|R| \rightarrow 0$ on $\partial M \rightarrow \partial X$. On the 4-manifold $\mathcal{M} = M \times_u S^1$, the Einstein equations give the inequality

$$\Delta_{\mathcal{M}}|R_{g_{\mathcal{M}}}| + c|R_{g_{\mathcal{M}}}|^2 \geq 0, \tag{4.38}$$

where c is a fixed numerical constant, $\Delta_{\mathcal{M}}$ is the 4-Laplacian and $R_{g_{\mathcal{M}}}$ is the curvature tensor on \mathcal{M} . One has $|R_{g_{\mathcal{M}}}| = |R|$, (up to a constant). We have proved above that $R \rightarrow 0$ in L^2 locally on $(\mathcal{M}, (g_{\mathcal{M}})_i)$ and $R \rightarrow 0$ pointwise at $\partial\mathcal{M}$. It follows from the deGiorgi-Nash-Moser estimates for domains with boundary, (cf [GT, theorem 8.25]), that $R \rightarrow 0$ pointwise on \mathcal{M} and hence on M , contradicting (4.7). This completes the proof. \square

Next, we show that the potential function u is also controlled by the boundary data of Π_B .

Corollary 4.2. *For $(M, g, u) \in \mathcal{E}^o$, there is a constant U_0 depending only on the boundary data $(\gamma, H) \in \Pi^o(\mathcal{E}^o)$ such that*

$$u \leq U_0, \tag{4.39}$$

on M . Moreover, if $H \geq H_0 > 0$ on ∂M , then there exists U_0 as above depending in addition only on H_0 , such that on M ,

$$u \geq U_0^{-1}. \tag{4.40}$$

Proof. Let $S(s) = \{x \in (M, g) : \text{dist}(x, \partial M) = s\}$ be the geodesic ‘sphere’ about ∂M . Choose a fixed base point $x_0 \in S(1)$ and suppose one has the bound

$$c_0^{-1} \leq u(x_0) \leq c_0. \tag{4.41}$$

By theorem 4.1, the geometry of the annular region $A(\frac{1}{2}, 2)$ about $S(1)$ is uniformly controlled by the boundary data (γ, H) and so by integration of the static vacuum equations $u \text{ Ric} = D^2u$ along paths in $A(\frac{1}{2}, 2)$, one has

$$C_0^{-1} \leq u \leq C_0, \tag{4.42}$$

in $A(\frac{3}{4}, \frac{3}{2})$, where C_0 depends only on c_0 and (γ, H) . \square

To remove the dependence of C_0 in (4.42) on c_0 in (4.41), we need better control on the large-scale behavior of u . To do this, it is proved in [A2, lemma 3.6], that there is a constant K , depending only on C_0 , such that for all $s \geq 1$,

$$\sup_{S_c(s)} |du| \leq K(v_c(s))^{-1}, \tag{4.43}$$

where $v_c(s) = \text{area } S_c(s)$ and $S_c(s)$ is any component of the geodesic sphere $S(s)$. We point out that (4.43) holds for general static vacuum solutions, not only those in \mathcal{E}^o for instance. The estimate (4.43) is proved by studying the behavior of the harmonic potential $\log u$ on the Ricci-flat 4-manifold $(\mathcal{M}, g_{\mathcal{M}}, \log u)$ and then reducing to (M, g_M, u) .

Consider now the conformally equivalent metric

$$\tilde{g} = u^2 g. \tag{4.44}$$

It is well-known that the static vacuum Einstein equations (1.1) are equivalent to the equations $\tilde{\text{Ric}} = 2(dv)^2 \geq 0$, $\Delta_{\tilde{g}} v = 0$, $v = \log u$. The metric \tilde{g} thus has non-negative Ricci curvature with harmonic potential v . These are exactly the properties used to prove (4.43), and a brief examination of its proof shows that (4.43) also holds with respect to \tilde{g} , i.e.

$$\sup_{\tilde{S}_c(s)} |dv|_{\tilde{g}} \leq K(\tilde{v}_c(s))^{-1}, \tag{4.45}$$

again with $K = K(C_0)$. Since (M, g) is asymptotically flat and $u \rightarrow \text{const}$ at infinity, the area growth of geodesic spheres $\tilde{v}(s)$ in (M, \tilde{g}) satisfies $\tilde{v}(s)/s^2 \rightarrow \omega_2$, where $\omega_2 = \text{area} S^2(1)$. It follows then from the volume comparison theorem for Ricci curvature, (cf [Pe]), that

$$\tilde{v}(s) \geq \omega_2 s^2, \tag{4.46}$$

for all $s \geq 1$. As above, by integration of (4.45) along a geodesic ray starting from a suitable base point $x_1 \in S(1)$ out to infinity, one sees that (4.39) holds then globally on $M \setminus B(1)$, with U_0 again depending only on c_0 in (4.41). Using the static vacuum equations, the same integration along paths gives such a bound within $B(1)$. Thus, we see that (4.39) follows from (4.41).

To prove (4.41), suppose one has a static vacuum solution (M, g, u) with

$$u(x_0) = \varepsilon. \tag{4.47}$$

Renormalize u to $\bar{u} = u/u(x_0)$, so that $\bar{u}(x_0) = 1$ and $\bar{u} \rightarrow \varepsilon^{-1}$ at infinity. Then (4.41) holds, and hence so does (4.45) and (4.46). Again by integration along geodesics starting at x_0 and diverging to infinity, it follows that

$$u \leq U_1,$$

where U_1 depends only on the boundary data of Π_B . This proves the lower bound in (4.41).

To prove the upper bound, suppose instead

$$u(x_0) = \varepsilon^{-1}. \tag{4.48}$$

Then again we renormalize u to \bar{u} as above, so that now $u \rightarrow \varepsilon$ at infinity. This does not directly give a lower bound on ε via (4.45) and (4.46) as above. However, one may proceed as follows. First, it is well-known that static vacuum solutions come in ‘dual’ pairs, in that if (M, g, \bar{u}) is a static vacuum solution, then so is (M, \hat{g}, \hat{u}) with $\hat{g} = \bar{u}^4 g$, $\hat{u} = \bar{u}^{-1}$, cf [A2] for instance. Then (4.45) and (4.46) hold for (M, \hat{g}, \hat{u}) which as before by integration gives an upper bound $\hat{u} \leq U_1$ at infinity. Since near infinity, $\hat{u} \simeq \varepsilon^{-1}$, this again gives a bound on ε^{-1} . This completes the proof of (4.39).

To prove (4.40), note that (4.41) has been proved above, and hence by the maximum principle and normalization $u \rightarrow 1$ at infinity, (4.40) holds in the exterior region $M \setminus B(1)$. Thus, one only needs to consider the behavior near ∂M . For this, suppose (M, g, u) is a static vacuum solution, $C^{2,\alpha}$ up to ∂M with $u \geq 0$ on $\bar{M} = M \cup \partial M$. If $H \geq H_0 > 0$ on ∂M , we claim that necessarily

$$u > 0 \text{ on } \partial M.$$

For if $u = 0$ at some point $z \in \partial M$, then by (4.6), $\Delta_{\partial M} u + HN(u) = 0$ at z . Since $0 = u(z)$ is a global minimum for u , one has $\Delta_{\partial M} u \geq 0$ and by the Hopf maximum principle, $N(u) > 0$ at z . This gives a contradiction if $H > 0$. The same arguments prove the existence of a lower bound (4.40) by a contradiction argument, taking a sequence and passing to a limit, using theorem 4.1. \square

The previous results now lead quite easily to the following main result of this section.

Corollary 4.3. *The boundary map*

$$\Pi^o : \mathcal{E}^o \rightarrow \text{Met}^{m,\alpha}(\partial M) \times C_+^{m-1,\alpha}(\partial M),$$

is almost proper.

Proof. Let (M, g_i, u_i) be a sequence of static vacuum solutions in \mathcal{E}^o , with $\Pi^o(g_i, u_i) = (\gamma_i, H_i)$. Supposing $(\gamma_i, H_i) \rightarrow (\gamma, H)$ in $\text{Met}^{m,\alpha}(\partial M) \times C_+^{m-1,\alpha}(\partial M)$, we need to prove that the sequence (g_i, u_i) has a subsequence converging in $C^{m,\alpha}(M)$, modulo diffeomorphisms, to a limit $(M, g, u) \in \mathcal{E}$.

The curvature bound (4.1) and control of the intrinsic and extrinsic geometries of the boundary metrics first implies the metrics g_i cannot collapse within bounded distance to ∂M ,

i.e. there is a fixed constant $i_0 > 0$ such that the injectivity radius of (M, g_i) satisfies

$$inj_{g_i}(x) \geq i_0,$$

for $\text{dist}_{g_i}(x, \partial M) \leq K$. By the compactness theorem in, for instance [AT], it follows that a subsequence of (M, g_i) converges in $C^{m,\alpha}$, (and C^∞ in the interior), uniformly on bounded domains containing ∂M , to a limit (M', g) . One has $\partial M' = \partial M$ and g is a complete Riemannian metric on M' , $C^{m,\alpha}$ up to ∂M and C^∞ in the interior.

By corollary 4.2, the potential functions u_i also converge in $C^{m,\alpha}$, (in a subsequence) to a limit potential function u on M' , and the pair (g, u) gives a solution of the static vacuum Einstein equations. Since $u = \lim u_i$, it follows that

$$u \leq U_0, \tag{4.49}$$

on M' , for U_0 as in (4.39). Clearly the boundary metric and mean curvature of (M', g, u) are given by the limit values (γ, H) . To prove that $(M', g, u) \in \mathcal{E}$, one then needs to prove that (M', g, u) is asymptotically flat and M' is diffeomorphic to M . Note that since the convergence above is only uniform on compact sets, *a priori* there need not be any relation between the asymptotic structure of (M', g, u) and (M, g_i, u_i) for any given i .

The equation (4.43) holds on each (M, g_i, u_i) and by corollary 4.2, the constant K is uniform, independent of i . Moreover, as in the proof of corollary 4.2, there is a geodesic ray $\sigma = \sigma_i$ starting at any fixed base point in $S(1)$ and diverging to infinity, such that on the component $S_c(s)$ of $S(s)$ containing σ , one has

$$\sup_{S_c(s)} |du_i| \leq Ks^{-2},$$

where K is independent of i . Since u_i is harmonic, by elliptic regularity, (and scaling), a similar estimate holds for higher derivatives of u_i , and via the static vacuum equations, it follows that

$$\sup_{S_c(s)} |R_{g_i}| \leq Cs^{-3},$$

with again C independent of i . This means that the metrics (M, g_i, u_i) become asymptotically flat at infinity uniformly, at a rate independent of i . For R sufficiently large, $R \geq R_0$ independent of i , the geodesic spheres $S(R)$ and annuli $A(R, 2R)$ are close to Euclidean spheres and annuli, (when scaled by R^{-1}), and hence the geometry is close to that of Euclidean space; there can be no branching or joining of different components of $S(R)$ for $R \geq R_0$. This implies that the limit (M', g, u) has a single asymptotically flat end, and M' is diffeomorphic to M . \square

Remark 4.4. The results above also show that the boundary map Π_B is almost proper not only on \mathcal{E}^o but also its closure $\overline{\mathcal{E}^o}$. In other words, if (M, g_i, u_i) is a sequence of static vacuum solutions in \mathcal{E}^o with boundary data (γ_i, H_i) and $(\gamma_i, H_i) \rightarrow (\gamma, H)$ in $\text{Met}^{m,\alpha}(\partial M) \times C^{m-1,\alpha}(\partial M)$, then (M, g_i, u_i) converges in $C^{m,\alpha}$ (in a subsequence) to a limit (M, g, u) in $\overline{\mathcal{E}^o}$, with $H \geq 0$.

To see this, note that the constant Λ in theorem 4.1 does not depend on a positive lower bound on H , and so (4.1) and (4.2) hold for the sequence (M, g_i, u_i) above. Of course we are using here the fact that ∂M is outer-minimizing on the sequence (M, g_i, u_i) . Similarly in corollary 4.2, the upper bound U_0 on u_i does not depend on a lower bound for H . One may then use the argument concerning (4.47) to show that u_i cannot go to 0 on M away from ∂M . The proof of corollary 4.3 also does not require a bound on H away from 0.

It is also worth pointing out a brief examination of the proof shows that the results of this section only require that ∂M is outer-minimizing in a neighborhood of arbitrarily small but fixed size (depending on (M, g, u)) about ∂M .

5. Degree of Π^P

By [Sm], a smooth proper Fredholm map $F : B_1 \rightarrow B_2$ of Fredholm index 0 between connected Banach manifolds B_1, B_2 has a well-defined degree (mod 2). Namely, if y is a regular value of F , then $F^{-1}(y)$ is a finite set of points, and $\deg_{\mathbb{Z}_2} F$ is just the cardinality of $F^{-1}(y)$ (mod 2). In fact if B_1 and B_2 are oriented, then F has a well-defined degree in \mathbb{Z} , cf [ET, BFP].

From the discussion in section 1, the boundary map $\Pi_B : \mathcal{E}^+ \rightarrow \text{Met}(\partial M) \times C_+(\partial M)$, although Fredholm, is not proper. However, by theorem 1.2, the restricted boundary map $\Pi^o : \mathcal{E}^o \rightarrow \text{Met}(\partial M) \times C_+(\partial M)$ is almost proper. This in fact suffices to obtain a \mathbb{Z}_2 -valued degree on natural domains within \mathcal{E}^o .

Thus, as in the introduction, let $\partial\mathcal{E}^o$ be the boundary of \mathcal{E}^o within the space \mathcal{E}_S of static vacuum solutions. Hence, $(M, g, u) \in \partial\mathcal{E}^o$ if and only if (M, g, u) satisfies (1.10) but not (1.8). Let $Z = \Pi_B(\partial\mathcal{E}^o) \subset \text{Met}^{m,\alpha}(\partial M) \times C^{m-1,\alpha}(\partial M)$ be the image of $\partial\mathcal{E}^o$ under the boundary map Π_B and set

$$\mathcal{E}^P = (\Pi^o)^{-1}([\text{Met}^{m,\alpha}(\partial M) \times C^{m-1,\alpha}(\partial M)] \setminus Z).$$

Then, by theorem 1.2 and construction, the induced boundary map

$$\Pi^P : \mathcal{E}^P \rightarrow [\text{Met}^{m,\alpha}(\partial M) \times C^{m-1,\alpha}(\partial M)] \setminus Z \tag{5.1}$$

is proper. In particular \mathcal{E}^P has only finitely many components \mathcal{E}^{P_i} and on each component the \mathbb{Z}_2 -valued Smale degree is well-defined.

This discussion leads to theorem 1.3. Let \mathcal{E}^{P_0} be the component of \mathcal{E}^P containing the standard round flat solution, equal to the exterior of the round ball in \mathbb{R}^3 , with

$$\Pi^{P_0} : \mathcal{E}^{P_0} \rightarrow \mathcal{T}_0.$$

Theorem 5.1. *The degree of Π^{P_0} satisfies*

$$\deg_{\mathbb{Z}_2} \Pi^{P_0} = 1. \tag{5.2}$$

Proof. The proof is based on the black hole uniqueness theorem [I, R, BM], that the Schwarzschild metrics

$$g_{\text{Sch}}(m) = \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2 g_{S^2(1)}, \quad u = \sqrt{1 - \frac{2m}{r}}, \tag{5.3}$$

$r \geq 2m$, are the unique AF static vacuum metrics with a smooth horizon $\mathcal{H} = \{u = 0\}$. The induced metric on ∂M is $S^2(2m)$ —the round metric γ_{2m} of radius $2m$ on S^2 . The mean curvature satisfies $H = 0$. Of course the Schwarzschild metrics are not in \mathcal{E}_S , but instead lie at the boundary $\partial\mathcal{E}_S$.

Consider any sequence $\{(g_i, u_i)\} \in \mathcal{E}^{P_0}$ for which $\Pi^{P_0}(g_i, u_i) = (\gamma_i, H_i) \rightarrow (\gamma_{2m}, 0)$ smoothly. Clearly, $\{(g_i, u_i)\}$ is a divergent sequence in \mathcal{E}^o . By corollary 4.3 and remark 4.4, a subsequence of $\{(g_i, u_i)\}$ converges smoothly to a static vacuum limit (M, g, u) . (Of course one may have $u = 0$ on ∂M). On this limit,

$$H = 0,$$

at ∂M , so that ∂M is a minimal surface. From (4.6) one has $2\Delta_{\partial M} u = u(|A|^2 + s_\gamma) \geq 0$, and hence it follows from the maximum principle that $u = 0$ on ∂M . Via the static vacuum equations (1.1), this implies further that $A = 0$ and $N(u) = \text{const}$ at ∂M . The black hole uniqueness theorem then implies that any such limit is the Schwarzschild metric, and so unique up to scaling. Thus one has uniqueness for the boundary data $(\gamma, 0)$, so that almost all boundary metrics γ cannot be realized with $H = 0$ at ∂M , (the no-hair result).

Given this background, suppose

$$\deg_{\mathbb{Z}_2} \Pi^{P_0} = 0. \tag{5.4}$$

Then for any regular value (γ, H) of Π^{P_0} , the finite set $(\Pi^{P_0})^{-1}(\gamma, H)$, if non-empty, consists of at least two distinct static vacuum solutions $(g^1, u^1), (g^2, u^2)$. The regular values of Π^{P_0} are generic (of second category) in the range space, by the Sard–Smale theorem. Choose then a sequence of regular values $(\gamma_i, H_i) \rightarrow (\gamma_{+1}, 0)$ smoothly. (We set $m = 1/2$ here). Suppose for now that $(\Pi^{P_0})^{-1}(\gamma_i, H_i)$ is non-empty; this will be proved to be the case later.

Let $(g_i^1, u_i^1), (g_i^2, u_i^2)$ be any pair of corresponding distinct sequences in $(\Pi^{P_0})^{-1}(\gamma_i, H_i)$. By corollary 4.3 and remark 4.4, the sequences $(g_i^1, u_i^1), (g_i^2, u_i^2)$ have $C^{m,\alpha}$ convergent subsequences to limits $(g_\infty^1, u_\infty^1), (g_\infty^2, u_\infty^2)$ in $\bar{\mathcal{E}}^{P_0}$ and by the uniqueness above

$$g_\infty^1 = g_\infty^2 = g_{\text{Sch}}(m),$$

with $m = 1/2$, with $u_\infty^1 = u_\infty^2 = u$ in (5.3).

This implies that near g_{Sch} , the boundary map Π^{P_0} is not locally 1-1, and so presumably $D\Pi_B$ has a non-trivial kernel at g_{Sch} . (Note however that $g_{\text{Sch}} \notin \mathcal{E}_S$). We claim this is impossible. To prove the claim, let

$$g_{\text{Sch}} = u^2 d\theta^2 + g_{\text{Sch}},$$

be the four-dimensional Schwarzschild metric on $\mathbb{R}^2 \times S^2$, and similarly let

$$g_i^j = (u_i^j)^2 d\theta^2 + g_i^j,$$

be the four-dimensional static Ricci-flat metrics associated to (g_i^j, u_i^j) . By lemma 2.2, without loss of generality we may assume that each g_i^j is in Bianchi gauge with respect to g_{Sch} , so that, as in (2.10) and (2.11),

$$\beta_{g_{\text{Sch}}}(g_i^j) = 0,$$

for $j = 1, 2$ and i sufficiently large. By the smoothness of the convergence above, one may write

$$g_i^j = g_{\text{Sch}} + \varepsilon_i^j \kappa_i^j + O((\varepsilon_i^j)^2), \tag{5.5}$$

where $L(\kappa_i^j) = 0$ and L is the linearized Einstein operator (2.6) at g_{Sch} . The data g_i^j, g_{Sch} and κ_i^j are all smooth, (up to the boundary). The forms κ_i^j are only unique up to multiplicative constants, which will be determined by choosing ε_i^j so that the $C^{1,\alpha}$ norm of $g_i^j - g_{\text{Sch}}$ equals ε_i^j . Thus the $C^{1,\alpha}$ norm of κ_i^j is on the order of 1. Note that κ_i^j decays to 0 at infinity, so it is basically supported within compact regions of M . Let $\varepsilon_i = \max(\varepsilon_i^1, \varepsilon_i^2)$. Then

$$\varepsilon_i^{-1}(g_i^2 - g_i^1) = \kappa_i + O(\varepsilon_i),$$

where $\kappa_i = \varepsilon_i^{-1}(\varepsilon_i^2 \kappa_i^2 - \varepsilon_i^1 \kappa_i^1) \rightarrow \kappa$, where the convergence is in $C^{1,\alpha'}$, (in a subsequence). As previously, we need to show that the convergence is strong, so that $\kappa \neq 0$. This follows from a standard linearization and bootstrap argument, as preceding (4.16). In more detail, dropping the index i , we have $\Delta_{g^j} u^j = 0$, so that

$$\Delta_{g^2}(u^1 - u^2) = (\Delta_{g^2} - \Delta_{g^1})u^1.$$

In local harmonic coordinates, the right side this equation is on the order of ε in $C^{1,\alpha}$, and hence by elliptic regularity, $u_1 - u_2$ is on the order of ε in $C^{3,\alpha}$. Substituting this in the difference of the static equations $u^j \text{Ric}_{g^j} = D_{g^j}^2 u^j$ and arguing in the same way shows that the difference $g^1 - g^2$ is then also on the order of ε in $C^{3,\alpha}$. This proves the strong convergence.

It follows that the limit form

$$\kappa = (h, u'), \tag{5.6}$$

is a non-zero $C^{1,\alpha}$ weak solution of the linearized static vacuum equations $L(\kappa) = 0$ at g_{Sch} and since $\Pi^{P_0}(g_i^1, u_i^1) = \Pi^{P_0}(g_i^2, u_i^2)$, one has

$$\gamma'_h = H'_h = 0 \text{ at } \partial M, \tag{5.7}$$

where $\gamma'_h = h^T = h|_{\partial M}$. As discussed following (4.16), elliptic regularity implies that (h, u') is smooth and so in particular a strong solution. Below we will use the fact that the data (h, u') are in fact real-analytic up to ∂M , (again by elliptic regularity).

We claim that

$$(h, u') = 0 \text{ on } M, \tag{5.8}$$

which will give a contradiction. This is of course a linearized version of the black hole uniqueness theorem. It is possible that (5.8) can be proved by linearizing one of the existing proofs of black hole uniqueness in [I, R, BM]. However, we have not succeeded in doing this and instead (5.8) is proved in a manner similar to the proof of theorem 5.1.

First, the linearization of (4.6) gives, at ∂M ,

$$u' s_\gamma = 2\Delta u'.$$

Since $s_\gamma > 0$, the maximum principle implies that $u' = 0$ at ∂M . Next we claim $A' = 0$. To see this, the vacuum equations give $u \text{Ric} = D^2 u$ and $D^2 u = N(u)A + (D^2 u)^T$ when evaluated on tangent vectors to ∂M . Taking then the variation and evaluating tangentially gives $(u \text{Ric})' = 0$ so $0 = (D^2 u)' = (D^2)'u + D^2 u'$. The first term on the right vanishes when evaluating tangentially and hence so does the second term. This implies $0 = (N(u)A)' = N(u)A' + N(u')A = N(u)A'$. Since $N(u) = \text{const} \neq 0$, it follows that $A' = 0$. Similarly, taking the variation of the divergence or vector constraint gives $N(u') = \text{const}$.

Clearly $N(u') = m'$, (up to constants). A simple examination of the proof of black hole uniqueness in [R] applied to an Einstein deformation as in (5.3) and satisfying (5.7), shows easily that $N(u') = 0$ at ∂M . (One does not obtain any further information, since the bulk data in the Robinson proof, via divergence identities, are quadratic in the deviation from Schwarzschild).

Thus the variations $(\gamma', u', A', N(u'))$ of all the Cauchy data are trivial. As in the proof of theorem 4.1, we use a bootstrap argument to prove that the data (h, u') vanish to infinite order at ∂M , in geodesic gauge.

Thus, using geodesic normal coordinates near ∂M , write

$$g = dt^2 + g_t,$$

where $t(x) = \text{dist}(x, \partial M)$. We may assume, (by adding an infinitesimal deformation of the form $\delta^* V$ if necessary), that h preserves this gauge, so that $h_{0\alpha} = 0$, i.e. $h(N, \cdot) = 0$, $N = \partial_t$, near ∂M . By the discussion above, we have $u' = N(u') = 0$ at ∂M and similarly $h = \nabla_N h = 0$ at ∂M , so that

$$u' = O(t^2) \quad \text{and} \quad h = O(t^2). \tag{5.9}$$

The variation of the potential equation $\Delta_M u = 0$ gives

$$\Delta u' = -\Delta' u = \langle D^2 u, h \rangle - \langle \beta(h), du \rangle, \tag{5.10}$$

where β is the Bianchi operator, (cf [Be]). Since $\beta(h) = 0$ at ∂M , this gives $\Delta u' = 0$ at ∂M and hence $NN(u') = 0$ at ∂M , so that

$$u' = O(t^3). \tag{5.11}$$

Next the linearization of the Riccati equation gives $(\nabla_N A)' = -(A^2)' - (R_N)' = -(R_N)'$ at ∂M . One computes $*R_N = -(\text{Ric}^T)' = -u^{-1}(D^2 u)^T$, so $(*R_N)' = u^{-2}u'D^2 u - u^{-1}(D^2)'u - u^{-1}D^2 u'$, as a form on $T(\partial M)$. It follows from (5.11) that $(*R_N)' = 0$ at ∂N and hence

$(\nabla_N A)' = 0$ so that

$$h = O(t^3). \quad (5.12)$$

Next taking the normal derivative of (5.10) gives

$$\Delta N(u') = \langle \nabla_N D^2 u, h \rangle + \langle D^2 u, \nabla_N h \rangle + \langle \nabla_N \beta(h), du \rangle + \langle \beta(h), \nabla_N du \rangle,$$

which vanishes at ∂M and hence $u' = O(t^4)$. Substituting this in the linearized Riccati equation above and using previous estimates gives $h = O(t^4)$, and so on. It follows that (h, u') vanish to infinite order at ∂M . Since (h, u') are real-analytic up to ∂M , (in geodesic gauge), this implies that $h = u' = 0$ on M , i.e. (5.8) holds.

It remains only to prove that there are regular values (γ, H) of Π^{P_0} near $(\gamma_{+1}, 0)$ whose inverse image under Π^{P_0} is non-empty. However, (5.8) proves that $\text{Ker } D\Pi_B = 0$ at the Schwarzschild metric with boundary data $(\gamma_{+1}, 0)$. By continuity and the black hole uniqueness theorem, it follows that $\text{Ker } D\Pi_B = 0$, for all solutions with boundary data (γ, H) near $(\gamma_{+1}, 0)$ in $\text{Im } \Pi^{P_0}$. Thus, all such boundary data are regular values of Π^{P_0} . This completes the proof. \square

Remark 5.2. Although the proof of theorem 5.2 implies that $D\Pi_B$ has trivial kernel at the Schwarzschild metric g_{Sch} , one does not expect this to be the case for the cokernel. In fact, one expects that $\text{Coker } D\Pi_B$ is infinite dimensional, in that any boundary variation of the form $(k, 0)$, where k is a variation of the boundary metric, is not tangent to a curve of static metrics with $H = 0$ at ∂M . This amounts to the linearized version of the no-hair theorem, (which has not been proved as far as we are aware). In particular, we expect $D\Pi_B$ is not Fredholm at g_{Sch} .

Remark 5.3. The proof of theorem 5.2 above shows that $\text{Im } \Pi^{P_0}$ and hence $\text{Im } \Pi_B$ contains regular values. In particular, this means that the image $\text{Im } \Pi_B$ has non-empty interior and is in fact an open map near the Schwarzschild metric with boundary the horizon $r = 2m$. We point out that it has been an open question as to whether Π_B has *any* regular values. For instance, the analysis of Miao in [M1] shows that the exterior of the standard round unit ball $B^3(1)$ in flat \mathbb{R}^3 is a critical point of Π_B . In fact it is still unknown whether $\text{Im } \Pi_B$ has open interior near such standard boundary data $(\gamma_{+1}, 2)$.

Remark 5.3 indicates how little has been understood, and how much remains to be understood, regarding the global behavior of the boundary map Π_B . The proof of the almost properness of Π^o in theorem 1.2 uses the outer-minimizing property (1.8) in two key, but essentially independent ways. First, it is used to obtain the *a priori* curvature and related estimates discussed in section 4, i.e. to prevent blow-up behavior at the boundary. For this, only a small-scale version of (1.8) is necessary, in that one needs stability of the area of the boundary only to fourth order at ∂M , and this only in small discs in ∂M . Second, it is used to keep the boundary ∂M properly embedded in M , i.e. to prevent the passage from embedded to immersed behavior. Again, as mentioned in remark 4.4, only a local version of (1.8) is needed for this.

For further progress, an important issue is to find some condition on the boundary data (γ, H) which ensures that the two properties above hold. A natural question is whether positivity of the Gauss and mean curvatures is sufficient.

Acknowledgments

The first author is partially supported by NSF Grant DMS 0905159 and 1205947 while the second author is partially supported by NSF Grant DMS 1007156 and a Sloan Research

Fellowship. We thank Robert Bartnik, Piotr Chruściel, Gerhard Huisken and Xin Zhou for their interest and comments on this work. We are especially grateful to Simon Brendle for pointing out an error in a previous version of the paper.

References

- [ADN] Agmon S, Douglis A and Nirenberg L 1959 Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions: I *Commun. Pure Appl. Math.* **12** 623–727
 Agmon S, Douglis A and Nirenberg L 1964 Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions: II *Commun. Pure Appl. Math.* **17** 35–692
- [A1] Anderson M 2000 On stationary solutions to the vacuum Einstein equations *Ann. Henri Poincaré* **1** 977–94
- [A2] Anderson M 2000 On the structure of solutions to the static vacuum Einstein equations *Ann. Henri Poincaré* **1** 995–1042
- [A3] Anderson M 2008 On boundary value problems for Einstein metrics *Geom. Topology* **12** 2009–45
- [AH] Anderson M and Herzlich M 2008 Unique continuation results for Ricci curvature and applications *J. Geom. Phys.* **58** 179–207
 Anderson M and Herzlich M 2010 Unique continuation results for Ricci curvature and applications *J. Geom. Phys.* **60** 1062–7 (erratum)
- [AT] Anderson M, Katsuda A, Kurylev Y, Lassas M and Taylor M 2004 Boundary regularity for the Ricci equation, geometric convergence and Gel'fand's inverse boundary problem *Invent. Math.* **158** 261–321
- [B1] Bartnik R 1986 The mass of an asymptotically flat manifold *Commun. Pure Appl. Math.* **39** 661–93
- [B2] Bartnik R 1989 New definition of quasi-local mass *Phys. Rev. Lett.* **62** 2346–8
- [B3] Bartnik R 1997 Energy in general relativity *Tsing Hua Lectures on Geometry and Analysis* (Cambridge, MA: International Press) pp 5–27
- [B4] Bartnik R 2002 Mass and 3-metrics of non-negative scalar curvature *Proc. Int. Congr. of Mathematics* vol 2 (Beijing: Higher Education Press) pp 231–40
- [BS] Beig R and Simon W 1980 Proof of a multipole conjecture due to Geroch *Commun. Math. Phys.* **78** 75–82
- [BFP] Benevieri P, Furi M and Pera M 2002 On the product formula for the oriented degree for Fredholm maps of index zero between Banach manifolds *Nonlinear Anal. A* **48** 853–67
- [Be] Besse A 1987 *Einstein Manifolds* (Berlin: Springer)
- [Br] Bray H L and Chruściel P T 2004 The Penrose inequality *The Einstein Equations and the Large Scale Behavior of Gravitational Fields* ed P T Chruściel and H Friedrich (Basel: Birkhäuser) pp 39–70
- [BM] Bunting G and Massoud-ul-Alam A 1987 Non-existence of multiple black holes in asymptotically Euclidean static vacuum space-times *Gen. Rel. Grav.* **19** 147–54
- [ET] Elworthy K D and Tromba A J 1970 Degree theory on Banach manifolds *Nonlinear Functional Analysis I Proc. Symp. in Pure Mathematics* vol 18 (Providence, RI: American Mathematical Society) pp 86–94
- [GT] Gilbarg D and Trudinger N 1983 *Elliptic Partial Differential Equations of Second Order* 2nd edn (New York: Springer)
- [HI] Huisken G and Ilmanen T 2001 The inverse mean curvature flow and the Riemannian Penrose inequality *J. Differ. Geom.* **59** 353–437
- [I] Israel W 1967 Event horizons in static vacuum space-times *Phys. Rev.* **164** 1776–9
- [KO] Kennefick D and O'Murchadha N 1995 Weakly decaying asymptotically flat static and stationary solutions to the Einstein equations *Class. Quantum Grav.* **12** 149
- [LP] Lee J and Parker T 1987 The Yamabe problem *Bull. Am. Math. Soc.* **17** 37–91
- [M1] Miao P 2003 On existence of static metric extensions in general relativity *Commun. Math. Phys.* **241** 27–46
- [M2] Miao P 2010 Some recent developments of the Bartnik mass *Proc. Int. Conf. Chinese Math. 2007* vol 3 (Boston: International Press) pp 331–40
- [M] Müller zum Hagen H 1970 On the analyticity of static vacuum solutions of Einstein's equations *Math. Proc. Camb. Phil. Soc.* **67** 415–21
- [P] Peters P C 1979 Toroidal black holes? *J. Math. Phys.* **20** 1481–5
- [Pe] Petersen P 2006 *Riemannian Geometry* 2nd edn (New York: Springer)
- [RT] Regge T and Teitelboim C 1974 Role of surface integrals in the Hamiltonian formulation of general relativity *Ann. Phys.* **88** 286–318
- [R] Robinson D 1977 A simple proof of the generalization of Israel's theorem *Gen. Rel. Grav.* **8** 695–8
- [Sm] Smale S 1965 An infinite dimensional version of Sard's theorem *Am. J. Math.* **87** 861–6
- [Th] Thorne K 1975 A toroidal solution of the vacuum Einstein field equations *J. Math. Phys.* **16** 1860–5
- [T] Treves F 1975 *Basic Linear Partial Differential Equations* (New York: Academic)