

**MAT 534: HOMEWORK 10**  
DUE WED, DEC 5

1. Prove that  $\mathbb{Z}[\sqrt{-2}]$  is a Euclidean domain.
2. Determine the greatest common divisor in  $\mathbb{Q}[x]$  of  $a(x) = x^3 + 4x^2 + x - 6$  and  $b(x) = x^5 - 6x + 5$  and write it as a linear combination of  $a(x)$  and  $b(x)$ .
3. (a) Prove that every  $a \in \mathbb{Z}$  can be uniquely written in the form

$$a = \pm p_1^{n_1} \cdots p_k^{n_k} (q_1 \bar{q}_1)^{m_1} \cdots (q_l \bar{q}_l)^{m_l}$$

where  $p_i \in \mathbb{Z}$  are integers which are prime(=irreducible) as elements of  $\mathbb{Z}[i]$ , and  $q_i \in \mathbb{Z}[i]$  are irreducible elements of  $\mathbb{Z}[i]$  which are not in  $\mathbb{Z}$ .

- (b) Prove that a prime number  $p \in \mathbb{Z}_+$  remains irreducible in  $\mathbb{Z}[i]$  iff equation  $a^2 + b^2 = p$  has no integer solutions. (Hint:  $a^2 + b^2 = (a + bi)(a - bi)$ .) Deduce from this that prime numbers of the form  $4k + 3$  remain irreducible in  $\mathbb{Z}[i]$ . (In fact, it is known that a prime integer number is irreducible in  $\mathbb{Z}[i]$  iff it has the form  $4k + 3$ .)
- (c) Assuming the statement given in the previous part, prove that for a positive integer  $n$  the following statements are equivalent:
  - $n$  can be written as sum of two squares of integer numbers
  - $n$  can be written in the form  $n = z\bar{z}$ ,  $z \in \mathbb{Z}[i]$ .
  - In the prime factorization for  $n$  (in  $\mathbb{Z}$ ), each prime factor of the form  $4k + 3$  has even exponent.

4. Consider the ring  $R = \mathbb{Z}[\sqrt{-5}]$ .

- (a) Prove that elements  $2, 3, 1 \pm \sqrt{-5}$  are irreducible in  $R$ . [Hint: if  $2 = zw$ , then  $N(z)N(w) = N(2) = 4$ , where  $N(z) = z\bar{z} \in \mathbb{Z}_+$ .]
- (b) Show that  $R$  is not UFD because  $6 = 2 \cdot 3 = (1 + \sqrt{-5})(1 - \sqrt{-5})$ .
- (c) Define the ideals

$$I = (2, 1 + \sqrt{-5})$$

$$J = (3, 2 + \sqrt{-5})$$

$$J' = (3, 2 - \sqrt{-5})$$

Prove that these ideals are prime (see hint in Exercise 8, p. 293 in the book).

- (d) Prove that  $(2) = I^2$ ,  $(3) = JJ'$ ,  $(1 - \sqrt{-5}) = IJ$ ,  $(1 + \sqrt{-5}) = IJ'$ . Deduce from this that both factorizations  $6 = 2 \cdot 3 = (1 + \sqrt{-5})(1 - \sqrt{-5})$  give the same presentation for  $(6)$  as a product of prime ideals:  $(6) = I^2 JJ'$ .

5. In each of the following cases, determine whether  $a$  is an irreducible element of the ring  $R$ . Is  $a$  prime?

- (a)  $R = \mathbb{F}_2[x]$ ,  $a = x^2 + x + 1$
- (b)  $R = \mathbb{Z}[\sqrt{5}]$ ,  $a = 2$
- (c)  $R = \mathbb{Q}[x]$ ,  $a = x^4 + x^3 + x^2 + x + 1$
- (d)  $R = \mathbb{Q}[x]$ ,  $a = x^4 + 4$