

# MAT 126 Calculus B Spring 2006 Practice Midterm I — Solutions

Name: \_\_\_\_\_

I.D.: \_\_\_\_\_ Section number: \_\_\_\_\_

Answer each question in the space provided and on the reverse side of the sheets. Show your work whenever possible. Unless otherwise indicated, **answers without justification will get little or no partial credit!** Cross out anything that grader should ignore and circle or box the final answer.

1. (a) (10 points) Estimate the area under the graph of  $f(x) = 16 - x^2$  from  $x = 0$  to  $x = 4$  using four rectangles and right endpoints. Sketch the graph and rectangles. Is your estimate and underestimate or an overestimate?

*Solution.*

$$R_4 = f(1) + f(2) + f(3) + f(4) = 15 + 12 + 7 = 34$$

The function  $f(x)$  is concave downward, so that  $R_4 < A$ , i.e., it is an underestimate (sketch the graph!)

- (b) (10 points) Repeat part (a) using left endpoints.

*Solution.*

$$L_4 = f(0) + f(1) + f(2) + f(3) = 50$$

It is an overestimate,  $L_4 > A$  (sketch the graph!)

2. (a) (10 points) Evaluate integral by interpreting it as area

$$\int_{-5}^5 \sqrt{25 - x^2} dx$$

*Solution.* It is the upper half of the circle of radius 5 centered at the origin, so

$$\int_{-5}^5 \sqrt{25 - x^2} dx = \frac{1}{2} \pi 5^2 = 12.5\pi$$

- (b) (5 points) Determine a region whose area is equal to

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{\pi}{4n} \tan \frac{i\pi}{4n}$$

Do not evaluate the limit.

*Solution.* We get  $\Delta x = \pi/4n$ , so that  $b - a = \pi/4$ . Since  $x_i = i\Delta x$ , we get  $a = x_0 = 0$ , so that  $b = \pi/4$ . The sum is

the right endpoint sum  $R_n$  for  $f(x) = \tan x$  on the interval  $\left[0, \frac{\pi}{4}\right]$ , and we get

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{\pi}{4n} \tan \frac{i\pi}{4n} = \int_0^{\pi/4} \tan x dx,$$

which is the area of the region under the graph of the function  $f(x) = \tan x$  from  $x = 0$  to  $x = \pi/4$ .

**3.** Given two functions  $f(x)$  and  $g(x)$  which satisfy

$$\begin{aligned} \int_0^3 f(x) dx &= 5, & \int_0^5 f(x) dx &= 7, \\ \int_3^5 g(x) dx &= 1, & \int_0^5 g(x) dx &= 9, \end{aligned}$$

find

(a) (5 points)

$$\int_3^5 (3f(x) - g(x)) dx$$

*Solution.* We have

$$\int_3^5 f(x) dx = \int_0^5 f(x) dx - \int_0^3 f(x) dx = 7 - 5 = 2$$

so that

$$\int_3^5 (3f(x) - g(x)) dx = 3 \int_3^5 f(x) dx - \int_3^5 g(x) dx = 6 - 1 = 5$$

(b) (5 points)

$$\int_0^3 (f(x) + 2g(x)) dx$$

*Solution.* Similarly,

$$\int_0^3 g(x) dx = 9 - 1 = 8$$

and

$$\int_0^3 (f(x) + 2g(x)) dx = 5 + 2 \times 8 = 21$$

**4.** (5 points) Express the limit as a definite integral on the given interval  $[0, 4]$ :

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{e^{x_i}}{1 + x_i} \Delta x$$

*Solution.*

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{e^{x_i}}{1+x_i} \Delta x = \int_0^4 \frac{e^x}{1+x} dx$$

**5.** Evaluate the following indefinite integrals

(a) (5 points)

$$\int (3 \cos x - 4 \sin x) dx$$

*Solution.*

$$\int (3 \cos x - 4 \sin x) dx = 3 \sin x + 4 \cos x + C$$

(b) (10 points)

$$\int \frac{\cos x}{1 - \cos^2 x} dx$$

*Solution.* Using the fundamental trigonometric identity,

$$\int \frac{\cos x}{1 - \cos^2 x} dx = \int \frac{\cos x}{\sin^2 x} dx = -\frac{1}{\sin x} + C$$

— and integral from the table of indefinite integrals.

**6.** Evaluate the following definite integrals

(a) (5 points)

$$\int_1^2 x^{-2} dx$$

*Solution.*

$$\int_1^2 x^{-2} dx = \left. \frac{x^{-1}}{-1} \right|_1^2 = -\frac{1}{2} + 1 = \frac{1}{2}$$

(b)

$$\int_1^8 \frac{x-1}{\sqrt[3]{x^2}} dx$$

*Solution.*

$$\begin{aligned} \int_1^8 \frac{x-1}{\sqrt[3]{x^2}} dx &= \int_1^8 (x^{1/3} - x^{-2/3}) dx \\ &= \left( \frac{3}{4} x^{4/3} - 3x^{1/3} \right) \Big|_1^8 = \left( \frac{3}{4} \times 8^{4/3} - 3 \times 8^{1/3} \right) - \left( \frac{3}{4} - 3 \right) = 8.25 \end{aligned}$$

where we have used that  $8 = 2^3$ .

(c) (5 points)

$$\int_1^{27} \frac{1}{9t} dt$$

*Solution.*

$$\int_1^{27} \frac{1}{9t} dt = \frac{1}{9} \ln t \Big|_1^{27} = \frac{1}{9} \ln 27 = \frac{1}{3} \ln 3,$$

where we have used that  $\ln 1 = 0$  and  $27 = 3^3$ .

(d) (5 points)

$$\int_{\ln 3}^{\ln 6} 5e^x dx$$

*Solution.*

$$\int_{\ln 3}^{\ln 6} 5e^x dx = 5e^x \Big|_{\ln 3}^{\ln 6} = 5(e^{\ln 6} - e^{\ln 3}) = 5(6 - 3) = 15$$

(e) (10 points)

$$\int_{\pi/3}^{\pi/2} \csc x \cot x dx$$

*Solution.*

$$\begin{aligned} \int_{\pi/3}^{\pi/2} \csc x \cot x dx &= \int_{\pi/3}^{\pi/2} \frac{\cos x}{\sin^2 x} dx \\ &= \left( -\frac{1}{\sin x} \right) \Big|_{\pi/3}^{\pi/2} = -1 + \frac{2}{\sqrt{3}}. \end{aligned}$$