

Sample Final Exam

MAT 131, Fall 1998

The final will be held on **Thursday, December 17** at **7:00 pm**. Be sure to bring your Stony Brook ID card and your calculator. You may also bring a single 8½" by 11" sheet of **handwritten** notes. This sheet must not be a photocopy or computer printout. The locations of the final exam are given in the table below.

1	Barcus/Perez	Physics P113	2	Barcus/Perez	Physics P118
3/4	Sutherland/Kim	Humanities 101	5	Mandell/Herrera-Guzman	SB Union 236
6	Mandell/Herrera-Guzman	SB Union 237	7/8	Bernhard	Earth+Space 001
9	Bishop/Panafidin	Lt Engineer 152	10	Bishop/Panafidin	Hvy Engineer 201
11/12	Barcus/Weinberg	SB Union Aud.	13	Schalm/Cheng	SB Union Aud.
14	Schalm/McKenzie	SB Union Aud.	15	Schalm/Cheng	SB Union Aud.
16	Schalm/Moraru	SB Union Aud.	17/18	Bernhard/Teo	Earth+Space 001

This sample is a collection of problems similar to the type of questions which will be on the final. However, you are responsible for **all** the material we have covered this semester— just because a topic isn't on this sample doesn't mean it won't be on the final. Also, merely completing this sample exam is **not** adequate preparation for the final. You must also do a large number of additional problems, both those assigned in the homework and others like them. You should ensure that you know how to do all the problems on the midterms and the previous sample exams, as well. This sample contains more problems than the actual final.

1. Write the equation of the linear function f with $f(0) = 1$ and $f(3) = 3$. Also write the equation of the exponential function g with $g(0) = 1$ and $g(3) = 3$.

2. Compute the derivatives with respect to x for each of the following:

- | | | |
|------------------------------|-------------------------------------|---------------------------|
| (a) $x^{-1/2} + x + x^{1/2}$ | (c) $\arcsin(5x^2)$ | (f) $\ln(\cos x)$ |
| (b) $x^2 \sin(x^2)$ | (d) $\int_x^{\ln \pi} \cos(e^t) dt$ | (g) $e^{\tan x} \cos x$ |
| (c) $\sin^2 x + \cos^2 x$ | (e) $x^5 - \cos x \sin x$ | (h) $\frac{1+x^3}{1+x^2}$ |

3. Compute each of the following anti-derivatives (indefinite integrals):

- | | | |
|----------------------------|-------------------------------|------------------------|
| (a) $\int x^5 dx$ | (c) $\int x \cos x^2 dx$ | (e) $\int \tan x dx$ |
| (b) $\int \frac{3}{2x} dx$ | (d) $\int \frac{x}{1+x^4} dx$ | (f) $\int e^{2x+1} dx$ |

4. Evaluate each of the following definite integrals:

- | | | |
|--|---|-----------------------------------|
| (a) $\int_0^1 x^5 dx$ | (c) $\int_0^{\sqrt{\pi}} x \sin x^2 dx$ | (e) $\int_{-1}^1 \sqrt{1-x^2} dx$ |
| (b) $\int_0^{\frac{\pi}{4}} \sin x dx$ | (d) $\int_{-1}^1 \frac{x^2}{1+x^3} dx$ | (f) $\int_{-2}^2 x dx$ |

5. What is the average value of $y = x^3$ over the interval $[0, 2]$?

6. Find the point on the graph of the curve $y = \sqrt{x}$ that is closest to the point $(5, 0)$. (Hint: if d is the distance from $(5, 0)$ to a point on the curve, then it is permissible (and easier!) to minimize d^2 .)

7. Give the left-hand sum, right-hand sum, and trapezoid approximation for the integral $\int_0^2 e^{\sqrt{x}} dx$, using $n = 4$ rectangles. What should n be to ensure that the right-hand sum is accurate to within 0.001? (Hint: compare the expressions for the left-hand and right-hand sums for arbitrary n — what does this tell you about the exact value of the integral?)

8. Write the equation of the line tangent to the curve $y = 3x^2 + 2x + 1$ at the point $(1, 6)$.

9. Write the equation of the line tangent to the curve $y^3 - 2xy + x^3 = 0$ at the point $(1, 1)$.

10. From physics, we know that the illumination at a point x which is provided by a light source at L is proportional to the intensity of the light at L divided by the square of the distance between x and L . Suppose that two lights L_1 and L_2 are placed 20 meters apart, and that the intensity of L_2 is 8 times the intensity of L_1 . Where is the point on the line between L_1 and L_2 where the illumination is at a minimum?

11. Find the maximum and minimum values of the function $f(x) = x^3 + 3x^2 - 42x - 22$ on the interval $-5 \leq x \leq 2$.

12. Calculate the area of the region bounded by the graphs of $y = x/2$ and $x = y^2 - 3$.

13. Write the equation of the parabola which best approximates $y = \sin(x)$ at $x = \frac{\pi}{2}$ (that is, the second Taylor polynomial). Use your polynomial to find approximations of the nonzero solutions to $\sin(x) = x/3$. (Hint: graph the relevant functions for $-2\pi \leq x \leq 2\pi$ to make sure your answers make sense. The fact that $\sin(-x) = -\sin(x)$ is helpful.)

14. Use Newton's method to determine the nonzero solutions of $\sin(x) - x/3 = 0$ to within 0.000005. You may either use your answer to the previous problem as x_0 , or use $x_0 = 2$.

15. Compute the following limits. Distinguish between $+\infty$, $-\infty$, and "does not exist".

(a) $\lim_{x \rightarrow +\infty} \tan x$

(c) $\lim_{x \rightarrow 3} \frac{x-3}{x^2-9}$


(e) $\lim_{x \rightarrow +\infty} x^{27}e^{-x}$

(b) $\lim_{x \rightarrow 3} \frac{x^2-9}{x-3}$

(d) $\lim_{x \rightarrow 3} \frac{x^2+9}{x-3}$

(f) $\lim_{x \rightarrow 3^+} \frac{x^2+9}{x-3}$

16. The figure below is the graph of a function $f(x)$. Use it to sketch the graph of $f'(x)$ and the graph of $\int_0^x f(t) dt$.



17. A coffee filter has the shape of an inverted cone. Water drains from it at a constant rate of $10 \text{ cm}^3/\text{min}$. When the depth is 8 cm , the water level drops at a rate of $2 \text{ cm}/\text{min}$. What is the ratio of the height of the cone to its radius? You may find it useful to recall that the volume of a cone of radius r and height h is $\pi r^2 h/3$.

18. Let \mathcal{C} be the parametric curve given by

$$x = t - \sin 2\pi t \quad y = \sqrt{t} + \cos \pi t$$

What is the slope of \mathcal{C} at the point $(4, 2)$, when $t = 4$?