

MAT 320: Introduction to Analysis, Spring 2019

Homework Assignment 6

Please read Section 13 of Ross's textbook thoroughly before starting on the problem set below.

Optional supplementary reading: pp30-42 of Rudin's book

Problem Set 6 (**due at the start of recitation on Wednesday, 3/27**): D-G (below), 13.12, 13.14, 13.3ba, 13.3c (below), 13.15

Problem D

Let $d_{\mathbb{R}}$ be the standard metric on \mathbb{R} , $d_{\mathbb{R}}(x, x') = |x - x'|$. What are the closures in $(\mathbb{R}, d_{\mathbb{R}})$ of

$$\mathbb{N}, \quad S = \{2^m : m \in \mathbb{Z}\}, \quad \text{and} \quad [-2, 2] \cap \mathbb{Q} \quad ?$$

Note that $\infty \notin \mathbb{R}$. Give examples (3 in total) of a closed non-compact subset of $(\mathbb{R}, d_{\mathbb{R}})$, of a bounded non-compact subset of $(\mathbb{R}, d_{\mathbb{R}})$, and of a closed bounded non-compact subset of $(\mathbb{Q}, d_{\mathbb{R}}|_{\mathbb{Q}})$. *Answers only.*

Problem E

Let (X, d) be a metric space, \mathcal{C} be some collection of subsets of X (i.e. each *element* $B \in \mathcal{C}$ is a *subset* $B \subset X$), and $A = \bigcup_{B \in \mathcal{C}} B$.

- (a) Show that $\overline{A} \supset \bigcup_{B \in \mathcal{C}} \overline{B}$, where $\overline{A}, \overline{B} \subset X$ are the closures of A and B , respectively, in (X, d) .
- (b) Show that the opposite inclusion holds if \mathcal{C} is finite, but may not hold if \mathcal{C} is countable.

Problem F

Let d and d' be two metrics on the same set X that are **uniformly equivalent**: there exists $C \in \mathbb{R}^+$ such that

$$C^{-1}d(x, x') \leq d'(x, x') \leq Cd(x, x') \quad \forall x, x' \in X.$$

- (a) Show that a subset $U \subset X$ is open/closed w.r.t. d if and only if $U \subset X$ is open/closed w.r.t. d' .
- (b) Show that a sequence $(x_n)_n$ converges to x (resp. is Cauchy) w.r.t. d if and only if $(x_n)_n$ converges to x (resp. is Cauchy) w.r.t. d' .
- (c) Show that (X, d) is bounded/complete/compact if and only if (X, d') is.

Problem G

Let (X, d_X) and (Y, d_Y) be two metric spaces. Define

$$d_1, d_2, d_3 : (X \times Y)^2 \longrightarrow \mathbb{R}, \quad d_i((x, y), (x', y')) = \begin{cases} \max(d_X(x, x'), d_Y(y, y')), & \text{if } i = 1; \\ (d_X(x, x')^2 + d_Y(y, y')^2)^{1/2}, & \text{if } i = 2; \\ d_X(x, x') + d_Y(y, y'), & \text{if } i = 3. \end{cases}$$

- (a) Show that these three functions are metrics and that any two of them are uniformly equivalent.
- (b) Take $(X, d_X), (Y, d_Y) = (\mathbb{R}, d_{\mathbb{R}})$, i.e. \mathbb{R} with the standard metric. On a *whole* page by itself, draw 3 *huge* (but separate) copies of the first quadrant of the xy -plane. On the i -th copy, clearly draw the open unit ball $B_1^i((2, 2))$ around $(2, 2) \in \mathbb{R}^2$ with respect to the metric d_i (make sure it comes out *large*). On the same copy, *clearly* indicate what it means for this ball to be also open with respect to the metric d_{i+1} (with $d_4 \equiv d_1$), as F-(a) says should be the case. You can add a few words clarifying the diagrams, but they should be mostly clear by themselves.

Problem 13.3c

Show that the metric space (B, d) in 13.3a is complete.