

# MAT 531: Topology & Geometry, II Spring 2010

## Problem Set 7

Due on Thursday, 4/8, in class

*Note:* This problem set has two pages. It covers 1.5 weeks, and so it is longer than usual. The first problem is a leftover from Chapter 4.

1. Let  $X$  be a path-connected topological space and let  $(\mathcal{S}_*(X), \partial)$  be the singular chain complex of *continuous* simplices into  $X$  with *integer* coefficients. Denote by  $H_1(X; \mathbb{Z})$  the corresponding first homology group.

(a) Show that there exists a well-defined surjective homomorphism

$$h: \pi_1(X, x_0) \longrightarrow H_1(X; \mathbb{Z}).$$

(b) Show that the kernel of this homomorphism is the commutator subgroup of  $\pi_1(X, x_0)$  so that  $h$  induces an isomorphism

$$\Phi: \pi_1(X, x_0) / [\pi_1(X, x_0), \pi_1(X, x_0)] \longrightarrow H_1(X; \mathbb{Z}).$$

This is the first part of the Hurewicz Theorem.

*Hint:* For each  $x \in X$ , choose a path from  $x_0$  to  $x$ . Use these paths to turn each 1-simplex into a loop based at  $x_0$  and construct a homomorphism

$$\mathcal{S}_1(X) \longrightarrow \pi_1(X, x_0) / [\pi_1(X, x_0), \pi_1(X, x_0)].$$

Show that it vanishes on  $\partial \mathcal{S}_2(X)$ , well-defined on  $\ker \partial$  (may not be necessary), and its composition with  $\Phi$  is the identity on  $\pi_1(X, x_0) / [\pi_1(X, x_0), \pi_1(X, x_0)]$ . Sketch something.

2. (a) Prove Mayer-Vietoris for Cohomology: If  $M$  is a smooth manifold,  $U, V \subset M$  open subsets, and  $M = U \cup V$ , then there exists an exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_{\text{deR}}^0(M) & \xrightarrow{f_0} & H_{\text{deR}}^0(U) \oplus H_{\text{deR}}^0(V) & \xrightarrow{g_0} & H_{\text{deR}}^0(U \cap V) & \xrightarrow{\delta_0} \\ & & \xrightarrow{\delta_0} & H_{\text{deR}}^1(M) & \xrightarrow{f_1} & H_{\text{deR}}^1(U) \oplus H_{\text{deR}}^1(V) & \xrightarrow{g_1} & H_{\text{deR}}^1(U \cap V) & \xrightarrow{\delta_1} \\ & & \xrightarrow{\delta_1} & \dots & & & & & \\ & & \vdots & & & & & & \end{array}$$

where

$$f_i(\alpha) = (\alpha|_U, \alpha|_V) \quad \text{and} \quad g_i(\beta, \gamma) = \beta|_{U \cap V} - \gamma|_{U \cap V}.$$

(b) Suppose  $M$  is a compact connected orientable  $n$ -dimensional submanifold of  $\mathbb{R}^{n+1}$ . Show that  $\mathbb{R}^{n+1} - M$  has exactly two connected components. How is the compactness of  $M$  used?

3. (a) Show that the inclusion map  $S^n \rightarrow \mathbb{R}^{n+1} - 0$  induces an isomorphism in cohomology.  
 (b) Show that for all  $n \geq 0$  and  $p \in \mathbb{Z}$ ,

$$H_{\text{deR}}^p(S^n) \approx \begin{cases} \mathbb{R}^2, & \text{if } p=n=0; \\ \mathbb{R}, & \text{if } p=0, n, n \neq 0; \\ 0, & \text{otherwise.} \end{cases}$$

*Hint:* Discuss the  $p \leq 0$ ,  $p > n$ ,  $n=0, 1$  cases separately, before starting an induction on  $n$ . The case  $n=1$  is the subject of 4.14.

(c) Show that  $S^n$  is not a product of two positive-dimensional manifolds.

*Note:* Do *not* use the Kunneth formula, unless you are intending to prove it. However, the cup/wedge product can be used and might be useful here.

4. (a) Use Mayer-Vietoris (*not* Kunneth formula) to compute  $H_{\text{deR}}^*(T^2)$ , where  $T^2$  is the two-torus,  $S^1 \times S^1$ . Find a basis for  $H_{\text{deR}}^*(T^2)$ ; justify your answer.  
 (b) Let  $\Sigma_g$  be a compact connected orientable surface of genus  $g$  (donut with  $g$  holes). Let  $B \subset \Sigma_g$  be a small closed ball or a single point. Relate  $H_{\text{deR}}^*(\Sigma_g - B)$  to  $H_{\text{deR}}^*(\Sigma_g)$  (do not compute  $H_{\text{deR}}^p$  for  $p=1, 2$  explicitly).  
 (c) Show that

$$H_{\text{deR}}^p(\Sigma_g) = \begin{cases} \mathbb{R}, & \text{if } p=0, 2; \\ \mathbb{R}^{2g}, & \text{if } p=1; \\ 0, & \text{otherwise.} \end{cases}$$

*Hint:* Discuss the cases  $g=0, 1$  before starting an induction on  $g$ . Note that  $\Sigma_{g_1+g_2} \approx \Sigma_{g_1} \# \Sigma_{g_2}$ .

5. (a) Suppose  $q: \tilde{M} \rightarrow M$  is a regular covering projection with a finite group of deck transformations  $G$  (so that  $M = \tilde{M}/G$ ). Show that

$$q^*: H_{\text{deR}}^*(M) \rightarrow H_{\text{deR}}^*(\tilde{M})^G \equiv \{\alpha \in H_{\text{deR}}^*(\tilde{M}) : g^*\alpha = \alpha \ \forall g \in G\}$$

is an isomorphism. Does the statement continue to hold if  $G$  is not assumed to be finite?

(b) Determine  $H_{\text{deR}}^*(K)$ , where  $K$  is the Klein bottle. Find a basis for  $H_{\text{deR}}^*(K)$ ; justify your answer.

*Hint:* see Exercise 3 on p454 of Munkres.

6. Chapter 5, #4 (p216)

7. Let  $K = \mathbb{Z}$  and let  $\pi: \mathcal{S}_0 \rightarrow \mathbb{R}$  be the corresponding skyscraper sheaf, with the only non-trivial stack over  $0 \in \mathbb{R}$ ; see 5.11. What is  $\mathcal{S}_0$  as a topological space?

*Hint:* it is something familiar.

**Exercises** (*figure these out, but do not hand them in*): Chapter 5, #11, 13, 16, 17 (pp 216,217); verify Lemma 5.14 (p172). The kernel of the first map in (2) of Lemma 5.14 is denoted by  $A'' * B$  or  $\text{Tor}(A'', B)$  and known as the torsion product of  $A''$  and  $B$ ;  $A'' * B = B * A''$ .