

MAT 542: Algebraic Topology, Fall 2016

Suggested Problems for Week 13

You may hand in solutions to at *most* 2 problems every 2 weeks and no later than 2 weeks after the necessary material for them is covered in class.

From Munkres: 61.1, 61.3, 52.3, 52.4, 53.3

Problem U

Give \mathbb{Z} the discrete topology. The module $S_0(\mathbb{Z} \times \mathbb{Z})$ is free with basis

$$\sigma_{ij}: \Delta^0 \longrightarrow \mathbb{Z} \times \mathbb{Z}, \quad \sigma_{ij}(\Delta^0) = \{(i, j)\}, \quad i, j \in \mathbb{Z}.$$

Define

$$\delta \in S^0(\mathbb{Z} \times \mathbb{Z}) \quad \text{by} \quad \delta(\sigma_{ij}) = \begin{cases} 1, & \text{if } i=j; \\ 0, & \text{if } i \neq j. \end{cases}$$

- (a) Show that δ is a cocycle.
- (b) Show that δ is not in the image of $\times: H^*(\mathbb{Z}) \otimes H^*(\mathbb{Z}) \longrightarrow H^*(\mathbb{Z} \times \mathbb{Z})$.
- (c) Let $S \subset \mathbb{Z}$ be a finite subset. Write $\delta|_{\mathbb{Z} \times S}$ explicitly as an element in the image of

$$\times: H^*(\mathbb{Z}) \otimes H^*(S) \longrightarrow H^*(\mathbb{Z} \times S).$$

This problem illustrates the necessity of the assumption in the Kenneth Theorem for H^* that H_* of one of the factors be finitely generated in each dimension.