

# MAT 542: Algebraic Topology, Fall 2016

## Suggested Problems for Week 6

You may hand in solutions to at *most* 2 problems every 2 weeks and no later than 2 weeks after the necessary material for them is covered in class.

From Munkres: 14.2, 16.1, 16.3, 19.3, 19.4, 20.2, 20.6

### Problem J

Let  $K$  and  $L$  be simplicial complexes and  $g: |K| \rightarrow |L|$  be a continuous map. For each  $v \in \text{Ver}(K)$ , let

$$\tau_v = \{w \in \text{Ver}(L) : g(\text{St}(v, K)) \subset \text{St}(w, L)\}.$$

For each simplex  $\sigma = \{v_0, \dots, v_p\} \in K$ , let

$$\tau_\sigma = \tau_{v_0} \cup \dots \cup \tau_{v_p} \subset \text{Ver}(L).$$

- (1) Suppose  $x \in \text{Int } \sigma$  for some  $\sigma \in K$  and  $g(x) \in \text{Int } \tau$  for some  $\tau \in L$ . Show that  $\tau_\sigma \subset \tau$  and that this inclusion may be strict even if  $\sigma$  is a vertex and  $\tau_v \neq \emptyset$  for every  $v \in \text{Ver}(K)$ .
- (2) Let  $\sigma \in K$  and  $\sigma'$  be a face of  $\sigma$ . Show that  $\tau_\sigma \in L$  and  $\tau_{\sigma'}$  is a face of  $\tau_\sigma$ .
- (3) Let  $\hat{g}: \text{Ver}(K) \rightarrow \text{Ver}(L)$  be a simplicial approximation to  $g$ , i.e.

$$g(\text{St}(v, K)) \subset \text{St}(\hat{g}(v), L) \quad \forall v \in \text{Ver}(K).$$

Show that  $\hat{g}$  is carried by  $\sigma \rightarrow \tau_\sigma$ , i.e.  $\hat{g}(\sigma) \subset \tau_\sigma$  for every  $\sigma \in K$ .

- (4) By definition,  $g$  admits a simplicial approximation if and only if  $\tau_v \neq \emptyset$  for every  $v \in \text{Ver}(K)$ . Conclude that the homomorphism

$$g_* \equiv \hat{g}: H_*(K) \rightarrow H_*(L)$$

is independent of the choice of the simplicial approximation  $\hat{g}$  to  $g$ , if one exists.

This gives a more systematic perspective on the proof of Lemma 14.1 in Munkres.