

**Sample Midterm 1, MAT 142, Fall 2005**  
**First Midterm is Friday, Oct 7, at the usual time and place**

(1) Place the letter corresponding to the correct answer in the box next to each question.

(i)  Which of the following functions is not one to one on the interval  $(0, 1)$ ?  
 (a)  $1/x$  (b)  $\cos x$  (c)  $x^2$  (d)  $\sin 2x$  (e)  $\ln x$  (f) none of these.

(ii)  Evaluate  $\int_0^1 \frac{1}{1+x^2} dx$ .  
 (a)  $\pi/4$  (b)  $\pi/3$  (c)  $\pi/2$  (d)  $\frac{3}{4}\pi$  (e)  $\pi$  (f) none of these.

(iii)  If we rewrite  $4x - x^2$  by completing the square we get  
 (a)  $(x+2)^2 - 4$  (b)  $(x-2)^2 - 4$  (c)  $(x-2)^2 + 4$  (d)  $4 - (x-2)^2$  (e)  $4 - (x+2)^2$   
 (f) none of these.

(iv)  What is  $\frac{d}{dx}(\sec^{-1}(x))$ ?  
 (a)  $|x|/\sqrt{x^2 - 1}$  (b)  $1/(|x|\sqrt{x^2 + 1})$  (c)  $|x|/\sqrt{x^2 + 1}$  (d)  $1/\sqrt{x^2 - 1}$  (e)  $1/(|x|\sqrt{x^2 - 1})$ .  
 (f) none of these.

(v)  The error estimate for Simpson's rule with  $n$  subdivisions is of the form  
 a constant (depending on  $f$  and the interval) times:  
 (a)  $n^{-1}$  (b)  $n^{-2}$  (c)  $n^{-3}$  (d)  $n^{-4}$  (e)  $n^{-5}$  (f) none of these.

(vi)  The function  $\frac{1}{x} \ln x$  has an absolute maximum at  $x =$   
 (a) 0 (b) 1 (c)  $e/2$  (d)  $e$  (e)  $2e$  (f) none of these.

(vii)  What is the definition of  $\cosh x$ ?  
 (a)  $2/(e^x - e^{-x})$  (b)  $\frac{1}{2}(e^x - e^{-x})$  (c)  $\frac{1}{2}(e^x + e^{-x})$  (d)  $(e^x - e^{-x})/(e^x + e^{-x})$  (e)  $(e^x + e^{-x})/(e^x - e^{-x})$  (f) none of these.

(viii)  Find the inverse function of  $f(x) = (1+x)^3$   
 (a)  $(1+x)^{-3}$  (b)  $(1-x)^3$  (c)  $x^{1/3} - 1$  (d)  $(x-1)^{1/3}$  (e)  $(1+x)^{1/3}$  (f) none of these.

(ix)  The function  $f(x) = ax + \sin(x)$  is one to one  
 (a) for all values of  $a$  (b) for no values of  $a$  (c) only if  $a > 0$  (d) only if  $a \leq -1$   
 or  $a \geq 1$  (e) only if  $-1 \leq a \leq 1$  (f) none of these.

(x)  Find the derivative of  $f(x) = (\ln x)^3$   
 (a)  $3(\ln x)^2/x$  (b)  $3(\ln x)^2$  (c)  $2 \ln x$  (d)  $3/x^2$  (e)  $(\ln x)/x$  (f) none of these.

(2) Find each of the following integrals.

- (i)  $\int_2^3 \frac{1}{x} dx$
- (ii)  $\int_1^2 \frac{2 \ln x}{x} dx$
- (iii)  $\int_{-\pi/2}^{\pi/2} \cos^3(x) dx$
- (iv)  $\int \frac{dy}{y^2 - 2y + 5}$
- (v)  $\int \sin(x) e^{\cos(x)} dx$

(3) Do each of the following

- (i) Expand by partial fractions:  $\frac{x+4}{x^2+5x-6}$
  - (ii) Expand by partial fractions:  $\frac{1}{(x+1)(x^2+1)}$
  - (iii) Explain whether the integral converges or diverges:  $\int_0^1 \ln(x) dx$
  - (iv) Explain whether the integral converges or diverges:  $\int_0^\infty \frac{x^2}{1+x^2+x^3} dx$
  - (v) Explain whether the integral converges or diverges:  $\int_0^1 x^{-\alpha} dx, 0 < \alpha < 1$
- (4) How many divisions are required for the trapezoid rule to evaluate  $\int_0^1 e^x dx$  with error less than .0001?
- (5) Using the definitions of the hyperbolic functions, prove the identity

$$\cosh^2 x = \frac{1}{2}(\cosh 2x + 1).$$

- (6) Show that for  $|x| < 1$ ,  $\frac{d}{dx} \tanh^{-1}(x) = \frac{1}{1-x^2}$
- (7) What values of  $p \in (0, \infty)$  have the property that the volume of the solid obtained by rotating the region  $\{(x, y) : x > 1, y < x^{-p}\}$  around the  $x$ -axis is finite?