

MAT 331 Project
The box-counting dimension of the Weierstrass function

The Weierstrass function on $[0, 1]$ is defined by

$$f_{b,\alpha}(x) = \sum_{k=n}^{\infty} b^{-\alpha k} \sin(2\pi b^k x).$$

where b is a integer greater or equal to 2 and $0 < \alpha < 1$. It is a famous example of a continuous function that is nowhere differentiable. The graph of this function is also an example of a “fractal” set. This project concerns numerically estimating the box-counting dimension of the graph. Box-counting dimension is also commonly called Minkowski dimension.

The basic idea is to count how many boxes from a $\frac{1}{n} \times \frac{1}{n}$ grid a set hits. As n grows and the grid squares get smaller, more squares will hit the set. The number of squares often grows like a negative power of n , say $N_n \approx n^d$, and d is called the box-counting dimension. To compute α we take the logarithm of this equation and solve for α to get $\alpha = \log(N_n)/\log(n)$ (the base does not matter as long as you use the same base for both logarithms).

- (1) Take $b = 2$ and $\alpha = 1/4, 1/2, 3/4$. Plot the graph of $f_{2,\alpha}$ on $[0, 1]$.
- (2) Let $N_n(f)$ be the number of boxes from the standard $\frac{1}{n} \times \frac{1}{n}$ grid of squares in the plane that are hit by the graph of f . Show that

$$N_n(f) \approx \sum_{k=1}^n n \cdot (\max(f, I_k) - \min(f, I_k)),$$

where $I_k = [\frac{k-1}{n}, \frac{k}{n}]$. Write a **MATLAB** function that takes f and n as inputs and returns value of the sum on the right.

- (3) The box-counting dimension of the graph of f is defined as

$$\lim_{n \rightarrow \infty} \frac{\log N_n(f)}{\log n}.$$

Estimate this limit for the Weierstrass function $f = f_{2,\alpha}$ for several α values, say $\alpha = .2, .3, \dots, .9$. Plot your estimates. Can you formulate a conjecture for what the box-counting dimension is as a function of α ?

Since $f_{b,\alpha}$ is given by an infinite series, you will have to replace the infinite sum by a finite sum in your experiments. Taking $k = 50$ or 100 should give good results for $n \leq 1,000,000$.

- (4) Repeat your experiments for other values of b , say $b = 3, 4$. For each α , does the box-counting dimension seem the same as before, or does it change when b changes?

Remark: The dimension of the graph of the Weierstrass function is discussed in Chapter 5 of “Fractals in Probability and Analysis”; a PDF of this book is available at

http://www.math.stonybrook.edu/~bishop/fractalbook_final.pdf