



DIMENSIONS OF TRANSCENDENTAL JULIA SETS

Christopher Bishop, Stony Brook

On geometric complexity of Julia sets - III

27 Sept 2021 - 01 Oct 021

www.math.sunysb.edu/~bishop/lectures



THE PLAN

- Hausdorff and packing dimensions
- Polynomial versus transcendental
- Dimension $d = 2$
- Dimension $1 < d < 2$
- Dimension $d = 1$
- Open problems.

Defn: Minkowski dimension. If K is a bounded set, let $N(K, \epsilon)$ be the minimal number of cubes of diameter ϵ needed to cover K .


Upper Minkowski dimension:

$$\overline{\text{Mdim}}(K) = \limsup_{\epsilon \rightarrow 0} \frac{\log N(K, \epsilon)}{\log 1/\epsilon},$$

Lower Minkowski dimension


$$\underline{\text{Mdim}}(K) = \liminf_{\epsilon \rightarrow 0} \frac{\log N(K, \epsilon)}{\log 1/\epsilon}.$$

If these agree, common value is **Minkowski dimension**, $\overline{\text{Mdim}}(K)$.



Two disadvantages:

- Not defined for unbounded sets
- Countable sets can have dimension > 0 .



$\{1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots\}$ needs \sqrt{n} balls of size $1/n$ balls to cover.

The packing dimension fixes these problems:

Defn: packing dimension

$$\text{Pdim}(A) = \inf \left\{ \sup_{j \geq 1} \overline{\text{Mdim}}(A_j) : A \subset \bigcup_{j=1}^{\infty} A_j \right\},$$

where the infimum is over all countable covers of A .

By definition $\text{Pdim}(\cup_n A_n) = \sup_n \text{Pdim}(A_n)$.


Defn: α -dimensional Hausdorff content

$$\mathcal{H}_\infty^\alpha(K) = \inf \left\{ \sum_i |U_i|^\alpha \right\},$$

$\{U_i\}$ = cover of K , $|E|$ = diameter of a set E .

Like Minkowski dimension, but allows covering sets of different sizes.

Defn: Hausdorff dimension $\dim(K) = \inf \{ \alpha : \mathcal{H}_\infty^\alpha(K) = 0 \}$.



Always true that $\text{Hdim} \leq \text{Pdim}$, but " $<$ " can sometimes hold.

For polynomials, $\text{Hdim}(\mathcal{J})$ can take any value in $(0, 2]$.

Same for meromorphic functions (Bergweiler-Cui).

For transcendental entire functions, $\text{Hdim}(\mathcal{J}) \in [1, 2]$.

Today, I will only discuss entire functions:

- Sketch proof that $\text{Hdim} \geq 1$
- Discuss examples for $d = 2$, $1 < d < 2$, $d = 1$.

Thm (Baker): $H_{\dim} \geq 1$ for transcendental entire functions.

Lemma: Non-trivial loops escape

- Suppose curve γ in Fatou set surrounds a point of \mathcal{J} .
- If $\{f^n\}$ bounded on γ , also bounded on interior by max principle.
- Hence interior of γ in Fatou set, a contradiction.
- So a point in γ escapes. By normality all γ escapes.

Lemma: Iterates of γ have non-zero index around 0

- Suppose not.
- Then minimum principle applies and interior of γ escapes.
- But γ surrounds \mathcal{J} and hence surrounds a pre-periodic point.
- Contradiction.
- \Rightarrow iterates of γ surround every compact set.

Lemma: Multiply connected Fatou components are bounded:

- Suppose Ω is multiply connected and unbounded.
- Suppose $\gamma \subset \Omega$ surrounds a Julia point.
- γ escapes, index non-zero $\Rightarrow \gamma_n = f^n(\gamma)$ intersects Ω for all large n .
- $\Rightarrow \gamma_n \subset \Omega$ for all n .

...

- Schwarz lemma \Rightarrow hyperbolic distance from γ_{n+1} to γ_n is bounded.
- Implies $\text{diam}(\gamma_{n+1}) \leq C \cdot \text{diam}(\gamma_n)$.
- Implies that f grows polynomially. Contradiction.
- Hence multiply connected Fatou components are bounded.

Thm: \mathcal{J} contains non-trivial continuum, so $\text{Hdim}(\mathcal{J}) \geq 1$

- Suppose not. Then \mathcal{J} is totally disconnected.
- \Rightarrow one multiply connected Fatou component.
- Such a component is bounded. Contradiction.
- Hence \mathcal{J} contains a continuum.

So $\text{Hdim}(\mathcal{J}) \in [1, 2]$. Next we will discuss examples of

- $\text{Hdim}(\mathcal{J}) = 2$
- $1 < \text{Hdim}(\mathcal{J}) < 2$
- $\text{Hdim}(\mathcal{J}) = 1$

Many transcendental functions have $\dim(\mathcal{J}) = 2$:

Thm (Misiurewicz): $\dim(\mathcal{J}) = \mathbb{C}$ for $f(z) = e^z$.

Thm (McMullen): $\dim(\mathcal{J}) = 2$ and $\text{area}(\mathcal{J}) = 0$ for $f(z) = \lambda e^z$.

Thm (McMullen): $\text{area}(\mathcal{J}) > 0$ for $f(z) = \lambda \cdot \cosh(z)$.

Singular set = closure of critical values and finite asymptotic values
= smallest set so that f is a covering map onto $\mathbb{C} \setminus S$


Eremenko-Lyubich class = bounded singular set = \mathcal{B}

Speiser class = finite singular set = $\mathcal{S} \subset \mathcal{B}$


$\lambda \cdot \exp(z)$, and $\lambda \cdot \cosh(z)$ are in Speiser class.

Defn: Escaping set $I(f) = \{z : f^n(z) \rightarrow \infty\}$.

Fact: In general, $\mathcal{J}(f) = \partial I(f)$. For f in EL-class, $\mathcal{J}(f) = \overline{I(f)}$.




Definition of $\exp(z)$.



Definition of $\cosh(z)$.

$$\cosh(-x + iy) = \overline{\cosh(x + iy)}$$

Proof that $\text{area}(\mathcal{J}) > 0$ for cosh:




Let $S = 2\pi(n + im) + [0, 2\pi]^2$.

$\cosh(S)$ approximately covers annulus A_n of area $\simeq 2^{2|n|}$.

Annulus contains $\simeq e^{2|n|}$ disjoint translates of S .

Proof that $\text{area}(\mathcal{J}) > 0$ for cosh:



Omit $\simeq |n| \cdot e^{|n|}$ squares near y -axis, $\simeq e^{|n|}$ near ∂A_n .

Remaining squares cover $1 - O(|n| \cdot e^{-|n|})$ area of annulus.

$\sum_{n>0} ne^{-n} < \infty \Rightarrow$ positive area escapes (so is in \mathcal{J} .)

Transcendental examples with $1 < \text{Hdim}(\mathcal{J}) < 2$:

Gwyneth Stallard gave examples in EL-class: all $1 < d < 2$.


Rippon-Stallard proved $\text{Pdim}(\mathcal{J}) = 2$ in EL-class.

$\Rightarrow \text{Hdim} \neq \text{Pdim}$ can occur.

Simon Albrecht and I gave sequence in Speiser class with $\text{Hdim} \rightarrow 1$.

Do all values $(1, 2]$ occur in Speiser class?

Theorem (Stallard): There are Eremenko-Lyubich functions whose Julia sets have Hausdorff dimension close to 1.




There is EL function with tract $\{z : |f(z)| > 1\} \approx$ half-strip.

- Cauchy integrals
- Solve $\bar{\partial}$ -equation
- Use models theorem for EL-class.




Assume we have g in EL-class so that:

- $g(0) = 0$ and $|g(z)| < 1$ outside $S = \text{half-strip}$.
- $\{|z| < 1\}$ attracted to 0 (in Fatou set).
- Inside S , $g(z) \approx \exp(\exp(z - K))$



Inside S , $g(z) \approx \exp(\exp(z - K))$

This is conformal map of S to half-plane, followed by \exp .




Fixed point $g(0) = 0$ attracts everything in complement of S

Thus the Julia set is inside S . More precisely,

$$\mathcal{J}(g) \subset \bigcap X_n,$$

$$X_n = \{z : |g^k(z)| \geq K, k = 1, \dots, n\}.$$




To prove $\dim(\mathcal{J}) \leq 1 + \delta$, it suffices to show: if K large enough, then

(1) X_1 can be covered by disks $\{D_j\}$ so that $\sum_j \text{diam}(D_j)^{1+\delta} < \infty$,


(2) if D hits $\mathcal{J} \cap \{|z| > K\}$, then its preimages satisfy

$$\sum_{W_j \in f^{-1}(D)} \text{diam}(W_j)^{1+\delta} \leq \epsilon \cdot \text{diam}(D)^{1+\delta}.$$

If K is large, we may take $\epsilon > 0$ as small as we wish.



This is what preimages of one disk look like.




Preimage of gold disk $D = D(w, r)$ defined in two steps:

- stack of regions of diameter $O(r/|w|)$ on line $\{x = \log |w|\}$.
- region at height $2\pi k$ in stack has single preimage U_k of diameter


$$O\left(\frac{r}{|w|(\log |w| + 2\pi|k|)}\right).$$

These estimates only use $(\log z)' = 1/z$.



If $\delta > 0$ is fixed and K is large enough, then


$$\begin{aligned}
 \sum_k \text{diam}(U_k)^{1+\delta} &\lesssim \left(\frac{r}{|w|}\right)^{1+\delta} \sum_k \frac{1}{(\log |w| + 2\pi|k|)^{1+\delta}} \\
 &\lesssim \left|\frac{r}{w}\right|^{1+\delta} \frac{1}{\delta \log^{1+\delta} |w|} \ll \left|\frac{r}{w}\right|^{1+\delta} \ll r^{1+\delta}
 \end{aligned}$$




If $\delta > 0$ is fixed and K is large enough, then

$$\begin{aligned} \sum_k \text{diam}(U_k)^{1+\delta} &\lesssim \left(\frac{r}{|w|}\right)^{1+\delta} \sum_k \frac{1}{(\log |w| + 2\pi|k|)^{1+\delta}} \\ &\lesssim \left|\frac{r}{w}\right|^{1+\delta} \frac{1}{\delta \log^{1+\delta} |w|} \ll \left|\frac{r}{w}\right|^{1+\delta} \ll r^{1+\delta} \end{aligned}$$

This proves $\dim(\mathcal{J}(g)) \leq 1 + \delta$.






To get $1 < d < 2$ in Speiser class, we want to repeat same argument.

Find g is Speiser class so that

- $g(z) \approx \exp(\exp(z - K))$ in half-strip
- $|g| < 1$ outside n half-strip




To get $1 < d < 2$ in Speiser class, we want to repeat same argument.

Find g is Speiser class so that


- $g(z) \approx \exp(\exp(z - K))$ in half-strip
- $|g| < 1$ outside n half-strip

Unfortunately, no such g exists (B. 2017).




Instead, we use several strip-like tracts.

Then method of quasiconformal folding can be applied.



- $g(z) \approx \exp(\exp(\omega \cdot z - K))$ in each half-strip,
- $|g| < 1$ on sectors.
- Zero attracts central disk and all sectors.



A single R-component.

For QC-folding experts: vertical slits chosen to give $\tau \simeq 1$ on segments between strips and sectors.

Preimages of a disk follow the boundary.


Estimates like before, but more intricate.

Transcendental Julia sets with dimension 1

Theorem: $\text{Hdim}(\mathcal{J}) = \text{Pdim}(\mathcal{J}) = 1$ is possible. I gave example using infinite products. Somewhat technical.


New, more geometric, proof by Burkart and Lazebnik (“folding-like”).

Both proofs based on similar geometry.



Suppose we have annuli $\{r_k < |z| < r_{k+1}\}$.


maps $z \rightarrow C_k \cdot z^{2^k}$ from A_k to A_{k+1} .



Approximate this by placing 2^k zeros evenly around k th circle.

Approximate polynomial p near origin so there is some Julia set near origin.


Annuli escape, each corresponds to a different Fatou component.



The k th annulus looks rotationally invariant.


Other zeros are very, very close to 0 or ∞ .

Its inner and outer boundaries should be nearly circular.




Fatou component has a boundary around each zero.

Must surround pre-image of component at zero.




These boundaries and inner boundary map to outer boundary.



The k th annulus maps to $(k + 1)$ st annulus.

The $(k + 1)$ st annulus also has ring of boundary components.

These have a preimage in the k th annulus; a second ring.



There is an infinite sequence of rings converging to outer boundary.

Estimates show component boundary is countable union of C^1 curves.

“Buried” points have small dimension $\Rightarrow \dim(\mathcal{J}) = 1$.

Open questions:

My example has finite spherical 1-measure, but infinite packing 1-measure.

\Rightarrow not subset of rectifiable curve on sphere.


Can a transcendental entire Julia set lie on such a curve?

True for meromorphic. Julia set of $\tan(z)$ is a line.

Are the boundary components better than C^1 ?

Any Jordan curve is boundary of s.c. wandering domain – Boc Thaler.

Open questions: Black = known, Green = unknown



Which pairs $(Hdim, Pdim)$ can occur for a transcendental entire function?

Burkart: any pair (s, s) , $1 < s < 2$ can be approximated.

Can we have $Hdim = 1$, $Pdim = 2$?

Does $Hdim = Pdim$ hold for all polynomials?

Open questions:

Speiser class Julia sets take dimensions as close to 1 as desired.

Do they take all dimensions in $(1, 2]$?

Can the escaping set have dimension 1?

Open questions:

Eremenko and Lyubich showed Speiser class functions QC equivalent to f (i.e., $g = \psi \circ f \circ \varphi$) are a finite dimensional manifold M_f .

Think of $\dim(\mathcal{J}(g))$ as a function on M_f .

Often this is the constant 2 (e.g., finite order of growth).

Otherwise is it always non-constant?

Is the supremum over M_f always 2?

Do Shishikura's methods for Mandelbrot set apply?

