Stiefel–Whitney Currents

By Reese Harvey and John Zweck

ABSTRACT. A canonically defined mod 2 linear dependency current is associated to each collection v of sections, v_1, \ldots, v_m , of a real rank n vector bundle. This current is supported on the linear dependency set of v. It is defined whenever the collection v satisfies a weak measure theoretic condition called "atomicity." Essentially any reasonable collection of sections satisfies this condition, vastly extending the usual general position hypothesis. This current is a mod 2 d-closed locally integrally flat current of degree q = n - m + 1 and hence determines a \mathbb{Z}_2 -cohomology class. This class is shown to be well defined independent of the collection of sections. Moreover, it is the qth Stiefel-Whitney class of the vector bundle.

More is true if q is odd or q = n. In this case a linear dependency current which is twisted by the orientation of the bundle can be associated to the collection v. The mod 2 reduction of this current is the mod 2 linear dependency current. The cohomology class of the linear dependency current is 2-torsion and is the qth twisted integral Stiefel–Whitney class of the bundle.

In addition, higher dependency and general degeneracy currents of bundle maps are studied, together with applications to singularities of projections and maps.

These results rely on a theorem of Federer which states that the complex of integrally flat currents mod *p* computes cohomology mod *p*. An alternate approach to Federer's theorem is offered in an appendix. This approach is simpler and is via sheaf theory.

1. Introduction

It is well known [3, 17, 18, 22, 24] that the linear dependency set of a collection of sections of a vector bundle is related to the characteristic classes of the bundle. In particular, the zero set of a regular section defines a cohomology class which is the Chern–Euler class of the bundle. In [12] canonically defined current representatives of the Chern classes of a complex vector bundle were associated to collections of smooth sections of the bundle. These currents are called linear dependency currents since they are supported on the linear dependency set of the collection of sections.

The main aim of this paper is to study the linear dependency currents of a collection of sections of a *real* vector bundle. These are either mod 2 or bundle-twisted currents which represent either the mod 2 or twisted-integer Stiefel–Whitney classes of the bundle. Since they are either mod 2 or 2-torsion, these currents were overlooked in [12].

The linear dependency current associated with an ordered collection v of sections of a real vector bundle is defined in Section 3 paralleling a standard construction in enumerative geometry (see, for example, [19, 7]). In general, the most one can say is that this linear dependency current,

 $LD^{mod2}(\nu)$, is a mod 2 current and that it determines a \mathbb{Z}_2 -cohomology class which is well defined independent of the particular collection of sections of the bundle (Theorem 3.14).

However, if the degree of $\mathbf{LD}^{\text{mod2}}(v)$ is odd or equal to the rank of the bundle it is also possible to define a (bundle-twisted) linear dependency current, $\mathbf{LD}(v)$, which encodes certain (twisted) integer multiplicities of dependency among the sections (Proposition 3.16). The mod 2 reduction of this current $\mathbf{LD}(v)$ is the mod 2 current $\mathbf{LD}^{\text{mod2}}(v)$. The current $\mathbf{LD}(v)$ determines a (twisted) integer cohomology class well defined independent of the choice of collection v (Theorem 3.6). If the degree of $\mathbf{LD}(v)$ is less than the rank of the bundle (which occurs when the collection consists of more than one section), this cohomology class is a torsion class of order 2 (Corollary 3.8).

A major advantage of the approach taken here is that the linear dependency current is defined whenever the collection ν satisfies a weak measure theoretic condition called "atomicity", which is vastly more general than the usual general position hypothesis. For example, a real analytic collection of *m* sections of a rank *n* bundle is atomic provided that, for all $j \in \{0, 1, ..., m - 1\}$, the codimension of the set of points over which exactly *j* of the sections are linear independent is at least the expected codimension n - j (see [12], Proposition 2.14).

Another important property of the (mod 2) linear dependency current is that it is a (mod 2) locally integrally flat current. Recall that the integrally flat currents are those of the form R + dS, where R and S are rectifiable. Federer [5] proved that the complex of locally integrally flat currents (or such currents mod p) can be used to compute integer (or mod p) cohomology. In the Appendix we offer an alternate approach to the theory of (mod p) integrally flat currents and their cohomological properties. This simple approach is via sheaf theory and is distinct from the form of the theory given in the geometric measure theory literature.

The theory of dependency currents relies heavily on the theory of zero divisor currents, which was originally developed in [14] for "atomic" sections of an oriented vector bundle over an oriented manifold. The notion of an atomic section provides a generalization of the notion of a section being transverse to zero, one that is both useful and vastly more general. The zero divisor is a *d*-closed locally integrally flat current which determines a unique integer cohomology class, the Euler class. In this paper it is crucial that the notion of a zero current be understood in the non-orientable case. This is done in Section 2 where the zero divisor is defined as a bundle twisted current. This current determines a cohomology class (Theorem 2.5), which is the twisted Euler class, $\tilde{e} \in H^n(X, \mathbb{Z})$, of the vector bundle. The reduction mod 2 of the zero divisor eliminates the twisting, yielding a mod 2 current which represents the top Stiefel-Whitney class, $w_n \in H^n(X, \mathbb{Z}_2)$.

In Section 4 we identify the \mathbb{Z}_2 -cohomology class of the degree $q \mod 2$ current $\mathbf{LD}^{\mod 2}(v)$ as w_q , the qth Stiefel-Whitney class of the bundle (Theorem 4.1). Moreover, if q is odd, the \mathbb{Z} class of the twisted current $\mathbf{LD}(v)$ is identified as the qth twisted integral Stiefel-Whitney class $\widetilde{W}_q \in H^q(X, \mathbb{Z})$ (Theorem 4.10). This result is a corollary of the fact that the Bockstein of the mod 2 dependency current $\mathbf{LD}^{\text{mod } 2}(v)$ of degree q - 1 is the degree q twisted dependency current associated with a subcollection of the collection v (c.f. [24, 21]).

The Stiefel-Whitney classes were originally defined [22, 24, 21] as the primary obstruction to the existence of certain collections of linearly independent sections of a bundle $F \rightarrow X$. In Section 5 we examine the relationship between linear dependency currents and obstruction cocycles. Given a triangulation of X it is possible to choose a particular atomic collection of sections of F so that the Steenrod obstruction cocycle of the collection is defined. The Poincaré dual of such a cocycle is a cycle which defines a current on X by integration. We then show that this obstruction current is equal to the linear dependency current of the particular collection of sections. Among other things, this provides an alternate proof of the results of Section 4. In Section 6 higher dependency currents and general degeneracy currents of vector bundle maps are discussed, further expanding the results of [12]. Some of the degeneracy currents studied in Section 6 were not included in [12] since they are either not defined as twisted currents or their real cohomology class is zero. In these cases we can define mod 2 and/or twisted degeneracy currents. The integer cohomology classes of the twisted degereracy currents were first studied by Ronga [20] who proved that they are uniquely determined by their torsion-free part and mod 2 reduction. We expand upon Ronga's result by explicitly identifying the integer cohomology classes of the higher dependency currents as certain polynomials in the integer Pontrjagin and Stiefel–Whitney classes (Theorem 6.15).

In Section 7 applications of the theory to singularities of projections and maps are given. In particular we recover the well-known fact that the Steifel–Whitney classes of the tangent bundle TX and normal bundle NX of a submanifold $X \subset \mathbb{R}^N$ can be defined in terms of singularities of projections. The original version of this result is due to [17, 18], [23] (see also [1]). Note, however, that they only consider generic projections whose critical sets are non-degenerate, with multiplicity ± 1 . The atomic theory enables us to consider degenerate critical sets of arbitrary integer multiplicity (Proposition 3.16). In particular, if X is a real analytic submanifold, the tangent and normal Stiefel–Whitney classes can be defined in terms of the singularities of any projection whose degeneracy subvarieties have at least the expected codimension. Integer and mod 2 cohomological obstructions to the existence of smooth immersions and surjections between manifolds are also given, c.f. [20].

Two further applications are worth noting. Following [12] we can define mod 2 and twisted integer degeneracy currents associated with higher self-intersections of plane fields and invariants of pairs of foliations. Mod 2 and integer umbilic currents of hypersurfaces can also be studied using these ideas. Details of these two applications are left to the reader.

Secondary (Cheeger–Chern–Simons) Stiefel–Whitney classes will be introduced in a later paper. Canonical L_{loc}^1 representatives of these classes will be associated to each atomic collection of sections of a bundle with Riemannian connection. In the case of a single section α , the secondary Euler class is represented by the Chern–Euler potential $\sigma(\alpha)$. As is discussed in [10] this potential satisfies the important equation $d\sigma(\alpha) = \chi - \text{Div}(\alpha)$, where χ is the Euler form and $\text{Div}(\alpha)$ the divisor of the section. If the collection ν consists of more than one section, then there is a canonical L_{loc}^1 current $T(\nu)$ satisfying the current equation $dT(\nu) = -\mathbf{LD}(\nu)$, which represents the appropriate secondary Stiefel–Whitney class. This current equation is related to a formula of Eells [4].

Finally, we would like to draw the attention of the reader to recent work of Fu and McCrory [6] who, in the spirit of [15], have constructed current representatives for the tangential Stiefel–Whitney homology classes of a *singular* variety.

2. Divisors and atomicity

Harvey and Semmes defined the zero divisor current of an atomic section of an *oriented* real rank *n* vector bundle over an *oriented* manifold. The divisor is a codimension *n* current which is supported on the zero set of the section and which encodes the integer multiplicity of vanishing of the section. Furthermore, it is a *d*-closed locally integrally flat current whose cohomology class in $H^n(X, \mathbb{Z})$ is well defined independent of the choice of section. This class is the Euler class of the bundle.

The aim of this section is to define and study the zero divisor current in the case in which neither the vector bundle nor the base manifold are assumed to be orientable. In this case, the zero divisor is defined to be a current that is twisted by the orientation bundle of the vector bundle. It is also useful to define the mod 2 divisor to be the mod 2 reduction of the divisor. Both of these notions of divisor will be important in the study of linear dependency currents in Section 3.

We begin by recalling some definitions. Let $V \to X$ be a real rank *n* vector bundle over an *N*dimensional manifold. No orientation assumptions will be made on *V* or *X*. Let \mathcal{O}_X and \mathcal{O}_V denote the principal \mathbb{Z}_2 -bundles of orientations of *TX* and *V* over *X*. An \mathcal{O} -twisted *k*-form is a section of $\mathcal{O} \otimes_{\mathbb{Z}_2} \Lambda^k T^* X \to X$. (Often the subscript \mathbb{Z}_2 will be dropped when tensoring with \mathcal{O} .) A **density** is a top degree smooth \mathcal{O}_X -twisted form on *X*. Note that densities can be integrated over *X*. A **generalized function** is a continuous linear functional on the space of compactly supported smooth densities on *X*. A **current** of degree *p* is a differential *p*-form on *X* whose coefficients (with respect to each coordinate system) are generalized functions. Equivalently, a degree *p* current is a continuous linear functional on the space of compactly supported \mathcal{O}_X -twisted (N - p)-forms. Similarly, an \mathcal{O}_V -twisted forms. An L^1_{loc} form is a form whose coefficients are generalized functions, i.e., it acts on $\mathcal{O}_V \otimes \mathcal{O}_X$ -twisted forms. An L^1_{loc} form is a form whose coefficients are L^1_{loc} functions. Therefore, L^1_{loc} forms are currents that are not twisted. On the other hand, an oriented compact submanifold of *X* defines an \mathcal{O}_X -twisted current by integrating (untwisted) forms over it. Note that exterior differentiation is a well defined operation on (twisted) currents.

On a contractable open subset U of X, each \mathcal{O}_V -twisted current \widetilde{T} can be written in the form $\widetilde{T} = [e] \otimes T$ where $[e] \in \mathcal{O}_V$ is the orientation class of a local frame e for V over U and where T is a current on U. If V is orientable, each choice of orientation defines an isomorphism between \mathcal{O}_V -twisted currents and currents. These two isomorphisms differ by a minus sign. Note that the definition of a current on a non-orientable manifold agrees with that given in [13] but disagrees with that in [25, 26].

A. Divisors in the nonorientable case

In this subsection we define and study the divisor of a section of $V \to X$. The divisor is defined to be an \mathcal{O}_V -twisted current. Note that if V and X are oriented, the definition of divisor given below agrees with that of [14].

The solid angle kernel, θ , is the L_{loc}^1 form on \mathbb{R}^n obtained by pulling back the normalized volume form on the unit sphere to $\mathbb{R}^n \sim \{0\}$ by the radial projection map. The current equation $d\theta = [0]$ on \mathbb{R}^n , where [0] denotes the point mass at the origin, motivates the definition of divisor.

Definition 2.1. Let X be a smooth manifold and let $y = (y_1, \ldots, y_n)$ denote coordinates on \mathbb{R}^n . In the case n > 1 a smooth function $u : X \to \mathbb{R}^n$ is called **atomic** if, for each form $\frac{dy^l}{|y|^p}$ on \mathbb{R}^n with $p = |I| \le n - 1$, the pullback $u^*(\frac{dy^l}{|y|^p})$ to X has an $L^1_{loc}(X)$ extension across the zero set Z of u. Also assume that u does not vanish identically in any connected component of its domain X. In the case n = 1, it is convenient to define a smooth function $u : X \to \mathbb{R}$ to be atomic if $\log |u| \in L^1_{loc}(X)$, (c.f. [14]).

If u is atomic, then the zero set Z has measure zero in X (see [14]) so that the $L^1_{loc}(X)$ extensions are unique. In particular, the smooth form $u^*(\theta)$ on $X \sim Z$ has a unique $L^1_{loc}(X)$ extension across Z, and therefore defines a current on X.

Definition 2.2. Let $u : X \to \mathbb{R}^n$ be an atomic function. The **divisor** of u is the degree n current Div(u) on X defined by

$$\operatorname{Div}(u) := d(u^*\theta)$$
.

Atomicity is a weak condition which ensures the existence of a zero divisor. Harvey and Semmes proved that a large class of smooth functions are atomic. More specifically those functions which

vanish algebraically and whose zero sets are not too big in the sense of Minkowski content are atomic. In particular, real analytic functions whose zero sets have codimension n are atomic.

Lemma 2.3. Let g be a smooth $GL(n, \mathbb{R})$ -valued function on an oriented manifold X and let $u : X \to \mathbb{R}^n$ be atomic. Then v := ug is atomic and

$$\operatorname{Div}(v) = \pm 1 \operatorname{Div}(u)$$
,

where $\pm 1 := \operatorname{sgn} \det(g)$ is constant on connected components of X.

This result of [14] allows one to extend the notion of divisor to sections of vector bundles. First, a section v of a smooth vector bundle $V \rightarrow X$ is called **atomic** if for each choice of local frame *e* for *V* the function *v*, defined by v = ve, is atomic.

Definition 2.4. Let v be an atomic section of a rank n bundle $V \to X$. The **divisor**, Div(v), of v is the \mathcal{O}_V -twisted current on X defined locally on an open subset U of X as follows. Choose a local frame e for V over U and let $v : U \to \mathbb{R}^n$ be the coordinate expression for v determined by e. Then

$$\operatorname{Div}(v) := [e] \otimes \operatorname{Div}(v)$$
 on U.

In particular, if V is oriented, then Div(v) is a current on X.

As described in the Appendix, the **locally integrally flat** currents are those currents that can be expressed as R + dS where R and S are locally rectifiable. Furthermore, the complex $\widetilde{\mathcal{F}}_{loc}^*(X)$ of \mathcal{O}_V -twisted currents on X which are locally integrally flat may be used to compute the cohomology, $H^*(X, \mathbb{Z}_V)$, of X with integer coefficients twisted by \mathcal{O}_V , i.e., $\mathbb{Z}_V := \mathcal{O}_V \otimes_{\mathbb{Z}_2} \mathbb{Z}$.

Theorem 2.5. Let v be an atomic section of a real rank n vector bundle $V \to X$. The zero divisor, Div $(v) \in \tilde{\mathcal{F}}_{loc}^n(X)$, of v is an \mathcal{O}_V -twisted d-closed locally integrally flat current of degree n on X, whose support is contained in the zero set of the section v. Furthermore, if μ is another atomic section of V, then there is an \mathcal{O}_V -twisted locally rectifiable current R so that

$$Div(v) - Div(\mu) = dR$$
. (2.5.1)

That is, the cohomology class of Div(v) in $H^n(X, \mathbb{Z}_V)$ is well defined independent of the choice of section v. This class is the twisted Euler class \tilde{e} of V. In particular, if V is oriented, the Euler class $e \in H^n(X, \mathbb{Z})$ of V is the cohomology class of Div(v).

Corollary 2.6. Suppose that v is an atomic section of an odd rank bundle $V \to X$. Then there is an \mathcal{O}_V -twisted locally rectifiable current R on X so that

$$2 \operatorname{Div}(v) = dR$$
.

Consequently, the cohomology class $\tilde{e} \in H^n(X, \mathbb{Z}_V)$ of Div(v) is a torsion class of order 2, (c.f. [16]).

Proof of Corollary 2.6. Since the antipodal map on the even dimensional sphere S^{n-1} is orientation reversing, $\text{Div}(-\nu) = -\text{Div}(\nu)$. The result now follows by applying Equation (2.5.1).

The proof of Theorem 2.5 relies on the following general remark.

Remark 2.7. Let $p: \widetilde{X} \to X$ denote the double cover $\mathcal{O}_V \to X$. Note that the pullback bundle $\widetilde{V} = p^*V$ on \widetilde{X} is oriented by choosing the orientation on $\widetilde{V}_{\widetilde{X}}$ to be the one determined by the point $\widetilde{x} \in \widetilde{X}$. A current T on \widetilde{X} is **odd** if $a_*T = -T$, where $a: \widetilde{X} \to \widetilde{X}$ is the natural involution. Then odd currents on \widetilde{X} are in 1–1 correspondence with \mathcal{O}_V -twisted currents on X.

Proof of Theorem 2.5. The proof in the case that V is oriented, is given in [14], Theorem 5.1. In the case that V is nonorientable let $p: \widetilde{X} \to X$ be the double cover of Remark 2.7 and let $\widetilde{\nu}$ denote the pullback of the section ν to $\widetilde{V} \to \widetilde{X}$. Since \widetilde{V} is oriented the theorem holds for the section $\widetilde{\nu}$. Furthermore, by Lemma 2.3, $\text{Div}(\widetilde{\nu})$ is an odd current on \widetilde{X} which corresponds to the \mathcal{O}_V -twisted current $\text{Div}(\nu)$ on X. Incorporating the double cover $p: \widetilde{X} \to X$ into the proof of [14], Theorem 5.1, we observe that the locally rectifiable current \widetilde{R} on \widetilde{X} can be chosen to be odd. Finally, we let R be the corresponding \mathcal{O}_V -twisted current on X.

The following structure theorem for divisors, which is a corollary of [14], Proposition 4.3, is proved in [25]. Set $Z := \text{Zero}(\nu)$ and let

$$\operatorname{Reg} Z = \left\{ x \in X : Z \text{ is a codimension-} n \ C^1 \text{ submanifold near } x \right\}$$

denote the set of regular points of Z and let Sing $Z := Z \sim \text{Reg } Z$ denote the set of singular points. Let $\{Z_i\}$ denote the family of connected components of Reg Z.

Theorem 2.8. Let v be an atomic section of $V \to X$. If $Z_j \subset spt(Div(v))$, then

$$\mathcal{O}_{TZ_j} \cong \mathcal{O}_V \otimes \mathcal{O}_X \Big|_{Z_i}$$
,

and, given such an isomorphism, the submanifold Z_j defines an \mathcal{O}_V -twisted current $[Z_j]$ by integration. Furthermore, there are integers $n_j \in \mathbb{Z}$ such that

$$\operatorname{Div}(v) = \sum n_j \left[Z_j \right]$$
(2.8.1)

as \mathcal{O}_V -twisted currents on $X \sim \text{Sing } Z$.

Remark 2.9. By [14], Proposition 4.3, the integers n_j in Equation (2.8.1) can be calculated as follows. Let $x \in Z_j$ and let \mathcal{U} be an open neighborhood of x in X. Choose orientations for V and TX over \mathcal{U} . These orientations induce orientations on TZ_j and on the normal bundle NZ_j over \mathcal{U} . Let $\rho : S(V) \to S(\mathbb{R}^n)$ be an orientation preserving trivialization of the sphere bundle S(V) over \mathcal{U} . Then, for almost all $x \in Z_j$,

$$n_i = \text{Deg}\left(\rho \circ \nu : S\left(N_x Z_i\right) \to S\left(\mathbb{R}^n\right)\right)$$

is the degree of the induced map between oriented (n - 1)-spheres.

B. Mod 2 divisors

In this subsection, the mod 2 divisor of an atomic section is defined to be the mod 2 reduction of the divisor of the section. The mod 2 reduction of an \mathcal{O}_V -twisted locally integrally flat current is a mod 2 locally integrally flat current. At the cohomology level, mod 2 reduction is simply the natural mapping $H^*(X, \mathbb{Z}_V) \to H^*(X, \mathbb{Z}_2)$. The idea behind the definition of a mod 2 current is to completely ignore orientation issues by declaring a current T and its negative -T to be the same. Although they encode less information than their twisted counterparts, mod 2 currents have the advantage that they push forward under proper smooth maps. This fact will be particularly useful in Section 3.

Definition 2.10. Let $\mathcal{F}_{loc}^{p}(X)$ denote the space of locally integrally flat currents of degree p on X. Then the space $\mathcal{F}_{loc}^{mod \, 2}(X)$ of **mod 2 locally integrally flat currents** of degree p on X is defined to be the quotient $\mathcal{F}_{loc}^{p}(X)/2\mathcal{F}_{loc}^{p}(X)$. The natural mapping $\mathcal{F}_{loc}(X) \to \mathcal{F}_{loc}^{mod \, 2}(X)$ is called **mod 2 reduction**.

The spaces $\mathcal{F}_{loc}(X)$ and $\mathcal{F}_{loc}^{mod\,2}(X)$ are studied in the Appendix. In particular, we show there that the complex $\mathcal{F}_{loc}^{mod\,2}(X)$ may be used to compute the cohomology, $H^*(X, \mathbb{Z}_2)$, of X with \mathbb{Z}_2 coefficients. In Subsection A we saw that $\text{Div}(\nu) \in \widetilde{\mathcal{F}}_{loc}^n(X)$ is an \mathcal{O}_V -twisted locally integrally flat current. Now, by Lemma A.26, there is a canonical isomorphism

$$\widetilde{\mathcal{F}}_{\rm loc}(X) / 2\widetilde{\mathcal{F}}_{\rm loc}(X) \cong \mathcal{F}_{\rm loc}^{\rm mod\,2}(X) .$$
(2.11)

The induced mapping $\widetilde{\mathcal{F}}_{\text{loc}}(X) \to \mathcal{F}^{\text{mod } 2}_{\text{loc}}(X)$ is also called **mod 2 reduction**.

Definition 2.12. The mod 2 divisor, $\operatorname{Div}^{\operatorname{mod} 2}(\nu) \in \mathcal{F}_{\operatorname{loc}}^{\operatorname{mod} 2}(X)$, of an atomic section ν of $V \to X$ is defined to be the mod 2 reduction of the \mathcal{O}_V -twisted current $\operatorname{Div}(\nu) \in \widetilde{\mathcal{F}}_{\operatorname{loc}}(X)$.

Now the mod 2 version of Theorem 2.5 is immediate.

Theorem 2.13. Let v be an atomic section of a real rank n vector bundle $V \rightarrow X$. The mod 2 divisor, $\text{Div}^{\text{mod}2}(v)$, is a *d*-closed mod 2 locally integrally flat current of degree n on X, whose support is contained in the zero set of the section v. Furthermore, if μ is another atomic section of V, then there is a mod 2 locally rectifiable current R so that

$$\operatorname{Div}^{\operatorname{mod}2}(\nu) - \operatorname{Div}^{\operatorname{mod}2}(\mu) = dR$$

That is, the cohomology class of $\text{Div}^{\text{mod}_2}(v)$ in $H^n(X, \mathbb{Z}_2)$ is well defined independent of the choice of section v. This class is the mod 2 Euler (or top Stiefel–Whitney) class, w_n , of V.

Remark 2.14. By definition, the divisor $\text{Div}(\nu) \in \widetilde{\mathcal{F}}_{\text{loc}}(X)$ is determined by a collection of local divisors $\text{Div}(v_{\alpha})$ defined on open subsets U_{α} of X. These local divisors satisfy $\text{Div}(v_{\alpha}) = \pm \text{Div}(v_{\beta})$ on $U_{\alpha} \cap U_{\beta}$. The mod 2 divisor $\text{Div}^{\text{mod } 2}(\nu)$ is the mod 2 current which is naturally associated to this collection of local divisors.

Example 2.15. In general, a current representative for a mod 2 divisor is not *d*-closed. In fact, it is easy to construct sections ν for which there are *no d*-closed *current* representatives $T \in \mathcal{F}_{loc}(X)$ of $\text{Div}^{\text{mod } 2}(\nu)$ (though of course $\text{Div}(\nu)$ is a *d*-closed \mathcal{O}_V -twisted current representative of $\text{Div}^{\text{mod } 2}(\nu)$). This can be done as follows.

Let v be an atomic section of a nontrivial real line bundle $L \to X$. Choose a metric on L. Then the divisor of v is the d-closed \mathcal{O}_L -twisted current $\operatorname{Div}(v) = d(\frac{1}{2}\frac{v}{|v|}) \in \widetilde{\mathcal{F}}_{\operatorname{loc}}^1(X)$. The mod 2 reduction of $\operatorname{Div}(v)$ is the mod 2 divisor $\operatorname{Div}^{\operatorname{mod} 2}(v)$ which represents the first Stiefel–Whitney class $w_1(L) \in H^1(X, \mathbb{Z}_2)$. A current representative $T \in \mathcal{F}_{\operatorname{loc}}^1(X)$ of $\operatorname{Div}^{\operatorname{mod} 2}(v)$ can be constructed as follows. For simplicity we assume that 0 is a regular value of v. Let $Z = \operatorname{Zero}(v)$. Choose an auxiliary section μ of $L \to X$ so that $W = \operatorname{Zero}(\mu)$ is a submanifold which is transverse to Z. Let $p: \widetilde{X} \to X$ be the double cover orienting L and let $\widetilde{v}, \ \widetilde{\mu} : \widetilde{X} \to \mathbb{R}$ be the pullbacks of the sections v, μ . Set $\chi := \frac{\widetilde{\mu}}{|\mu|}$. Now $\chi \operatorname{Div}(\widetilde{v}) \in \mathcal{F}_{\operatorname{loc}}^1(\widetilde{X})$ is well defined and $T := \frac{1}{2}p_*(\chi \operatorname{Div}(\widetilde{v})) \in \mathcal{F}_{\operatorname{loc}}^1(X)$ is a current on X whose mod 2 reduction is $\operatorname{Div}^{\operatorname{mod} 2}(v)$. Finally $dT = 2[Z \cap W] \neq 0$ on X.

Reese Harvey and John Zweck

Next suppose that S = T - 2R is a current representative of $\text{Div}^{\text{mod }2}(\nu)$ for which dS = 0. This forces $[Z \cap W] = dR$ to be zero in $H^2(X, \mathbb{Z})$. But this is not always possible, since if $L \to \mathbb{RP}^3$ is the tautological line bundle, then $[Z \cap W]$ can be chosen to be the generator \mathbb{RP}^1 of $H^2(X, \mathbb{Z}) \cong H_1(X, \mathbb{Z}) = \mathbb{Z}_2$.

3. Linear dependency currents

In this section we associate to each atomic collection v of n - q + 1 sections of a real rank n bundle $F \to X$ a degree q current on X which is supported on the linear dependency set of the collection of sections. This current will be called the **linear dependency current** of the collection v. In all cases the linear dependency current exists as a mod 2 current. However, in the case where q is odd or where q = n, it can also be defined to be an \mathcal{O}_F -twisted current. Note that if q = n, the linear dependency current is simply the divisor of the section v (see Section 2). Henceforth we assume that q < n.

The linear dependency currents are defined using the construction of such currents in [12] which we now briefly recall. Let $F \to X$ be a real rank *n* vector bundle. Fix $q \in \{1, 2, ..., n-1\}$ and let $v = (v_1, ..., v_m)$ be a collection of m = n - q + 1 > 1 sections of $F \to X$. (Such collections will always be ordered.) These sections define a bundle map $v : \mathbb{R}^m \to F$ by $v(t_1, ..., t_m) := \sum_{i=1}^m t_i v_i$ which drops rank on the set where $v_1, ..., v_m$ are linearly dependent. Let $\pi : \mathbb{P}(\mathbb{R}^m) \to X$ denote the trivial bundle of (m-1)-dimensional real projective spaces and let $U \subset \mathbb{R}^m$ be the tautological line bundle over $\mathbb{P}(\mathbb{R}^m)$. Using π to pull back the bundle map $v : \mathbb{R}^m \to F$ to $\mathbb{P}(\mathbb{R}^m)$ and then restricting to the subbundle $U \subset \mathbb{R}^m$ we obtain an **induced section** \tilde{v} of the bundle $H := \text{Hom}(U, \pi^*F)$ over $\mathbb{P}(\mathbb{R}^m) \equiv \mathbb{P}(\mathbb{R}^m) \times X$. By construction the projection by π to X of the zero set of \tilde{v} is the linear dependency set of $v_1, ..., v_m$.

Definition 3.1. The collection v of sections v_1, \ldots, v_m is called **atomic** if the induced section \tilde{v} of $H \to \mathbb{P}(\underline{\mathbb{R}}^m)$ is atomic.

If the collection ν is atomic, then the \mathcal{O}_H -twisted divisor current, $\text{Div}(\tilde{\nu})$, and its mod 2 reduction, $\text{Div}^{\text{mod } 2}(\tilde{\nu})$, are well defined. The linear dependency current is defined to be the current push forward of $\text{Div}^{\text{mod } 2}(\tilde{\nu})$, or whenever possible the push forward of $\text{Div}(\tilde{\nu})$.

Generally speaking, it is not possible to push forward twisted currents on $\mathbb{P}(\mathbb{R}^m) \times X$ to X by the projection π . However, the push forward by π of an $\mathcal{O}_{\mathbb{P}(\mathbb{R}^m)} \otimes \pi^* \mathcal{O}_F$ -twisted current on $\mathbb{P}(\mathbb{R}^m) \times X$ is well defined and is an \mathcal{O}_F -twisted current on X. This observation together with the following elementary lemma, will be used to determine when it is possible to push forward the \mathcal{O}_H -twisted current Div($\tilde{\nu}$).

Lemma 3.2. If $n \equiv m \mod 2$, then there is a canonical isomorphism

$$\mathcal{O}_H \cong \mathcal{O}_{\mathbb{P}(\mathbb{R}^m)} \otimes \pi^* \mathcal{O}_F$$
.

Proof. First recall that if V and W are oriented finite dimensional vector spaces, then there is a canonical choice of orientation on $V \otimes W$. Furthermore, if the dimension of W is even this choice is independent of the orientation on V and so there is a canonical isomorphism $\mathcal{O}_W \cong \mathcal{O}_{V \otimes W}$. The canonical choice of orientation on $V \otimes W$ is defined as follows. Choose ordered bases $v = (v_1, \ldots, v_p)$ and $w = (w_1, \ldots, w_q)$ for V and W. Then the canonical orientation on $V \otimes W$ is given by the

ordered basis

$$v \otimes w = (v_1 \otimes w_1, v_2 \otimes w_1, \dots, v_p \otimes w_1, v_1 \otimes w_2, \dots, v_p \otimes w_2, \dots, v_p \otimes w_q) .$$
(3.3)

Second, recall [16] that there is a canonical vector bundle isomorphism $U^* \otimes U^{\perp} \cong T\mathbb{P}(\mathbb{R}^m)$, where U^{\perp} denotes the orthogonal complement of U in \mathbb{R}^m .

So if *n* and *m* are both even, then the result is true since there are canonical isomorphisms $\mathcal{O}_H \cong \mathcal{O}_{\pi^*F}$ (as *n* is even) and $\mathcal{O}_{\mathbb{P}(\mathbb{R}^m)} = \mathcal{O}_{U^* \otimes U^{\perp}} \cong \mathbb{R}$. Here the last isomorphism is well defined by sending $[u^* \otimes u^{\perp}]$ to 1, where u^{\perp} is chosen so that (u, u^{\perp}) is a positively oriented frame for $U \oplus U^{\perp} = \mathbb{R}^m$. Similarly, if *n* and *m* are both odd, then $\mathcal{O}_H \cong \mathcal{O}_{U^*} \otimes \mathcal{O}_{\pi^*F}$ and since m - 1 is even $\mathcal{O}_{U^*} \cong \mathcal{O}_{U^* \otimes U^{\perp}} = \mathcal{O}_{\mathbb{P}(\mathbb{R}^m)}$, as required.

A. Linear dependency currents (q odd)

Throughout this subsection we assume that q < n is odd, and hence, $m \equiv n \pmod{2}$. Then, by Lemma 3.2, the push forward by π of the $\mathcal{O}_{\mathbb{P}(\mathbb{R}^m)} \otimes \pi^* \mathcal{O}_F$ -twisted current $\text{Div}(\tilde{\nu})$ on $\mathbb{P}(\mathbb{R}^m) \times X$ exists and is an \mathcal{O}_F -twisted current on X.

Definition 3.4. Let q be odd. The linear dependency current, $LD(\nu)$, of an atomic collection ν of n - q + 1 sections of $F \to X$ is the \mathcal{O}_F -twisted current on X defined by

$$\mathbf{LD}(v) := \pi_* (\operatorname{Div} (\widetilde{v}))$$
.

Remark 3.5. The following equivalent definition of the linear dependency current is often useful, especially when *n* and *m* are both odd as in this case the fibers $\operatorname{Hom}(U, F_x)$ and $\mathbb{P}(\mathbb{R}^m)_x$ are nonorientable. Let $p : S(\mathbb{R}^m) \to \mathbb{P}(\mathbb{R}^m)$ be the double cover by the unit sphere and let $\rho :=$ $\pi \circ p : S(\mathbb{R}^m) \to X$. Let $NS(\mathbb{R}^m)$ denote the normal bundle to $S(\mathbb{R}^m)$ in \mathbb{R}^m , with its canonical orientation. Then, as above, there is associated to the collection ν a section $\hat{\nu}$ of the bundle $\hat{H} :=$ $\operatorname{Hom}(NS(\mathbb{R}^m), \rho^*F)$ over $S(\mathbb{R}^m)$. Note that the sections $\hat{\nu}$ and $\tilde{\nu}$ are simultaneously atomic. As above there is a canonical isomorphism $\mathcal{O}_{\hat{H}} \cong \rho^* \mathcal{O}_F$. Consequently $\operatorname{Div}(\hat{\nu})$ is a well-defined $\rho^* \mathcal{O}_F$ -twisted current on $S(\mathbb{R}^m)$. Then, if q is odd, we have that

$$\mathbf{LD}(\nu) = \frac{1}{2} \rho_* \left(\text{Div}\left(\widehat{\nu}\right) \right) \ . \tag{3.5.1}$$

We verify (3.5.1) as follows. First note that $\widehat{H} = p^*H$ and that $\widehat{\nu} = p^*\widetilde{\nu}$. Since $n \equiv m \pmod{2}$, the $\rho^*\mathcal{O}_F$ -twisted current $\text{Div}(p^*\widetilde{\nu})$ is even in that

$$a_* \operatorname{Div}(p^* \widetilde{\nu}) = \operatorname{Div}(p^* \widetilde{\nu}) \quad \text{on } S(\mathbb{R}^m),$$

where $a: S(\underline{\mathbb{R}}^m) \to S(\underline{\mathbb{R}}^m)$ is the antipodal map. Now even currents on $S(\underline{\mathbb{R}}^m)$ are in 1–1 correspondence with $\mathcal{O}_{\mathbb{P}(\underline{\mathbb{R}}^m)}$ -twisted currents on $\mathbb{P}(\underline{\mathbb{R}}^m)$. In particular, if q is odd,

$$\operatorname{Div}(\widetilde{\nu}) = \frac{1}{2} p_* \left(\operatorname{Div} \left(p^* \widetilde{\nu} \right) \right) \qquad \text{on } \mathbb{P} \left(\underline{\mathbb{R}}^m \right) \,.$$

This fact immediately implies (3.5.1). Note that if q is even, then $\text{Div}(p^*\tilde{\nu})$ is an odd twisted current on $S(\mathbb{R}^m)$ and so its current push forward is zero. In general, odd currents on $S(\mathbb{R}^m)$ are in 1–1 correspondence with $\mathcal{O}_U \otimes \mathcal{O}_{\mathbb{P}(\mathbb{R}^m)}$ -twisted currents on $\mathbb{P}(\mathbb{R}^m)$, see [26].

The following result generalizes Theorem 2.5.

Theorem 3.6. Let $F \to X$ be a real rank *n* bundle and let *q* be odd. For each atomic collection ν of n - q + 1 sections of $F \to X$ the linear dependency current $LD(\nu)$ is an \mathcal{O}_F -twisted *d*-closed locally integrally flat current of degree *q* on *X* whose support is contained in the linear dependency set of the collection of sections. Furthermore, if μ is another atomic collection of sections of $F \to X$, then there is an \mathcal{O}_F -twisted locally rectifiable current *R* so that

$$\mathbf{LD}(\nu) - \mathbf{LD}(\mu) = dR.$$

That is, the cohomology class of LD(v) in $H^q(X, \widetilde{\mathbb{Z}}_F)$ is well defined independent of the choice of sections.

Proof. Since the push forward of a locally rectifiable current is locally rectifiable, the current $LD(\nu)$ inherits its properties from those of the divisor of the induced section $\tilde{\nu}$ (see Theorem 2.5).

Note. In the next section the cohomology class of $LD(\nu)$ is shown to be $\widetilde{W}_q \in H^q(X, \mathbb{Z}_F)$, the (twisted) integer Stiefel–Whitney class of F, whose mod 2 reduction is the standard Stiefel–Whitney class $w_q \in H^q(X, \mathbb{Z}_2)$ of F.

Proposition 3.7. Let $v : \underline{\mathbb{R}}^m \to F$ be as above and let $\psi : \underline{\mathbb{R}}^m \to \underline{\mathbb{R}}^m$ and $\varphi : F \to F$ be bundle isomorphisms. Then the collection of sections corresponding to the bundle map $\varphi \circ v \circ \psi : \underline{\mathbb{R}}^m \to F$ is also atomic. Furthermore, if q is odd,

$$\mathbf{LD}(\varphi \circ \nu \circ \psi) = \operatorname{sgn} \operatorname{det}(\psi) \operatorname{sgn} \operatorname{det}(\varphi) \mathbf{LD}(\nu) ,$$

as \mathcal{O}_F -twisted currents on X.

The proof of the proposition will be given at the end of this subsection. The following result generalizes Corollary 2.6.

Corollary 3.8. Under the same hypothesis as in Theorem 3.6, with q < n, there is an \mathcal{O}_F -twisted locally rectifiable current R on X so that

$$2\mathbf{L}\mathbf{D}(\mathbf{v}) = dR.$$

Consequently, the cohomology class \widetilde{W}_q of LD(v) in $H^q(X, \mathbb{Z}_F)$ is a torsion class of order 2.

Proof of Corollary. Define $\psi : \mathbb{R}^m \to \mathbb{R}^m$ by $\psi(t_1, \ldots, t_m) = (-t_1, t_2, \ldots, t_m)$ and let $\mu := \nu \circ \psi$. Then, by Proposition 3.7, $\mathbf{LD}(\mu) = -\mathbf{LD}(\nu)$. The result now follows from Theorem 3.6. For an alternate proof, see Theorem 4.10.

Next we study the case q = 1 in more detail.

Remark 3.9. Let $\mathcal{O}_F^{\mathbb{R}} := \mathcal{O}_F \otimes_{\mathbb{Z}_2} \mathbb{R}$ denote the orientation line bundle of F. The divisor $\text{Div}(\sigma) = d(\frac{1}{2}\frac{\sigma}{|\sigma|})$ of an atomic section σ of $\mathcal{O}_F^{\mathbb{R}}$ is called the **orientation current** of F associated with σ . Note that $\text{Div}(\sigma)$ is an \mathcal{O}_F -twisted current whose cohomology class in $H^1(X, \widetilde{\mathbb{Z}}_F)$ is \widetilde{W}_1 .

To each collection v of n sections of a rank n bundle F there is an associated section σ of $\mathcal{O}_F^{\mathbb{R}}$ well defined as follows. Choose a local frame f for F and let A be the matrix defined by v = Af. Then $\sigma := [f] \otimes \det A$.

Proposition 3.10 (The case q = 1). Let v be an atomic collection of n sections of a rank n bundle F. Suppose that the associated section σ of $\mathcal{O}_F^{\mathbb{R}}$ is also atomic. Then

$$LD(v) = Div(\sigma)$$

as \mathcal{O}_F -twisted currents on X.

Proof. Choose a local frame f for F and define A by $\nu = Af$. Then the local expression for the induced section $\hat{\nu}$ of ρ : Hom $(NS(\mathbb{R}^n), \rho^*F) \to S(\mathbb{R}^n)$ is the mapping $\psi : X \times S(\mathbb{R}^n) \to \mathbb{R}^n$ defined by $\psi(x, y) = yA(x)$. Let θ denote the normalized solid angle kernel on \mathbb{R}^n . Now, by the Change of Variables and Stokes's Theorems,

$$\int_{\rho^{-1}(x)} \psi^* \theta = \frac{\det A(x)}{|\det A(x)|} \quad \text{for each } x \notin \operatorname{Zero}(\det A) \ .$$

This implies the proposition since $\text{Div}(\sigma) = [f] \otimes d\left(\frac{1}{2} \frac{\det A}{|\det A|}\right)$ and $\text{LD}(\nu) = [f] \otimes \frac{1}{2}\rho_*(d\psi^*\theta))$.

Proof of Proposition 3.7. By (3.5.1) it suffices to show that

$$\operatorname{Div}\left(\widehat{\varphi \circ \nu}\right) = \operatorname{sgn}\operatorname{det}(\varphi)\operatorname{Div}\left(\widehat{\nu}\right) \quad \text{on } S\left(\underline{\mathbb{R}}^{m}\right) \tag{3.11}$$

and, if q is odd, that

$$\Psi_* \operatorname{Div}\left(\widehat{\nu \circ \psi}\right) = \operatorname{sgn} \operatorname{det}(\psi) \operatorname{Div}\left(\widehat{\nu}\right), \quad \text{on } S\left(\underline{\mathbb{R}}^m\right), \quad (3.12)$$

where $\Psi : S(\mathbb{R}^m) \to S(\mathbb{R}^m)$ is the diffeomorphism induced by ψ . Now since $\widehat{\varphi \circ \nu} = \varphi \circ \widehat{\nu}$, (3.11) follows from Lemma 2.3. To prove (3.12) let $\mu = \nu \circ \psi$ and note that the pullback of the section $\widehat{\nu}$ of \widehat{H} by Ψ is a section $\Psi^* \widehat{\nu}$ of $\Psi^* \widehat{H} = \text{Hom}(\Psi^* NS(\mathbb{R}^m), \rho^* F)$. Let $\psi^* : \text{Hom}(\Psi^* NS(\mathbb{R}^m), \rho^* F) \to \text{Hom}(NS(\mathbb{R}^m), \rho^* F)$ be the bundle isomorphism defined by $\psi^*(\alpha) := \alpha \circ \psi$. Then

$$\widehat{\mu} \;=\; \psi^* \left(\Psi^* \widehat{
u}
ight) \,.$$

Clearly, $\hat{\nu}$, $\Psi^*\hat{\nu}$ and $\hat{\mu}$ are simultaneously atomic. Since Equation (3.12) is local on X we can assume that F and X are oriented. Then $\hat{\mu}$ and $\hat{\nu}$ are sections of the oriented bundle \hat{H} over the oriented manifold $S(\mathbb{R}^m)$. Furthermore, since q is odd the orientation induced on $\Psi^*\hat{H}$ by the diffeomorphism Ψ is the same as that induced by the bundle isomorphism ψ^* . Equation (3.12) now follows immediately by applying Lemma 2.3 to ψ , the Change of Variables theorem to Ψ , and noting that sgn det $(D\Psi) = \text{sgn det}(\psi)$.

B. Mod 2 linear dependency currents

Since mod 2 currents can always be pushed forward by proper maps we can define the mod 2 linear dependency current for an even as well as an odd number of sections.

Definition 3.13. Suppose $1 \le q \le n$. The mod 2 linear dependency current, $LD^{mod 2}(\nu)$, of an atomic collection ν of m = n - q + 1 sections of $F \to X$ is defined to be the current push forward of the mod 2 divisor of the induced section $\tilde{\nu}$ of $H \to \mathbb{P}(\mathbb{R}^m)$,

$$\mathbf{LD}^{\mathrm{mod}\,2}(\nu) := \pi_*\left(\mathrm{Div}^{\mathrm{mod}\,2}(\widetilde{\nu})\right)$$
.

Note. If q is odd or q = n the mod 2 linear dependency current, $LD^{mod 2}(v)$, is the mod 2 reduction of the \mathcal{O}_F -twisted linear dependency current, LD(v).

The mod 2 analogs of Theorem 3.6 and Proposition 3.7 hold. In particular,

Theorem 3.14. For each atomic collection ν of n - q + 1 sections of a real rank n bundle $F \rightarrow X$ the linear dependency current $LD^{mod2}(\nu)$ is a d-closed mod 2 locally integrally flat current of degree q on X whose support is contained in the linear dependency set of the collection of sections. Furthermore, if μ is another atomic collection of sections of $F \rightarrow X$, then there is a mod 2 locally rectifiable current R so that

$$\mathbf{LD}(\nu)^{\mathrm{mod}2} - \mathbf{LD}^{\mathrm{mod}2}(\mu) = dR.$$

That is, the cohomology class of $LD^{mod2}(v)$ in $H^q(X, \mathbb{Z}_2)$ is well defined independent of the choice of sections.

Note. In the next section the cohomology class of $LD^{mod 2}(v)$ is shown to be $w_q(F) \in H^q(X, \mathbb{Z}_2)$, the *q*th Steifel–Whitney class of *F*.

C. The structure of linear dependency currents

The following result concerning the structure of the twisted linear dependency current builds on Proposition 2.8 of [12] and Theorem 2.8 above. Let q be odd and let v be an atomic collection of m = n - q + 1 sections of $F \to X$. Suppose that the zero set, $Z(\tilde{v})$, of \tilde{v} is a smooth submanifold of $\mathbb{P}(\mathbb{R}^m) \times X$. Let $\{Z_j\}$ denote the connected components of $Z(\tilde{v})$. Then, by Theorem 2.8, there are integers $n_j \in \mathbb{Z}$ so that

$$\operatorname{Div}\left(\widetilde{\nu}\right) = \sum n_{j} \left[Z_{j} \right]$$
(3.15)

as $\mathcal{O}_{\mathbb{P}(\mathbb{R}^m)} \otimes \pi^* \mathcal{O}_F$ -twisted currents on $\mathbb{P}(\mathbb{R}^m)$. By [12], Proposition 2.8, the subset

$$RK_{m-1}(v) := \{x \in X : \operatorname{rank} v_x = m-1\}$$

of the linear dependency set of ν is a locally rectifiable set. Let $RK_j := RK_{m-1}(\nu) \cap \pi(Z_j)$. If $n_j \neq 0$, then

$$\mathcal{O}_{TRK_j} \cong \mathcal{O}_F \otimes \mathcal{O}_X \big|_{RK_j}$$

(wherever it makes sense) and, given an isomorphism of these two bundles, RK_j defines an \mathcal{O}_F -twisted current $[RK_j]$ by integration. Arguing as in the proof of [12], Proposition 2.8, it follows that $\pi_*[Z_j] = [RK_j]$. Consequently we have the following:

Proposition 3.16. Let v be as above. Then

$$\mathbf{LD}(v) = \sum n_j \left[RK_j \right]$$

as \mathcal{O}_F -twisted currents on X, where the integers n_i are given by (3.15).

Next we examine the structure of the twisted and mod 2 linear dependency currents in the case that m - 1 of the *m* sections are everywhere linearly independent.

Theorem 3.17. Let μ_1, \ldots, μ_m be a collection of m = n - q + 1 sections of $F \to X$. Suppose that μ_1, \ldots, μ_{m-1} are everywhere linearly independent. Choose a metric on F and let μ_m^{\perp} denote

Stiefel-Whitney Currents

the projection of μ_m onto the orthogonal complement of μ_1, \ldots, μ_{m-1} in *F*. Then μ_m^{\perp} is atomic if and only if the induced section $\tilde{\mu}$ of $H \to \mathbb{P}(\mathbb{R}^m)$ is atomic. Furthermore, if *q* is odd, then

$$\mathbf{LD}(\mu) = \operatorname{Div}\left(\mu_m^{\perp}\right) \quad \text{as } \mathcal{O}_F \text{-twisted currents on } X , \qquad (3.17.1)$$

and, for any q,

$$\mathbf{LD}^{\mathrm{mod}\,2}(\mu) = \operatorname{Div}^{\mathrm{mod}\,2}\left(\mu_m^{\perp}\right) \quad \text{as mod } 2 \text{ currents on } X . \tag{3.17.2}$$

Proof. We present the proof in the twisted case. The mod 2 case follows similarly. The first step is to choose local coordinates and frames and to relate the local coordinate expression for the induced section $\tilde{\mu}$ of $H \to X \times \mathbb{P}(\mathbb{R}^m)$ to that of the section μ_m^{\perp} .

First note that the linear dependency set of μ_1, \ldots, μ_m is equal to the zero set $Z(\mu_m^{\perp})$ of μ_m^{\perp} . Fix a point $x_0 \in Z(\mu_m^{\perp})$ and let \mathcal{U} be a sufficiently small open neighborhood of x_0 in X. Choosing orientations for TX and F over \mathcal{U} we can regard $\mathbf{LD}(\mu)$ and $\mathrm{Div}(\mu_m^{\perp})$ as currents on \mathcal{U} . Now, since μ_1, \ldots, μ_{m-1} are linearly independent, there is precisely one point \tilde{x} of the zero set $Z(\tilde{\mu})$ of $\tilde{\mu}$ in $X \times \mathbb{P}(\mathbb{R}^m)$ lying over each point x of $Z(\mu_m^{\perp})$ in X. Let $\mathcal{W} \subset \pi^{-1}(\mathcal{U})$ be a sufficiently small open neighborhood of \tilde{x}_0 in $X \times \mathbb{P}(\mathbb{R}^m)$ which contains $Z(\tilde{\mu}) \cap \pi^{-1}(\mathcal{U})$.

Choose the coordinate chart $\psi : \mathbb{R}^{m-1} \to \mathbb{P}(\mathbb{R}^m)$ defined by $\psi(s) = [s, 1]$ and the local frame u for $U \to \mathbb{P}(\mathbb{R}^m)$ over $\psi(\mathbb{R}^{m-1})$ defined by $u(s) = (s, 1) \in U_{[s,1]} \subset \mathbb{R}^m$. Note that the orientations induced on $T\mathbb{P}(\mathbb{R}^m)$ and U by ψ and u are compatible and that $\mathcal{W} \subset \psi(\mathbb{R}^{m-1}) \times \mathcal{U}$ since μ_1, \ldots, μ_{m-1} are linearly independent on \mathcal{U} . Also note that the orientations on U and F induce a natural orientation on H over \mathcal{W} .

Choose a positively oriented local frame f_1, \ldots, f_n for F over \mathcal{U} so that $f_i = \mu_i$ for $1 \le i \le m-1$ and $f_i \perp \text{Span}\{f_1, \ldots, f_{m-1}\}$ for i > m-1. Define $a_i : \mathcal{U} \to \mathbb{R}$ by

$$\mu_m = \sum_{i=1}^n a_i f_i \; .$$

Let $a' = (a_1, \ldots, a_{m-1})$ and $a'' = (a_m, \ldots, a_n)$.

Then the local coordinate expression for μ_m^{\perp} in terms of the local frame f_m, \ldots, f_n is $a'': \mathcal{U} \to \mathbb{R}^q$ and the coordinate expression for $\tilde{\mu}$ in terms of the frames u, f is $(a'', s+a'): \mathcal{U} \times \mathbb{R}^{m-1} \to \mathbb{R}^n$. These two coordinate expressions can be related as follows. Let $\Psi: \mathcal{U} \times \mathbb{R}^{m-1} \to \mathcal{U} \times \mathbb{R}^{m-1}$ be the orientation preserving change of variables $\Psi(x, s) = (x, s + a'(x))$ and let $\mathrm{Id}: \mathbb{R}^{m-1} \to \mathbb{R}^{m-1}$ denote the identity map. Then

$$(a'', s + a') = (a'' \times \operatorname{Id}) \circ \Psi$$
.

By the Change of Variables Theorem and Lemma 3.18 below it follows that $\pi_*(\text{Div}(\widetilde{\mu})) = \text{Div}(\mu_m^{\perp})$ as required.

The following elementary fact about divisors is included for the sake of completeness.

Lemma 3.18. Let X be an oriented manifold and $f : X \to \mathbb{R}^n$ a smooth map. Let Id : $\mathbb{R}^m \to \mathbb{R}^m$ be the identity map and let $\pi : X \times \mathbb{R}^m \to X$ denote projection onto X. Then f is atomic if and only if $f \times \text{Id}$ is atomic. Furthermore, if $X \times \mathbb{R}^m$ is given the induced orientation, then

$$\pi_*(\operatorname{Div}(f \times \operatorname{Id})) = \operatorname{Div}(f)$$

Proof. By induction we may assume that m = 1. Let t denote the coordinate on \mathbb{R} . First, $\frac{dtdf^{I}}{|(t,f)|^{p+1}} \in L^{1}_{loc}(X \times \mathbb{R})$ for $p = |I| \le n-1$ iff $\log |f| \in L^{1}_{loc}(X)$ in case p = 0, and iff $\frac{df^{I}}{|f|^{p}} \in L^{1}_{loc}(X)$ in case p > 0. This is because

$$\int_{|t|$$

Of course, $\frac{df^I}{|(t,f)|^p}$ is dominated by $\frac{df^I}{|f|^p}$. Consequently $f \times \text{Id}$ is atomic if and only if f is atomic.

Let θ_n denote the normalized solid angle kernel on \mathbb{R}^n , and recall that $\text{Div}(f) = d(f^*\theta_n)$ where d is exterior differentiation of generalized forms. Let ∂S denote the boundary of a current S. Then $dS = (-1)^{k+1} \partial S$, where $k = \deg S$. Consequently $\pi_* d = -d\pi_*$. Therefore, it suffices to show that

$$\pi_*\left((f\times \mathrm{Id})^*\theta_{n+1}\right) = -f^*\theta_n .$$

Let $\omega_n = \operatorname{Vol}(S^{n-1})$ and $\lambda(f) = df_1 \wedge \ldots \wedge df_n$. Then

$$(f \times \mathrm{Id})^* \theta_{n+1} = \frac{1}{\omega_{n+1}} \frac{t \lambda(f)}{\left(t^2 + |f|^2\right)^{\frac{n+1}{2}}} - \frac{\omega_n}{\omega_{n+1}} \frac{|f|^n dt}{\left(t^2 + |f|^2\right)^{\frac{n+1}{2}}} f^* \theta_n$$

Now, since the push forward of $(f \times \text{Id})^* \theta_{n+1}$ by π is equal to the integral of $(f \times \text{Id})^* \theta_{n+1}$ over the fibers of π ,

$$\pi_* \left((f \times \mathrm{Id})^* \theta_{n+1} \right) = -\frac{\omega_n}{\omega_{n+1}} \int_{-\infty}^{\infty} \frac{|f|^n dt}{\left(t^2 + |f|^2\right)^{\frac{n+1}{2}}} f^* \theta_n$$

= $-\frac{2\omega_n}{\omega_{n+1}} \int_{0}^{\pi/2} \cos^{n-1} t \, dt \, f^* \theta_n = -f^* \theta_n$

as required.

4. Stiefel–Whitney currents

The purpose of this section is to identify the cohomology class of a linear dependency current. First we consider the mod 2 case, and recall from Example A.25 of the Appendix that \mathbb{Z}_2 cohomology can be computed using mod 2 integrally flat currents, $\mathcal{F}_{loc}^{mod 2}(X)$.

Theorem 4.1. Given an atomic collection v_1, \ldots, v_m of m = n - q + 1 sections of a real rank n vector bundle F over X the mod 2 linear dependency current $\mathbf{LD}^{\text{mod } 2}(v) \in \mathcal{F}^{\text{mod } 2}_{\text{loc}}(X)$ represents the qth Stiefel–Whitney class $w_q(F) \in H^q(X, \mathbb{Z}_2)$.

Before proving this result, we note that the analog of a theorem of Bott for complex vector bundles and Chern classes is valid for real vector bundles and Stiefel–Whitney classes. Let w(E) := $1+w_1(E)+\ldots+w_m(E)$ denote the total Stiefel–Whitney class of a real rank *m* bundle *E* over *X*. Let *U* denote the universal line bundle on the projectivization $\mathbb{P}(E)$, and let $a := w_1(U) \in H^1(\mathbb{P}(E), \mathbb{Z}_2)$ denote the first Stiefel–Whitney class of *U* on $\mathbb{P}(E)$. Let $\pi : \mathbb{P}(E) \to X$ denote the natural projection, and let $\pi_* : H^{q+m-1}(\mathbb{P}(E), \mathbb{Z}_2) \to H^q(X, \mathbb{Z}_2)$ denote the induced map. Stiefel-Whitney Currents

Lemma 4.2.

$$\pi_*\left((1+a)^{-1}\right) = w(E)^{-1}.$$

Proof. Choose an inner product for E. Let \mathbf{E} denote the pullback of the bundle E to $\mathbb{P}(E)$, and let U^{\perp} denote the orthogonal bundle to $U \subset \mathbf{E}$. The product formula for Stiefel–Whitney classes implies that

$$w(\mathbf{E}) = w(U) w\left(U^{\perp}\right)$$
(4.3)

so that

$$w(U)^{-1} = w(\mathbf{E})^{-1} w(U^{\perp}).$$
 (4.4)

Since the fibre dimension of $\mathbb{P}(E)$ is m-1,

$$\pi_*\left(w_j\left(U^{\perp}\right)\right) = 0 \quad \text{if } j < m-1.$$
(4.5)

Therefore, $\pi_*(w(U^{\perp})) = \pi_*(w_{m-1}(U^{\perp}))$. It remains to show that

$$\pi_*\left(w_{m-1}\left(U^{\perp}\right)\right) = 1.$$
(4.6)

First note that $H^0(X, \mathbb{Z}_2) = \mathbb{Z}_2$ for X connected. One can verify (4.6) by choosing a section α of U^{\perp} and computing that $\pi_*(\text{Div}(\alpha)) \neq 0 \mod 2$. (Note that $\text{Div}(\alpha)$ represents the mod 2 Euler class of U^{\perp} which is equal to the top Stiefel–Whitney class, $w_{m-1}(U^{\perp})$.)

An alternate proof of (4.6) can be given as follows. Equations (4.4) and (4.5) imply that $\pi_*(w_{m-1}(U^{\perp})) = \pi_*(a^{m-1})$. Using the standard fact that if *a* is the nonzero element of $H^1(\mathbb{P}(\mathbb{R}^m), \mathbb{Z}_2)$, then a^{m-1} is the non-zero element of $H^{m-1}(\mathbb{P}(\mathbb{R}^m), \mathbb{Z}_2)$, we conclude that $\pi_*(a^{m-1}) = 1$.

We only need Lemma 4.2 in the special case that $E = \mathbb{R}^m$ is trivial.

Corollary 4.7. Consider the tautological line bundle U on $\mathbb{P}(\mathbb{R}^m)$. Then

$$\pi_* (w_1(U))^j = 0$$
 if $j \neq m-1$

and

$$\pi_* (w_1(U))^{m-1} = 1$$
.

Proof of Theorem 4.1. As in Section 3, let $\nu : \mathbb{R}^m \to F$ denote the bundle map corresponding to the sections ν_1, \ldots, ν_m . Let $\nu : \mathbb{R}^m \to \mathbf{F}$ denote the pullback of ν to the projectivization $\mathbb{P}(\mathbb{R}^m)$. Let $\tilde{\nu}$ denote the restriction of ν to the tautological line bundle $U \subset \mathbb{R}^m$. Then, considering $\text{Div}^{\text{mod } 2}(\tilde{\nu}) \in \mathcal{F}_{\text{loc}}^{\text{mod } 2}(\mathbb{P}(\mathbb{R}^m))$ as a mod 2 current, the linear dependency current is defined to be the current push forward

$$\mathbf{LD}^{\mathrm{mod}\,2}(\nu) := \pi_*\left(\mathrm{Div}^{\mathrm{mod}\,2}\left(\widetilde{\nu}\right)\right) \in \mathcal{F}^{\mathrm{mod}\,2}_{\mathrm{loc}}(X) \;.$$

Now, by Theorem 2.13, the mod 2 divisor $\operatorname{Div}^{\operatorname{mod} 2}(\widetilde{\nu})$ represents the top Stiefel–Whitney class of $H := \operatorname{Hom}(U, \mathbf{F})$ over $\mathbb{P}(\mathbb{R}^m)$. The standard formula for the Stiefel–Whitney classes of a tensor product (see [16]) says that

$$w_n(H) = w_n(U^* \otimes \mathbf{F}) = \sum_{j=0}^n w_{n-j}(\mathbf{F}) (w_1(U))^j$$
.

Reese Harvey and John Zweck

So, by Corollary 4.7, $\pi_*(w_n(H)) = w_{n-m+1}(F)$. Therefore, $LD^{\text{mod } 2}(v)$ represents $w_q(F)$ as desired.

Now we consider the case that q is odd and identify the cohomology class of the \mathcal{O}_F -twisted current $\mathbf{LD}(\nu) \in \widetilde{\mathcal{F}}_{\mathrm{loc}}(X)$ in $H^q(X, \widetilde{\mathbb{Z}})$, where $\widetilde{\mathbb{Z}} = \widetilde{\mathbb{Z}}_F := \mathcal{O}_F \otimes_{\mathbb{Z}_2} \mathbb{Z}$. Consider the short exact triple $0 \to \widetilde{\mathbb{Z}} \xrightarrow{2} \widetilde{\mathbb{Z}} \to \mathbb{Z}_2 \to 0$ and the induced long exact sequence

$$\dots \to H^{q-1}(X, \mathbb{Z}_2) \xrightarrow{\beta} H^q(X, \widetilde{\mathbb{Z}}) \xrightarrow{2} H^q(X, \widetilde{\mathbb{Z}}) \xrightarrow{\rho} H^q(X, \mathbb{Z}_2) \to \dots$$
(4.8)

Define the q th \mathbb{Z} -Stiefel–Whitney class $\widetilde{W}_q \in H^q(X, \mathbb{Z})$ to be the Bockstein of w_{q-1} ,

$$\widetilde{W}_q := \beta \left(w_{q-1} \right) \,. \tag{4.9}$$

Recall from Example A.16 of the Appendix that $\widetilde{\mathbb{Z}}$ -cohomology can be computed using \mathcal{O}_F -twisted integrally flat currents, $\widetilde{\mathcal{F}}_{loc}(X)$.

Theorem 4.10 (q odd). Given an atomic collection v of m = n - q + 1 sections of a real vector bundle $F \to X$ the linear dependency current LD(v) represents the qth \mathbb{Z} -Stiefel-Whitney class, $\widetilde{W}_q(F) \in H^q(X, \mathbb{Z})$, of the bundle F. Moreover, the mod 2 reduction of \widetilde{W}_q equals w_q , i.e., $\rho(\widetilde{W}_q) = w_q$. Hence, $\widetilde{Sq}^1 := \rho \circ \beta$ maps w_{q-1} to w_q for q odd.

Proof. A representative for $\widetilde{W}_q(F)$ can be computed as follows. Choose an atomic collection $\mu = (\mu_1, \ldots, \mu_{m+1})$ of m + 1 = m - (q - 1) + 1 sections. By Theorem 4.1, the mod 2 linear dependency current $\mathbf{LD}^{\mathrm{mod}\,2}(\mu) \in \mathcal{F}_{\mathrm{loc}}^{\mathrm{mod}\,2}(X)$ represents the (q - 1)th Stiefel–Whitney class $w_{q-1}(F) \in H^{q-1}(X, \mathbb{Z}_2)$. Choose an \mathcal{O}_F -twisted current representative $T \in \widetilde{\mathcal{F}}_{\mathrm{loc}}(X)$ of the mod 2 current $\mathbf{LD}^{\mathrm{mod}\,2}(\mu)$. Then, by the definition of β , the current $\frac{1}{2}dT$ represents $\widetilde{W}_q(F)$.

Suppose that the subcollection $\eta = (\mu_1, \dots, \mu_m)$ is atomic and that μ satisfies two additional assumptions described below. Then it is possible to choose T so that

$$\frac{1}{2}dT = \mathbf{L}\mathbf{D}(\eta) \quad \text{on } X . \tag{4.11}$$

Consequently $LD(\eta)$ represents $\widetilde{W}_q(F)$. The theorem now follows from Theorem 3.6. Also note that since the mod 2 reduction of $LD(\eta)$ is $LD^{\text{mod } 2}(\eta)$, Theorem 4.1 implies that $\rho(\widetilde{W}_q) = w_q$.

The \mathcal{O}_F -twisted current T representing $\mathbf{LD}^{\mathrm{mod}2}(\mu)$ can be chosen as follows. First, if q = 1 define T to be the \mathcal{O}_F -twisted generalized function $T := \frac{\sigma}{|\sigma|}$, where σ is the section of $\mathcal{O}_F^{\mathbb{R}}$ associated to η (c.f. Remark 3.9). In this case (4.11) is simply a restatement of Proposition 3.10. Second, if q > 1 is odd, T is defined as follows. Let $\hat{\mu}$ be the induced section of the rank n bundle $\hat{H} = \mathrm{Hom}(NS(\mathbb{R}^{m+1}), \rho^*F)$ over $S(\mathbb{R}^{m+1})$. Embed $\mathbb{R}^m \hookrightarrow \mathbb{R}^{m+1} = \mathbb{R}^m \oplus \mathbb{R}$ and let t be the coordinate on \mathbb{R} . Set $\chi := \frac{t}{|t|} \in L^{\infty}_{\mathrm{loc}}(S(\mathbb{R}^{m+1}))$ and let $p : S(\mathbb{R}^{m+1}) \to \mathbb{P}(\mathbb{R}^{m+1})$ and $\pi : \mathbb{P}(\mathbb{R}^{m+1}) \to X$ denote the projection maps. Assume that the current $\mathrm{Div}(\hat{\mu})$ on $S(\mathbb{R}^m)$ and so $\chi \mathrm{Div}(\hat{\mu})$ is a well defined even \mathcal{O}_F -twisted flat current on $S(\mathbb{R}^{m+1})$. Let $\widetilde{T} := \frac{1}{2}p_*(\chi \mathrm{Div}(\hat{\mu}))$ be the corresponding $\mathcal{O}_{\mathbb{P}(\mathbb{R}^m)}$ -twisted current on $\mathbb{P}(\mathbb{R}^{m+1})$. Since the mod 2 reduction of \widetilde{T} is $\mathrm{Div}^{\mathrm{mod}\,2}(\widetilde{\mu})$ its push forward $T := \pi_*\widetilde{T}$ is an \mathcal{O}_F -twisted current on X whose mod 2 reduction is $\mathrm{LD}^{\mathrm{mod}\,2}(\mu)$.

Next we verify that Equation (4.11) holds for this choice of *T*. Suppose that the codimension n-1 Hausdorff measure of $\text{Zero}(\widehat{\eta})$ is zero. Then, since $\widehat{\eta} = \widehat{\mu} |_{S(\underline{\mathbb{R}}^m)}$, Lemma 4.12 below implies that

$$d(\chi \operatorname{Div}(\widehat{\mu})) = 2i_*\operatorname{Div}(\widehat{\eta}) \quad \text{on } S(\mathbb{R}^{m+1}),$$

where $i: S(\mathbb{R}^m) \hookrightarrow S(\mathbb{R}^{m+1})$ is the inclusion map. Finally, by (3.5.1), this equation pushes forward to give Equation (4.11) on X.

Lemma 4.12. Let $u : X \times \mathbb{R} \to \mathbb{R}^n$ be atomic and suppose that Div(*u*) has locally finite mass. Suppose that the function $v : X \to \mathbb{R}^n$ defined by v(x) = u(x, 0) is atomic and that the codimension n - 1 Hausdorff measure of Zero(*v*) in X is zero. Let t denote the coordinate on \mathbb{R} and define $i : X \hookrightarrow X \times \mathbb{R}$ by i(x) = (x, 0). Then

$$d\left(\frac{t}{|t|}\operatorname{Div}(u)\right) = 2i_*\operatorname{Div}(v) \quad on \ X \times \mathbb{R} .$$
(4.12.1)

Proof. Let θ denote the solid angle kernel on \mathbb{R}^n . First note that since $i_*d = -di_*$, Equation (4.12.1) is the exterior derivative of the degree *n* current equation

$$d\left(\frac{t}{|t|}u^*\theta\right) = \frac{t}{|t|}\operatorname{Div}(u) + 2i_*\left(v^*\theta\right) \quad \text{on } X \times \mathbb{R}.$$

To verify this equation we argue as follows. First, the equation holds on $(X \times \mathbb{R}) \sim (\text{Zero}(v) \times \{0\})$ since it is true if $t \neq 0$ and if t = 0 and $x \notin \text{Zero}(v)$, then $d(\frac{t}{|t|}u^*\theta) = 2[X]u^*\theta = 2i_*(v^*\theta)$. Finally, since the codimension *n* Hausdorff measure of Zero(v) in $X \times \mathbb{R}$ is zero, the Federer Support Theorem for flat currents implies that the equation holds on all of $X \times \mathbb{R}$.

Remark 4.13 (Non-injectivity currents). Let $m = \operatorname{rk} E \leq \operatorname{rk} F = n$ and set q = n - m + 1. In this remark we study the non-injectivity current of a bundle map $v : E \to F$. This is a degree q current which is supported on the set of points of X over which the bundle map v fails to be injective. The mod 2 non-injectivity current, $\mathbf{D}_{\mathrm{NI}}^{\mathrm{mod}\,2}(v)$, is defined by replacing \mathbb{R}^m by E in Definition 3.13. The analog of Theorem 3.14 holds for $\mathbf{D}_{\mathrm{NI}}^{\mathrm{mod}\,2}(v)$. Furthermore, the cohomology class of $\mathbf{D}_{\mathrm{NI}}^{\mathrm{mod}\,2}(v)$ in $H^q(X, \mathbb{Z}_2)$ is $\{w(F) w(E)^{-1}\}_q$, the degree q part of $w(F) w(E)^{-1}$.

If q is odd, the non-injectivity current, $\mathbf{D}_{NI}(\nu)$, is defined as in Definition 3.4. (Note that, in the case that q = n is odd, the current $\mathbf{D}_{NI}(\nu)$ is simply the divisor of the induced section of Hom $(E, F) \rightarrow X$.) This current is an $\mathcal{O}_E \otimes \mathcal{O}_F$ -twisted current on X. The analogs of all the results of Section 3A hold for $\mathbf{D}_{NI}(\nu)$. Furthermore, the cohomology class of $\mathbf{D}_{NI}(\nu)$ in $H^q(X, \widetilde{\mathbb{Z}}_{E \oplus F})$ is $\beta (\{w(F) w(E)^{-1}\}_{q-1})$, the Bockstein of the degree q - 1 part of $w(F) w(E)^{-1}$.

5. Obstruction currents

The Stiefel-Whitney classes were originally defined (see [22, 24]) as obstruction classes. The qth obstruction class of a real rank n vector bundle $F \rightarrow X$ is a cohomology class, which is the obstruction to the existence of a collection of n - q + 1 linearly independent sections of F over the q-skeleton of a cell decomposition of X. It is defined to be the cohomology class of a certain obstruction q-cocycle which is associated to each suitable collection of n - q + 1 sections of F. If this obstruction cocycle is defined for an atomic collection of sections, then, by Poincaré duality, there is a canonical obstruction current defined on X as well. The aim of this section is to show that this obstruction current is equal to the linear dependency current of this special collection of sections.

We begin by recalling the definition of the **obstruction cocycle** as given by Steenrod [21]. Fix $q \in \{1, ..., n\}$. Let $V_{n-q+1}(F) \to X$ be the bundle whose fibre over $x \in X$ is the Stiefel manifold consisting of all (n - q + 1)-tuples of linearly independent vectors of F_x . Choose a smooth locally finite simplicial decomposition K of X and let K' be the first barycentric subdivision of K. Each

barycentrically subdivided q-simplex, a^q , of K is a simplicial subcomplex of K'. In fact, since it is diffeomorphic to a q-ball, a^q is a q-cell. If q is odd or q = n, choose an orientation on each cell. The collection of such (oriented) cells forms a cellular subdivision K_a of K'. Since $\pi_i(V_{n-q+1}(\mathbb{R}^n)) = 0$ for all i < q - 1, there is a section η of $V_{n-q+1}(F)$ over the (q - 1)-skeleton K_a^{q-1} of K_a .

Now

 $\pi_{q-1}\left(V_{n-q+1}\left(\mathbb{R}^n\right)\right) = \begin{cases} \mathbb{Z} & \text{if } q \text{ is odd or } q = n \\ \mathbb{Z}_2 & \text{if } q \text{ is even and } q < n \end{cases}.$

Consequently there is no guarantee that η will extend to a section of $V_{n-q+1}(F)$ over the q-skeleton K_a^q . Steenrod defines a twisted (or mod 2) cellular q-cochain, $w_q(\eta)$, which is zero iff η can be extended over K_a^q . Fix a point x_a in each q-cell a^q of K_a^q . The cochain $w_q(\eta)$ assigns an element $w_q(\eta)(a^q)$ of $\pi_{q-1}(V_{n-q+1}(F_{x_a}))$ to each q-cell a^q . It is defined as follows. Choose a trivialization of F over a^q and let $\psi : V_{n-q+1}(F) \to V_{n-q+1}(F_{x_a})$ be the induced map. Then $w^q(\eta)(a^q)$ is defined to be the homotopy class of $\psi \circ \eta : \partial a^q \to V_{n-q+1}(F_{x_a})$. This class is well defined independent of the choice of trivialization of F. Steenrod shows that $w_q(\eta)$ is a cocycle whose cohomology class in $H^q(X, \pi_{q-1}(V_{n-q+1}(F)))$ is well defined independent of the choice of section η . By definition this class is the qth obstruction class of F. In keeping with the notation of Section 4 the qth obstruction class will be denoted by $w_q(F)$ when q is even and by $\widetilde{W}_q(F)$ when q is odd.

In summary, if q is odd or q = n (resp. q is even and q < n) Steenrod associates a $\pi_{q-1}(V_{n-q+1}(F))$ -twisted cochain (resp. mod 2 cochain), $w_q(\eta)$, to each section η of the Stiefel bundle $V_{n-q+1}(F) \to K_a^{q-1}$. On the other hand, in Section 3 we associated the \mathcal{O}_F -twisted current $\mathbf{LD}(\nu)$ (resp. mod 2 current $\mathbf{LD}^{\text{mod } 2}(\nu)$) to a collection of n - q + 1 sections of the vector bundle $F \to X$. Our goal is to relate these two constructions. We begin by considering the case that q is odd or q = n.

The case that q is odd or q = n.

First note that there is a bundle isomorphism

$$\varphi: \pi_{q-1}(V_{n-q+1}(F)) \longrightarrow \widetilde{\mathbb{Z}}_F$$

defined, in terms of a generator $[\sigma]$ of $\pi_{q-1}(V_{n-q+1}(\mathbb{R}^n))$, as follows. (See [21] Section 25.6, for a definition of the homotopy generator $[\sigma]$.) Fix $x \in X$ and let $\eta : S^{q-1} \to V_{n-q+1}(F_x)$ represent an element of $\pi_{q-1}(V_{n-q+1}(F))_x$. For each frame f of F_x we obtain a map $\psi : V_{n-q+1}(F_x) \to$ $V_{n-q+1}(\mathbb{R}^n)$. Define $\lambda \in \mathbb{Z}$ by $[\psi \circ \eta] = \lambda[\sigma]$ in $\pi_{q-1}V_{n-q+1}(\mathbb{R}^n)$ and let [f] denote the orientation class of the frame f in \mathcal{O}_F . Then $\varphi(\eta)$ is defined to be the class of $([f], \lambda)$ in \mathbb{Z}_F . (Recall that \mathbb{Z}_F is the space of orbits of the \mathbb{Z}_2 -action $\rho([f], \lambda) = (-[f], -\lambda)$ on $\mathcal{O}_F \times \mathbb{Z}$.)

Let $N = \dim X$. Now there is a dual cellular decomposition K_b of K' characterized by the fact that to each q-cell a^q of K_a there is a unique (N - q)-cell, b^{N-q} , of K_b so that the dual of each face of a^q has b^{N-q} as a face. Choose an orientation on each cell of K_b . Note that the intersection of a cell and its dual is the common centerpoint of both cells and that cells a^q and b^{N-q} which are not dual to each other do not intersect. Therefore, the (q - 1)-cells of K_a do not intersect the (N - q)-cells of K_b .

Using the fact that $\pi_i(V_{n-q+1}(\mathbb{R}^n)) = 0$ for i < q-1 we can construct smooth sections ν_1, \ldots, ν_n of $V \to X$ so that for each $q \in \{1, \ldots, n\}$ the linear dependency set of $\nu_1, \ldots, \nu_{n-q+1}$ is a cellular subcomplex $K_b^{N-q}(\nu)$ of K_b^{N-q} . Note that $\nu_1, \ldots, \nu_{n-q+1}$ define a section η of $V_{n-q+1}(F)$ over K_a^{q-1} .

Now the cohomology group $H^*(X, \mathbb{Z}_F)$ can be computed using the smooth $\mathcal{O}_F \otimes \mathcal{O}_X$ -twisted

infinite integral dual cellular chain complex. The (N-q)-dimensional chains of this complex are formal infinite combinations of the form $\sum_{i} \lambda_j b_j^{N-q}$ where the λ_j are $\mathcal{O}_F \otimes \mathcal{O}_X$ -twisted integers

and the b_j^{N-q} are the oriented (N-q)-cells of K_b . Our intermediate goal is to associate to the cocycle $w_q(\eta)$ (defined by the collection of sections v_1, \ldots, v_{n-q+1}) an $\mathcal{O}_F \otimes \mathcal{O}_X$ -twisted integral cellular (N-q)-cycle, $w_{N-q}(v)$, which represents $\widetilde{W}_q(F) \in H^q(X; \mathbb{Z}_F)$. It is defined as follows. First let a^q and b^{N-q} be oriented dual cells and set $\{x\} := a^q \cap b^{N-q}$. Let [a, b] denote the local section of $\mathcal{O}_X \to b^{N-q}$ induced by the decomposition $T_x X \cong T_x a^q \oplus T_x b^{N-q}$. Then

$$w_{N-q}(\nu) := \sum_{j} \lambda_j \, b_j^{N-q} \tag{5.1}$$

where the sum is taken over the cells b_j^{N-q} of the linear dependency subcomplex $K_b^{N-q}(v)$ of v_1, \ldots, v_{n-q+1} , and where λ_j is the $\mathcal{O}_F \otimes \mathcal{O}_X$ -twisted integer

$$\lambda_j := \left[a_j^q, b_j^{N-q}\right] \otimes \varphi\left(w^q(\eta)\left(a_j^q\right)\right) \,.$$

That is, $w_{N-q}(v)$ is the Poincaré dual of $w^q(\eta)$.

Since an \mathcal{O}_X -twisted smooth oriented cellular chain defines a locally rectifiable current, the twisted cellular cycle $w_{N-q}(\nu)$ defines an \mathcal{O}_F -twisted locally integrally flat current on X, which we also denote by $w_{N-q}(\nu)$. Now, by [14] Theorem 3.2, we can choose the sections $\nu_1, \ldots, \nu_{n-q+1}$ so that the induced section $\tilde{\nu}$ of $H \to \mathbb{P}(\mathbb{R}^m)$ is atomic. Then we have the following result.

Theorem 5.2. For the collection of sections v_1, \ldots, v_{n-q+1} described above,

$$\mathbf{LD}(v) = w_{N-q}(v)$$

as \mathcal{O}_F -twisted locally integrally flat currents on X. Consequently, the obstruction class $\widetilde{W}_q(F) \in H^q(X, \widetilde{\mathbb{Z}}_F)$ is the cohomology class of $LD(\nu)$.

Proof. Let m = n - q + 1. By construction v_1, \ldots, v_{m-1} are linearly independent sections of F over $X \sim K_b^{N-q-1}$. Let E^{m-1} denote the oriented span of v_1, \ldots, v_{m-1} over $X \sim K_b^{N-q-1}$ and let G^q denote the orthogonal complement of E^{m-1} in F with respect to some metric on F. Let v_m^{\perp} denote the orthogonal projection of v_m onto G^q over $X \sim K_b^{N-q-1}$. Now we can choose v_1, \ldots, v_{m-1} so that the section v_m^{\perp} is atomic. Then we have the following:

Lemma 5.3.

$$w_{N-q}(v) = \operatorname{Div}\left(v_m^{\perp}\right) \quad \text{over } X \sim K_b^{N-q-1}.$$

Proof of Lemma 5.3. To prove the lemma we need to calculate each twisted integer λ_j of Equation (5.1) in terms of the degree of a certain map between (q-1)-spheres. This was done by Halperin and Toledo [15] as follows. Since K_b^{N-q-1} and K_a^q are disjoint, the bundle G^q is defined over the q-cell a_j^q . Choose a trivialization $\rho: G^q \to \mathbb{R}^q$ of G^q over a_j^q . This induces an orientation on $G^q \to a_j^q$. Let [f] denote the induced orientation on $F = E^{n-q} \oplus G^q$ over a_j^q . Then

$$\lambda_j = \left[a_j^q, b_j^{N-q}\right] \otimes [f] \ n_j$$

where n_i is the degree of the induced map between oriented (q-1)-spheres,

$$n_j := \operatorname{Deg}\left(\rho \circ v_{n-q+1}^{\perp} : \partial a_j^q \to S\left(\mathbb{R}^q\right)\right) \,.$$

The lemma now follows from Theorem 2.8 and Remark 2.9.

Completion of the proof of Theorem 5.2. By Theorem 3.17 and Lemma 5.3

$$w_{N-q}(v) = \mathbf{L}\mathbf{D}(v)$$
 over $X \sim K_h^{N-q-1}$

Let $S := w_{N-q}(v) - \mathbf{LD}(v)$. Then S is a flat current of dimension N - q which is supported on the N - q - 1 dimensional submanifold K_b^{N-q-1} . So, by the Federer Support Theorem for flat currents, [5], S = 0 on X, as required.

The case that q is even and q < n.

In the case that q is even and q < n the obstruction class, $w_q(F)$, is an element of $H^q(X, \mathbb{Z}_2)$. Now the cohomology group $H^q(X, \mathbb{Z}_2)$ can be computed using the smooth infinite mod 2 dual cellular chain complex. The (N - q)-dimensional chains of this complex are formal infinite combinations of the form $\sum \lambda_i b_i^{N-q}$ where $\lambda_i \in \mathbb{Z}_2$ and the b_i^{N-q} are the (unoriented) (N - q)-cells of K_b .

Just as in the case described above, we construct sections $v_1, \ldots v_{n-q+1}$ of F and a mod 2 cellular (N-q)-cycle, $w_{N-q}^{\text{mod }2}(v) = \sum \lambda_j b_j^{N-q}$, which is supported on the linear dependency set of $v_1, \ldots v_{n-q+1}$ and which represents the obstruction class $w_q(F) \in H^q(X, \mathbb{Z}_2)$. This cycle defines a mod 2 locally integrally flat current in the obvious way. Arguing as in the proof of Theorem 5.2, we have the following:

Theorem 5.4. For the collection of sections v_1, \ldots, v_{n-q+1} described above,

$$\mathbf{L}\mathbf{D}^{\mathrm{mod}\,2}(\mathbf{v}) = w_{N-a}^{\mathrm{mod}\,2}(\mathbf{v})$$

as mod 2 locally integrally flat currents on X. Consequently, the obstruction class $w_q(F) \in H^q(X, \mathbb{Z}_2)$ is the cohomology class of $LD^{mod 2}(v)$.

Remark 5.5 (Obstructions to injective bundle maps). The results of this section can be generalized to the case of vector bundle maps $v : E \to F$ where $m = \operatorname{rk} E \leq \operatorname{rk} F$. Let $\operatorname{Hom}^{\times}(E, F) \to X$ denote the bundle of injective bundle maps from E to F. The Steenrod obstruction class of the bundle $\operatorname{Hom}^{\times}(E, F) \to X$ is a degree q = n - m + 1 cohomology class which is the obstruction to the existence of an injective bundle map $v : E \to F$ over the q-skeleton of X. Combining Remark 4.13 with the analogs of Theorems 5.2 and 5.4, we conclude that this class is equal to

- 1. the twisted Euler class \tilde{e} of Hom(E, F) when q = n,
- 2. the degree q part of $w(F) w(E)^{-1}$ when q < n is even, and
- 3. the Bockstein of the degree q 1 part of $w(F) w(E)^{-1}$ when q < n is odd.

6. Higher dependency currents

The aim of this section is to study the currents associated with higher dependencies, c.f. [12]. Let v be an ordered collection of m sections v_1, \ldots, v_m of a rank n bundle $F \to X$. In Section 3 we studied the linear dependency current, $\mathbf{LD}(v)$, which is supported on the set of points x where at least one of the vectors $v_1(x), \ldots, v_m(x)$ depends linearly on the remaining ones. Fix an integer ℓ with max $\{0, m - n\} \le \ell \le m$. In this section we study the higher dependency current, $\mathbf{LD}_{\ell}(v)$, which is supported on the set where at least ℓ of the sections depends linearly on the remaining ones,

i.e., on the set of points over which the induced bundle map $v : \mathbb{R}^m \to F$ has rank $\leq m - \ell$. The higher dependency current $\mathbf{LD}_{\ell}(v)$ has degree $q := \ell(n - m + \ell)$, and so the dimension of $\mathbf{LD}_{\ell}(v)$ decreases as ℓ increases.

The current $\mathbf{LD}_{\ell}(\nu)$ is defined as follows. Let $\pi : G_{\ell}(\mathbb{R}^m) \to X$ denote the trivial Grassmann bundle of unoriented ℓ -dimensional linear subspaces of the trivial bundle $\mathbb{R}^m \to X$, and let $U \subset \mathbb{R}^m$ be the tautological rank ℓ bundle over $G_{\ell}(\mathbb{R}^m)$. The collection ν is called ℓ -**dependency atomic** if the induced section $\tilde{\nu}$ of the bundle $H = \operatorname{Hom}(U, \pi^*F)$ over $G_{\ell}(\mathbb{R}^m)$ is atomic.

The mod 2 higher dependency current, $LD_{\ell}^{mod 2}(\nu)$, is defined by

$$\mathbf{LD}_{\ell}^{\mathrm{mod}\,2}(\nu) := \pi_* \operatorname{Div}^{\mathrm{mod}\,2}(\widetilde{\nu}) \qquad \text{on } X \,. \tag{6.1}$$

In this mod 2 case, orientation issues are irrelevant.

If $m \equiv n \mod 2$, it is possible to push forward the \mathcal{O}_H -twisted current $\operatorname{Div}(\tilde{v})$ by the projection π . This is because there is a canonical isomorphism

$$\mathcal{O}_H \cong \begin{cases} \mathcal{O}_{G_{\ell}(\mathbb{R}^m)} \otimes \pi^* \mathcal{O}_F & \text{if } \ell \text{ is odd,} \\ \mathcal{O}_{G_{\ell}(\mathbb{R}^m)} & \text{if } \ell \text{ is even.} \end{cases}$$

In this case, the **higher dependency current**, $LD_{\ell}(\nu)$, is defined by

$$\mathbf{LD}_{\ell}(\nu) := \pi_* \operatorname{Div}(\widetilde{\nu}) \quad \text{on } X.$$
(6.2)

Note that

$$\mathbf{LD}_{\ell}(\nu) \text{ is } \begin{cases} \text{ an } \mathcal{O}_F - \text{twisted current on } X & \text{ if } \ell \text{ is odd,} \\ \text{ a current on } X & \text{ if } \ell \text{ is even} \end{cases}$$

The following equivalent definition of $\mathbf{LD}_{\ell}(v)$ is often useful (c.f. Remark 3.5). Let ρ : $\widehat{G}_{\ell}(\mathbb{R}^m) \to X$ be the Grassmann bundle of *oriented* ℓ -dimensional linear subspaces of \mathbb{R}^m . Note that the fibers $\widehat{G}_{\ell}(\mathbb{R}^m)$ of $\widehat{G}_{\ell}(\mathbb{R}^m)$ are canonically oriented manifolds. Let \widehat{U} be the canonically oriented tautological bundle over $\widehat{G}_{\ell}(\mathbb{R}^m)$, and let $\widehat{\nu}$ denote the induced section of the bundle $\widehat{H} = \operatorname{Hom}(\widehat{U}, \rho^* F)$ over $\widehat{G}_{\ell}(\mathbb{R}^m)$. Then, if $m \equiv n \mod 2$,

$$\mathbf{LD}_{\ell}(\nu) = \frac{1}{2} \rho_* \operatorname{Div}(\widehat{\nu}) .$$
(6.3)

Proposition 6.4. Let $v : \underline{\mathbb{R}}^m \to F$ be as above and let $\psi : \underline{\mathbb{R}}^m \to \underline{\mathbb{R}}^m$ be a bundle isomorphism. Then the collection of sections corresponding to the bundle map $v \circ \psi : \underline{\mathbb{R}}^m \to F$ is also ℓ -dependency atomic. Furthermore, if $n \equiv m \mod 2$,

$$\mathbf{LD}_{\ell}(\nu \circ \psi) = \begin{cases} \operatorname{sgn} \det(\psi) \ \mathbf{LD}_{\ell}(\nu) & \text{if } \ell \text{ is odd,} \\ \mathbf{LD}_{\ell}(\nu) & \text{if } \ell \text{ is even.} \end{cases}$$

Note. The analogs of the main Theorems 3.6 and 3.14 for (mod 2) linear dependency currents also hold for the (mod 2) higher dependency currents. This fact, together with Proposition 6.4, implies the following corollary.

Corollary 6.5. Let $n \equiv m \mod 2$ and let ℓ be odd. Then there is a locally rectifiable \mathcal{O}_F -twisted current T so that

$$2\mathbf{L}\mathbf{D}_{\ell}(\mathbf{v}) = dT.$$

Reese Harvey and John Zweck

Proof of Proposition 6.4. Let $\mu = \nu \circ \psi$ and let $\Psi : \widehat{G}_{\ell}(\mathbb{R}^m) \to \widehat{G}_{\ell}(\mathbb{R}^m)$ be the diffeomorphism induced by ψ . Arguing as in the proof of Proposition 3.7 it suffices to show that

$$\Psi_* (\operatorname{Div} (\widehat{\mu})) = \operatorname{sgn} \det(D\Psi) \operatorname{Div} (\widehat{\nu}) \quad \text{on } \widehat{G}_\ell (\mathbb{R}^m) , \qquad (6.6)$$

and that

$$\operatorname{sgn} \det(D\Psi) = \begin{cases} \operatorname{sgn} \det(\psi) & \text{if } \ell \text{ is odd,} \\ 1 & \text{if } \ell \text{ is even .} \end{cases}$$
(6.7)

The proof of (6.6) is the same as that of Equation (3.12). Next we prove (6.7). Clearly, it suffices to consider orthogonal linear maps $\psi : \mathbb{R}^m \to \mathbb{R}^m$. Now there is a canonical orientation preserving bundle isomorphism $\phi : \operatorname{Hom}(\widehat{U}, \widehat{U}^{\perp}) \to T\widehat{G}_{\ell}(\mathbb{R}^m)$ defined as follows. Fix $P \in \widehat{G}_{\ell}(\mathbb{R}^m)$ and consider the canonical map $\varphi_P : \operatorname{Hom}(P, P^{\perp}) \hookrightarrow \widehat{G}_{\ell}(\mathbb{R}^m)$ which sends a linear map to its graph. Let Id : $\operatorname{Hom}(P, P^{\perp}) \to T_0 \operatorname{Hom}(P, P^{\perp})$ be the canonical isomorphism. Then $\phi_P := D\varphi_P \circ \operatorname{Id}$ defines ϕ pointwise. Let $\overline{\Psi} : \operatorname{Hom}(P, P^{\perp}) \to \operatorname{Hom}(\psi(P), \psi(P^{-}))$ be the map defined by $\overline{\Psi} :=$ $\varphi_{\psi(P)} \circ \Psi \circ \varphi_P$. Now $\overline{\Psi}(\alpha) = \psi \circ \alpha \circ \psi^{-1}$ is a linear map and so sgn det $(D\Psi) = \operatorname{sgn} \operatorname{det}(\overline{\Psi})$. The result now follows by applying (3.3).

A. The mod 2 cohomology class of $LD_{\ell}^{mod2}(v)$

The goal of this subsection is to identify the mod 2 cohomology class of the mod 2 higher dependency current. Let $w(F) = 1 + w_1(F) + w_2(F) + \ldots + w_n(F)$ denote the total Stiefel-Whitney class of F. The **Shur polynomial**, $\Delta_r^{(\ell)}(w(F)) \in H^{rl}(X, \mathbb{Z}_2)$, is the polynomial in $w_j(F)$ defined by

$$\Delta_r^{(\ell)}(w(F)) := \det(w_{r-i+j}(F))_{1 \le i, j \le \ell} .$$
(6.8)

Theorem 6.9. Let $v : \mathbb{R}^m \to F$ be as above and let $q = \ell(n - m + \ell)$. Then the cohomology class of the mod 2 higher dependency current $\mathbf{LD}_{\ell}^{\text{mod } 2}(v)$ in $H^q(X, \mathbb{Z}_2)$ is the Shur polynomial $\Delta_{n-m+\ell}^{(\ell)}(w(F))$.

Example 6.10. (a) Since $\Delta_1^{(3)}(w(T\mathbb{RP}^4)) \neq 0$, it is not possible to find a collection of six vector fields on \mathbb{RP}^4 so that at each point of \mathbb{RP}^4 at least four of the six vectors are linearly independent.

(b) Since $\Delta_3^{(3)}(w(T\mathbb{RP}^{10})) \neq 0$, for any collection of 10 vector fields on \mathbb{RP}^{10} there is a point of \mathbb{RP}^{10} so that at least 3 of the vectors at that point depend linearly on the remaining ones.

Proof of Theorem 6.9. Let $\pi : G_{\ell}(\underline{\mathbb{R}}^m) \to X$. As in the proof of Theorem 4.1, it suffices to show that

$$\pi_* w_{\ell n}(H) = \Delta_{n-m+\ell}^{(\ell)}(w(F)) \qquad \text{in } H^q(X, \mathbb{Z}_2)$$

Now the standard formula for the Stiefel-Whitney class of a tensor product says that

$$w_{\ell n}(H) = w_{\ell n} \left(U^* \otimes F \right) = \Delta_n^{(\ell)} \left(w(F) w(U)^{-1} \right) \,.$$

(One way to see this is to apply Equation (A.26) of [12] to Problem 7C of [16].) Let $k = m - \ell$. Then, arguing as in the proof of Theorem 4.4 of [12], we see that

$$w_{\ell n}(H) = \Delta_{n-k}^{(\ell)}(w(F)) \pi_*\left(w_k(U^{\perp})^\ell\right).$$

The proof is completed by observing that $w_k(U^{\perp})^{\ell}$ is the generator of $H^{k\ell}(G_{\ell}(\mathbb{R}^{k+\ell}), \mathbb{Z}_2) = \mathbb{Z}_2$.

Remark 6.11 (Mod 2 degeneracy currents). Let $v : E^m \to F^n$ be a bundle map. Fix an integer k with $0 \le k < \min\{m, n\}$ and let $\ell = m - k$. The mod 2 degeneracy current, $\mathbf{D}_k^{\text{mod } 2}(v)$, of the bundle map v is defined as in Equation (6.1), with \mathbb{R}^m replaced by E. It is a degree q = (m-k)(n-k) current which is supported on the set of points over which the bundle map has rank $\le k$. The cohomology class of $\mathbf{D}_k^{\text{mod } 2}(v)$ in $H^q(X, \mathbb{Z}_2)$ is given by $\Delta_{n-k}^{(m-k)}(w(F)w(E)^{-1})$.

Remark 6.12 (Non-surjectivity currents). Next we specialize Remark 6.11 to the case that $m = \operatorname{rk} E \ge \operatorname{rk} F = n$ and k = n - 1 so that q = m - n + 1. The mod 2 non-surjectivity current, $\mathbf{D}_{NS}^{\operatorname{mod} 2}(v) := \mathbf{D}_{n-1}^{\operatorname{mod} 2}(v)$, is supported on the set over which the bundle map v fails to be surjective. The cohomology class of $\mathbf{D}_{NS}^{\operatorname{mod} 2}(v)$ in $H^q(X, \mathbb{Z}_2)$ is given by $\Delta_1^{(q)}(w(F)w(E)^{-1}) = \Delta_q^{(1)}(w(E)w(F)^{-1}) = \{w(E)w(F)^{-1}\}_q$, the degree q part of $w(E)w(F)^{-1}$.

If q is odd, the non-surjectivity current, $\mathbf{D}_{NS}(v) := \mathbf{D}_{n-1}(v)$, is an $\mathcal{O}_E \otimes \mathcal{O}_F$ -twisted current defined as in Equation (6.2). Let $v^* : F^* \to E^*$ be the adjoint map. Then, at least for generic maps, $\mathbf{D}_{NS}(v) = \mathbf{D}_{NI}(v^*)$, where $\mathbf{D}_{NI}(v^*)$ is the non-injectivity current of v^* as defined in Remark 4.13. So, by Remark 4.13, the cohomology class of $\mathbf{D}_{NS}(v)$ in $H^q(X, \mathbb{Z}_{E \oplus F})$ is $\beta(\{w(E) \ w(F)^{-1}\}_{q-1})$, the Bockstein of the degree q - 1 = m - n part of $w(E) \ w(F)^{-1}$.

B. The integer cohomology class of $LD_{\ell}(v)$. The case $n \equiv m \mod 2$

The aim of this subsection is to identify the (twisted) integer cohomology class $[\mathbf{LD}_{\ell}(\nu)]$ of the higher dependency current $\mathbf{LD}_{\ell}(\nu)$, which is defined whenever $n \equiv m \mod 2$. The torsion-free part of $[\mathbf{LD}_{\ell}(\nu)]$ is well known. If ℓ is even it is a certain Shur polynomial in the total Pontrjagin class p(F) (see, for example, [12], Theorem 6.9) while if ℓ is odd it is zero, by Corollary 6.5 above. The mod 2 reduction of $[\mathbf{LD}_{\ell}(\nu)]$ is given in Theorem 6.9 above. We will prove that the (twisted) integer class of $\mathbf{LD}_{\ell}(\nu)$ is the sum of its torsion-free part and 2-torsion term, $T_{n-m+\ell}^{(\ell)}(\widetilde{W})$, defined below. This result builds on work of Ronga [20] who showed that the integer class of $\mathbf{LD}_{\ell}(\nu)$ is determined by its mod 2 and rational reductions. Our contribution is to explicitly identify the 2-torsion term as a certain polynomial in the Pontrjagin and twisted integral Stiefel–Whitney classes of F.

Throughout this subsection, \mathbb{Z} denotes the \mathcal{O}_F -twisted integers, $\mathbb{Z} := \mathcal{O}_F \otimes_{\mathbb{Z}_2} \mathbb{Z}$. Let $p_i(F) \in H^{4i}(X, \mathbb{Z})$ denote the *i*th integral Pontrjagin class of F and $\widetilde{W}_{2j+1}(F) \in H^{2j+1}(X, \mathbb{Z})$ the (2j+1)th twisted integral Stiefel–Whitney class, defined by (4.9). Recall that $\rho(\widetilde{W}_{2j+1}) = w_{2j+1}$. Note that, since $2\widetilde{W}_{2j+1} = 0$, the subring of $H^*(X, \mathbb{Z}) \oplus H^*(X, \mathbb{Z})$ generated by $p_i(F)$ and $\widetilde{W}_{2j+1}(F)$ is commutative. Also note that the product of two elements of $H^*(X, \mathbb{Z})$ is an element of $H^*(X, \mathbb{Z})$. Let \widetilde{W}_{2j} denote the *formal* square root of the *j*th Pontrjagin class, i.e., $\widetilde{W}_{2j} := \sqrt{p_j}$. We make this definition because w_{2j}^2 is the mod 2 reduction of p_j (see [16] Problem 15A). Of course, the formal symbol \widetilde{W}_{2j} has no cohomological meaning.

Definition 6.13. Let S_{ℓ} denote the symmetric group on ℓ elements. Define $\tau \in S_{\ell}$ by $\tau(i) = \ell + 1 - i$, and note that $\tau^2 = \text{Id}$. Let $R : S_{\ell} \to S_{\ell}$ be the involution defined by $R(\sigma) = \tau \sigma^{-1} \tau$. Define an index set $\mathcal{J} \subset S_{\ell}$ by

 $\mathcal{J} := \{ \sigma \in S_{\ell} : R(\sigma) = \sigma \text{ and, if } \ell \text{ is even, then } \sigma(i) \neq i \mod 2 \text{ for some } i. \}$

Set $r := n - m + \ell$ and note that $\ell \equiv r \mod 2$. Then we define

$$T_r^{(\ell)}\left(\widetilde{W}\right) := \sum_{\sigma \in \mathcal{J}} \prod_{i=1}^{\ell} \widetilde{W}_{r+i-\sigma(i)}.$$

Lemma 6.14. Suppose $\ell \equiv r \mod 2$. Then

(1) $T_r^{(\ell)}(\widetilde{W})$ is a polynomial in the p_i and \widetilde{W}_{2j+1} .

(2) If ℓ is odd (resp. even), then each term of $T_r^{(\ell)}(\widetilde{W})$ is of odd (resp. even) degree in the variables \widetilde{W}_{2j+1} . Therefore, $T_r^{(\ell)}(\widetilde{W}(F))$ is an element of $H^q(X, \widetilde{\mathbb{Z}})$ (resp. $H^q(X, \mathbb{Z})$), where $q = \ell r$.

(3) Each term of the polynomial $T_r^{(\ell)}(\widetilde{W})$ has a factor of the form \widetilde{W}_{2j+1} . Therefore, the class $T_r^{(\ell)}(\widetilde{W}(F))$ is a torsion class of order 2.

In the case that ℓ is even set $\ell = 2\ell_0$ and $r = 2r_0$. The main result is the following:

Theorem 6.15. The(twisted) integer cohomology class of the higher dependency current $LD_{\ell}(v)$ is

$$[\mathbf{LD}_{\ell}(\nu)] = \begin{cases} T_r^{(\ell)}(\widetilde{W}(F)) & \text{in } H^q(X, \widetilde{\mathbb{Z}}) \text{ when } \ell \text{ is odd,} \\ \Delta_{r_0}^{(\ell_0)}(p(F)) + T_r^{(\ell)}(\widetilde{W}(F)) & \text{in } H^q(X, \mathbb{Z}) \text{ when } \ell \text{ is even }. \end{cases}$$

Remark 6.16. For ℓ even (resp. ℓ odd) let $\iota^* : H^*(X, \mathbb{Z}) \to H^*(X, \mathbb{R})$ (resp. $\iota^* : H^*(X, \widetilde{\mathbb{Z}}) \to H^*(X, \widetilde{\mathbb{R}})$) denote the usual coefficient homomorphism, and let $\rho^* : H^*(X, \mathbb{Z}) \to H^*(X, \mathbb{Z}_2)$ (resp. $\rho^* : H^*(X, \widetilde{\mathbb{Z}}) \to H^*(X, \mathbb{Z}_2)$) denote mod 2 reduction. Then

$$\iota^* \left([\mathbf{LD}_{\ell}(\nu)] \right) = \begin{cases} 0 & \text{if } \ell \text{ is odd,} \\ \Delta_{r_0}^{(\ell_0)}(p(F)) & \text{if } \ell \text{ is even .} \end{cases}$$
(6.16.1)

and

$$\rho^*\left([\mathbf{L}\mathbf{D}_{\ell}(\nu)]\right) = \left[\mathbf{L}\mathbf{D}_{\ell}^{\text{mod }2}(\nu)\right] = \Delta_r^{(\ell)}(w(F)) .$$
(6.16.2)

These two formulae are well known (see [20]) and will be used to prove the more general result of Theorem 6.15.

Example 6.17.

(1) If $\ell = 2$, then $r = 2r_0 = n - m + 2$ and $[\mathbf{LD}_2(\nu)] = p_{r_0} + \widetilde{W}_{r-1}\widetilde{W}_{r+1}$. (2) If $\ell = r = 3$, then $[\mathbf{LD}_3(\nu)] = p_1\widetilde{W}_5 + p_2\widetilde{W}_1 + \widetilde{W}_3^3 + \widetilde{W}_1\widetilde{W}_3\widetilde{W}_5$. (3) If $\ell = r = 4$, then $[\mathbf{LD}_4(\nu)]$ $= p_2^2 - p_1p_3 + p_1\widetilde{W}_5\widetilde{W}_7 + p_2\left(\widetilde{W}_3\widetilde{W}_5 + \widetilde{W}_1\widetilde{W}_7\right) + p_3\widetilde{W}_1\widetilde{W}_3 + \left(\widetilde{W}_1\widetilde{W}_5 + \widetilde{W}_3^2\right)\left(\widetilde{W}_3\widetilde{W}_7 + \widetilde{W}_5^2\right)$.

Remark 6.18. The following equivalent definition of $T_r^{(\ell)}(\widetilde{W})$ will be useful. Define an $\ell \times \ell$ matrix (a_{ij}) by

$$a_{ij} := W_{r+i-j} \qquad 1 \le i, j \le \ell$$
 (6.18.1)

Note that the matrix (a_{ij}) is symmetric under reflection in the antidiagonal $i + j = \ell + 1$, i.e., $a_{\ell+1-j}, \ell+1-i = a_{ij}$. Let \mathcal{I} denote the collection of subsets I of the index set $\{(i, j) : 1 \le i, j \le \ell\}$ which satisfy the following three properties:

- 1. For each *i* (resp. *j*) in $\{1, \ldots, \ell\}$ there there is exactly one element *j* (resp. *i*) of $\{1, \ldots, \ell\}$ so that $(i, j) \in I$.
- 2. The subset *I* is symmetric under reflection in the antidiagonal $i + j = \ell + 1$, i.e., $(i, j) \in I$ iff $(\ell + 1 i, \ell + 1 j) \in I$, and

3. If ℓ is even, then there is at least one element $(i, j) \in I$ for which $i \neq j \mod 2$.

Then

$$T_r^{(\ell)}(\widetilde{W}) = \sum_{I \in \mathcal{I}} \prod_{(i,j) \in I} \widetilde{W}_{r+i-j}.$$
(6.18.2)

To see that Definition 6.13 and (6.18.2) agree, note that each $\sigma \in \mathcal{J}$ defines a subset $I(\sigma)$ of $\{(i, j) : 1 \le i, j \le \ell\}$ by $I(\sigma) := \{(i, \sigma(i)) : 1 \le i \le \ell\}$. Furthermore, the set associated with $R(\sigma)$ is the reflection in the line $i + j = \ell + 1$ of the subset associated with σ .

Proof of Lemma 6.14. Let $\sigma \in \mathcal{J}$. Then, since $R(\sigma) = \sigma$, we have $(i, j) \in I(\sigma)$ iff $(\tau(j), \tau(i)) \in I(\sigma)$, where $\tau(i) = \ell + 1 - i$. So, since $a_{\tau(j)\tau(i)} = a_{ij}$,

$$T_r^{(\ell)}\left(\widetilde{W}\right) \;=\; \sum_{\sigma \in \mathcal{J}} \prod_{i \in U(\sigma)} \widetilde{W}_{r+i-\sigma(i)}^2 \prod_{i \in \Delta(\sigma)} \widetilde{W}_{r+i-\sigma(i)} \;,$$

where $U(\sigma) = \{i : i + \sigma(i) < \ell + 1\}$ and $\Delta(\sigma) = \{i : i + \sigma(i) = \ell + 1\}$. Note that $|\Delta(\sigma)| \equiv \ell \mod 2$. Conclusions (1, 2, and 3) now follow from the fact that if $i \in \Delta(\sigma)$, then $r + i - \sigma(i) = r - \ell + 2i - 1$ is odd. Conclusion 4 follows from the fact that $|\Delta(\sigma)| \neq 0$ in the case that ℓ is odd, and from the definition of \mathcal{J} in the case that ℓ is even.

Proof of Theorem 6.15. By naturality we can reduce to the case in which the bundle $F \to X$ is the tautological rank *n* bundle *U* over a sufficiently high dimensional approximation, $G_n(\mathbb{R}^N)$, to the classifying space $G_n(\mathbb{R}^\infty)$. Now, if *N* is large enough, the torsion subgroup of $H^q(G_n(\mathbb{R}^N), \mathbb{Z})$ is a direct sum of cyclic groups of order 2 (see [2]). Furthermore, choosing *N* to be odd, $\mathcal{O}_{TG_n(\mathbb{R}^N)} \cong$ \mathcal{O}_U and so $H^*(G_n(\mathbb{R}^N), \mathbb{Z}) \cong H_*(G_n(\mathbb{R}^N), \mathbb{Z})$. So, by the universal coefficient theorem, the torsion subgroup of $H^q(G_n(\mathbb{R}^N), \mathbb{Z})$ and $H^q(G_n(\mathbb{R}^N), \mathbb{Z})$ are completely determined by their mod 2 and real reductions. So, setting

$$Q_r^{(\ell)} := \begin{cases} T_r^{(\ell)}(\widetilde{W}(F)) & \text{if } \ell \text{ is odd,} \\ \Delta_{r_0}^{(\ell_0)}(p(F)) + T_r^{(\ell)}(\widetilde{W}(F)) & \text{if } \ell \text{ is even,} \end{cases}$$

it suffices to prove that

$$\iota^*(\mathcal{Q}_r^{(\ell)}) = \begin{cases} 0 & \text{if } \ell \text{ is odd,} \\ \Delta_{r_0}^{(\ell_0)}(p(F)) & \text{if } \ell \text{ is even,} \end{cases}$$
(6.19)

and

$$\rho^*\left(\mathcal{Q}_r^{(\ell)}\right) = \Delta_r^{(\ell)}(w(F)), \qquad (6.20)$$

that is, that $Q_r^{(\ell)}$ and $[\mathbf{LD}_{\ell}(\nu)]$ have the same torsion-free part and mod 2 reduction.

Now (6.19) follows immediately from Lemma 6.14 (3). Let $w(\sigma) := \prod_{i=1}^{\ell} w_{r+i-\sigma(i)}$. Then, since the matrix $a_{ij} = w_{r+i-j}$ is symmetric under reflection in the antidiagonal,

$$w(R(\sigma)) = w(\sigma) . \tag{6.21}$$

In the case that ℓ is odd, we verify (6.20) by observing that, by Definition 6.13, $\Delta_r^{(\ell)}(w(F)) - \rho^*(Q_r^{(\ell)}) = \sum_{\sigma \notin \mathcal{J}} w(\sigma)$, which is zero, since, by (6.21), the sum is a sum of terms of the form $w(\sigma) + w(R(\sigma)) = 2w(\sigma) = 0$.

Reese Harvey and John Zweck

Finally, we verify (6.20) in the case that $\ell = 2\ell_0$ is even. Let $\psi : S_{\ell_0} \to S_{\ell}$ be the injection defined for $j \in \{1, 2, ..., \ell_0\}$ by

$$\psi(\eta)(2j-1) := 2\eta(j) - 1$$
 and $\psi(\eta)(2j) := 2\eta(j)$.

The map ψ can be interpreted as follows. Let (b_{ij}) denote the $\ell_0 \times \ell_0$ matrix $b_{ij} := p_{r_0+i-j}$ and let $C(\eta) = \{b_{j,\eta(j)} : 1 \le j \le \ell_0\}$ be the subset of entries of (b_{ij}) defined by $\eta \in S_{\ell_0}$. Then $C(\psi(\eta)) = \{a_{i,\psi(\eta)(i)} : 1 \le i \le \ell\}$ is the set of those entries of (a_{ij}) obtained from $C(\eta)$ by replacing each element $b_{j,\eta(j)}$ of $C(\eta)$ by the diagonal entries of the corresponding 2×2 submatrix $\begin{pmatrix} w_{2(r_0+j-\eta(j))} & 0 \\ 0 & w_{2(r_0+j-\eta(j))} \end{pmatrix}$ of the matrix (a_{ij}) . So, since $\rho^*(p_j) = w_{2j}^2$, the mod 2 reduction of the product of the elements of $C(\eta)$ equals the product, $w(\psi(\eta))$, of the elements of $C(\psi(\eta))$. Summing over $\eta \in S_{\ell_0}$ we conclude that

$$\rho^*\left(\Delta_{r_0}^{(\ell_0)}(p(F))\right) = \sum_{\sigma \in \psi(S_{\ell_0})} w(\sigma) + \sum_{\sigma \in \psi(S_$$

Now let $\mathcal{K} := S_{\ell} \sim (\psi(S_{\ell_0}) \cup \mathcal{J})$. To verify (6.20) it suffices to show that

$$\sum_{\sigma \in \mathcal{K}} w(\sigma) = 0.$$
(6.22)

To prove (6.22) we study the index set \mathcal{K} . Define $\beta \in S_{\ell}$ by

$$\beta(2j-1) = 2j$$
 and $\beta(2j) = 2j-1$ for $j \in \{1, ..., \ell_0\}$.

and define $P: S_{\ell} \to S_{\ell}$ by $P(\sigma) := \beta \sigma \beta$. The involution P can be interpreted as follows. First, each entry a_{ij} of the $\ell \times \ell$ matrix (a_{ij}) has a pair $P(a_{ij})$ defined as follows. Partition (a_{ij}) into 2×2 submatrices. Let $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be one such submatrix. Then P(a) = d and P(b) = c. Let $C(\sigma)$ denote the set of entries of (a_{ij}) defined by $\sigma \in S_{\ell}$. Then $C(P(\sigma)) = C(\sigma)$. The pairing involution P is introduced because

$$\psi\left(S_{\ell_0}\right) = \{\sigma \in S_{\ell} : P(\sigma) = \sigma \text{ and } i \equiv \sigma(i) \text{ mod } 2 \text{ for all } i\}.$$
(6.23)

Define

$$\mathcal{K}_R = \{ \sigma \in \mathcal{K} : R(\sigma) \neq \sigma \}$$
 and $\mathcal{K}_P = \{ \sigma \in \mathcal{K} : R(\sigma) = \sigma \text{ and } P(\sigma) \neq \sigma \}.$

We claim that \mathcal{K} is the disjoint union $\mathcal{K} = \mathcal{K}_R \cup \mathcal{K}_P$. To see this choose $\sigma \in \mathcal{K} \sim \mathcal{K}_R$. Then, since $\sigma \notin \mathcal{J}$, $i \equiv \sigma(i) \mod 2$ for all *i*. Therefore, since $\sigma \notin \psi(S_{\ell_0})$, Equation (6.23) implies that $P(\sigma) \neq \sigma$, as required.

Note that since the pair of the reflection of an entry of (a_{ij}) is the reflection of the pair of that entry,

$$R(P(\sigma)) = P(R(\sigma)).$$
(6.24)

Hence, *R* preserves the decomposition $S_{\ell} = \psi(S_{\ell_0}) \cup \mathcal{J} \cup \mathcal{K}_R \cup \mathcal{K}_P$. Now, since the involution $R : \mathcal{K}_R \to \mathcal{K}_R$ has no fixed points, it follows that $\sum_{\sigma \in \mathcal{K}_R} w(\sigma) = 0$, since it is the sum of terms of the form $w(\sigma) + w(R(\sigma)) = 2w(\sigma) = 0$. Finally, by (6.24), $P : \mathcal{K}_P \to \mathcal{K}_P$ is an involution with no fixed points, and once again $\sum_{\sigma \in \mathcal{K}_P} w(\sigma) = 0$. Hence, (6.22) holds, as desired.

7. Applications

In this section we apply the general results of the previous sections to study singularities of projections and singularities of maps (c.f. [12]). The results of this section hold whenever the projections and maps in question are atomic, by which we mean that the induced section of $\text{Hom}(U, F) \rightarrow G_{\ell}(E)$ is atomic. This hypothesis is assumed throughout. In particular, in the real analytic case we simply require that the degeneracy subvarieties of the map have codimension greater than or equal to the expected codimension in X, (see [12], Propostion 2.14).

A. Singularities of projections

Let $j: X \to \mathbb{R}^N$ be an immersion of a smooth *m*-manifold. Fix an integer n < N and let $P: \mathbb{R}^N \to \mathbb{R}^n$ be a linear map. We study the singularities of the smooth projection $\widehat{P} = P \circ j: X \to \mathbb{R}^n$. Fix an integer *k* with $0 \le k < \min\{m, n\}$. The *k*th **mod 2 degeneracy current** of the projection *P* on *X* is defined to be $\mathbf{D}_k^{\text{mod 2}}(P) := \mathbf{D}_k^{\text{mod 2}}(d\widehat{P})$ (c.f. Remark 6.11). This current is a degree q := (m - k)(n - k) current which is supported on the set where the differential $d\widehat{P}: TX \to \mathbb{R}^n$ has rank $\le k$. By Remark 6.11,

$$\left[\mathbf{D}_{k}^{\text{mod }2}(P)\right] = \Delta_{n-k}^{(m-k)}\left(w(TX)^{-1}\right) = \Delta_{m-k}^{(n-k)}(w(TX)) \quad \text{in } H^{q}\left(X, \mathbb{Z}_{2}\right) \,. \tag{7.1}$$

If $n \equiv m \mod 2$, the *k*th **degeneracy current** of the projection *P* on *X*, $\mathbf{D}_k(P) := \mathbf{D}_k(d\widehat{P})$ can also be defined [as in (6.2)]. Let $(d\widehat{P})^* : \mathbb{R}^n \to T^*X$ denote the adjoint map. Then, at least for generic *P*, $\mathbf{D}_k(P) = \mathbf{D}_k((d\widehat{P})^*)$. Therefore, by Theorem 6.15,

$$[\mathbf{D}_{k}(P)] = \begin{cases} T_{m-k}^{(n-k)}(\widetilde{W}(T^{*}X)) & \text{in } H^{q}(X,\widetilde{\mathbb{Z}}), \text{ when } n-k \text{ is odd,} \\ \Delta_{m_{0}-k_{0}}^{(n_{0}-k_{0})}(p(T^{*}X)) + T_{m-k}^{(n-k)}(\widetilde{W}(T^{*}X)) & \text{in } H^{q}(X,\mathbb{Z}), \text{ when } n-k \text{ is even,} \end{cases}$$

where $2(n_0 - k_0) = n - k$ and $2(m_0 - k_0) = m - k$.

Example 7.2 (Tangential Stiefel–Whitney classes). Fix $1 \le q \le m$ and let $P : \mathbb{R}^N \to \mathbb{R}^{m-q+1}$ be linear. The mod 2 **non–submersion current** of the projection P on X is defined by $\mathbf{D}_{\mathrm{NS}}^{\mathrm{mod}\,2}(P) := \mathbf{D}_{m-q}^{\mathrm{mod}\,2}(d\widehat{P})$ on X. This is a degree q current which is supported on the subset of X on which the map $\widehat{P} : X \to \mathbb{R}^{m-q+1}$ fails to be a submersion. By (7.1),

$$\left[\mathbf{D}_{\mathrm{NS}}^{\mathrm{mod}\,2}(P)\right] = w_q(TX) \qquad \text{in } H^q(X,\mathbb{Z}_2) \ .$$

Furthermore, if q is odd, then the non-submersion current, $\mathbf{D}_{NS}(P)$, can also be defined and, by Remark 6.12,

$$[\mathbf{D}_{\rm NS}(P)] = \widetilde{W}_q(TX) \qquad \text{in } H^q(X, \mathbb{Z}) .$$

Example 7.3 (Normal Stiefel–Whitney classes). Fix $1 \le q \le N - m$ and let $P : \mathbb{R}^N \to \mathbb{R}^{m+q-1}$ be linear. The mod 2 **non–immersion current** of the projection P on X is defined by $\mathbf{D}_{\mathrm{NI}}^{\mathrm{mod}\,2}(P) := \mathbf{D}_{m-1}^{\mathrm{mod}\,2}(d\widehat{P})$ on X. This is a degree q current which is supported on the subset of X on which the map $\widehat{P} : X \to \mathbb{R}^{m+q-1}$ fails to be a immersion. By (7.1),

$$\left[\mathbf{D}_{\mathrm{NI}}^{\mathrm{mod}\,2}(P)\right] = w_q(NX) \qquad \text{in } H^q(X,\mathbb{Z}_2)$$

where NX is the normal bundle to X in \mathbb{R}^N . Furthermore, if q is odd, then the non-immersion current, $\mathbf{D}_{NI}(P)$, can also be defined and, by Remark 4.13,

$$[\mathbf{D}_{\mathrm{NI}}(P)] = \widetilde{W}_q(NX) \qquad \text{in } H^q(X,\widetilde{\mathbb{Z}}) \ .$$

B. Singularities of maps

Let X and Y be smooth manifolds of dimensions m and n respectively, and let $f : X \to Y$ be a smooth mapping. Let $0 \le k < \min\{m, n\}$. The kth mod 2 degeneracy current of the map f is defined to be $\mathbf{D}_k^{\text{mod }2}(f) := \mathbf{D}_k^{\text{mod }2}(df)$. This is a degree q = (m - k)(n - k) current supported on the set where $df : TX \to TY$ has rank $\le k$. By Remark 6.11,

$$\left[\mathbf{D}_{k}^{\text{mod }2}(f)\right] = \Delta_{n-k}^{(m-k)} \left(f^{*}(w(TY)) w(TX)^{-1}\right) \quad \text{in } H^{q}(X, \mathbb{Z}_{2}) .$$
(7.4)

Example 7.5 (Non-submersion currents). Suppose that $m = \dim X \ge \dim Y = n$ and let q = m - n + 1. Then, by Remark 6.12, the cohomology class of the mod 2 non-submersion current $\mathbf{D}_{NS}^{\text{mod } 2}(f) := \mathbf{D}_{n-1}^{\text{mod } 2}(f)$ is

$$\left[\mathbf{D}_{\mathrm{NS}}^{\mathrm{mod}\,2}(f)\right] = \left\{w(TX) f^*\left(w(TY)^{-1}\right)\right\}_{\mathrm{deg}\,q} \qquad \text{in } H^q\left(X,\mathbb{Z}_2\right) \,,$$

and, if q is odd,

$$[\mathbf{D}_{\rm NS}(f)] = \beta \left(\left\{ w(TX) f^* \left(w(TY)^{-1} \right) \right\}_{\deg q - 1} \right) \qquad \text{in } H^q \left(X, \widetilde{\mathbb{Z}} \right) \,.$$

Example 7.6 (Non-immersion currents). Suppose that $m = \dim X \le \dim Y = n$ and let q = n - m + 1. Then, by Remark 4.13, the cohomology class of the mod 2 non-immersion current $\mathbf{D}_{\text{NI}}^{\text{mod } 2}(f) := \mathbf{D}_{m-1}^{\text{mod } 2}(f)$ is

$$\left[\mathbf{D}_{\mathrm{NI}}^{\mathrm{mod}\,2}(f)\right] = \left\{f^*w(TY)\,w(TX)^{-1}\right\}_{\mathrm{deg}\,q} \qquad \text{in } H^q(X,\mathbb{Z}_2) \ ,$$

and, if q is odd,

$$[\mathbf{D}_{\mathrm{NI}}(f)] = \beta \left(\left\{ f^* w(TY) \, w(TX)^{-1} \right\}_{\deg q - 1} \right) \qquad \text{in } H^q \left(X, \widetilde{\mathbb{Z}} \right) \,.$$

A. Appendix. Computing cohomology with currents

This appendix is included for two reasons; first for the sake of completeness, second for although the approach taken here is both simple and natural (via standard sheaf theory) it does not appear in the geometric measure theory literature.

Definition A.1. The complex

$$0 \to \mathcal{S} \to \mathcal{F}^0 \xrightarrow{d} \mathcal{F}^1 \xrightarrow{d} \dots \xrightarrow{d} \mathcal{F}^n \to 0$$

of sheaves is called an acyclic resolution of the sheaf S if

- 1. the complex is exact, and
- 2. each sheaf \mathcal{F}^p is acyclic, i.e., $H^j(X, \mathcal{F}^p) = 0$ for j = 1, ...

The basic result is that cohomology with coefficients in S can be computed from such a resolution. That is,

$$H^{p}(X, \mathcal{S}) = \frac{\{\varphi \in \Gamma(X, \mathcal{F}^{p}) : d\varphi = 0\}}{d\Gamma(X, \mathcal{F}^{p-1})}$$

and

$$H^{p}_{\rm cpt}(X,S) = \frac{\left\{\varphi \in \Gamma_{\rm cpt}\left(X,\mathcal{F}^{p}\right) : d\varphi = 0\right\}}{d\Gamma_{\rm cpt}\left(X,\mathcal{F}^{p-1}\right)}$$

A classical reference for this and other standard results from sheaf theory is Godement [8].

In this paper the cases and coefficient sheaves of most interest are:

The integer case with coefficient sheaf \mathbb{Z} , the sheaf of germs of locally constant integer valued functions (see Corollary A.13).

The mod 2 case with coefficient sheaf $\mathbb{Z}_2 := \mathbb{Z} / 2\mathbb{Z}$ (see Example A.25), and

The twisted integer case with coefficient sheaf $\widetilde{\mathbb{Z}} := \mathbb{Z} \otimes_{\mathbb{Z}_2} \mathcal{O}_V$, where \mathcal{O}_V is the orientation sheaf of a real vector bundle $V \to X$ (see Example A.16).

Throughout this appendix X is a C^{∞} (paracompact) *n*-dimensional manifold. Let \mathcal{O}_{TX} or \mathcal{O}_X denote the orientation sheaf of X and let $\widetilde{\mathbb{Z}}_X := \mathbb{Z} \otimes_{\mathbb{Z}_2} \mathcal{O}_X$ and $\widetilde{\mathbb{R}}_X := \mathbb{R} \otimes_{\mathbb{Z}_2} \mathcal{O}_X$.

Example A.2 (Differential forms). Let \mathcal{E} denote the sheaf of germs of C^{∞} differential *p*-forms with *d* taken to be exterior differentiation. Then

$$H^p(X,\mathbb{R}) = \frac{\{\varphi \in \mathcal{E}^p(X) : d\varphi = 0\}}{d\mathcal{E}^{p-1}(X)} \,.$$

Each \mathcal{E}^p is acyclic because \mathcal{E}^p is fine (i.e., there exists a partition of unity). Also the sequence, $0 \to \mathbb{R} \to \mathcal{E}^0 \to \ldots \to \mathcal{E}^n \to 0$ is exact by the Poincaré lemma for exterior differentiation.

Example A.3 (Currents). Let \mathcal{D}'^p denote the sheaf of germs of degree *p* currents (defined in Section 2). Then

$$0 o \mathbb{R} o \mathcal{D'}^0 o \mathcal{D'}^1 o \ldots o \mathcal{D'}^n o 0$$

is an acyclic resolution of \mathbb{R} and hence can be used to compute real cohomology $H^p(X, \mathbb{R})$. Note that $\mathcal{E}^p(X) \subset \mathcal{D}'^p(X)$, i.e. each *p*-form, φ , is a current of degree *p*, defined by $\varphi(\psi) \equiv \int_X \varphi \wedge \psi$, for all twisted n - p forms, ψ .

The spaces $\mathcal{R}^p_{loc}(X) \subset \mathcal{D}'^p(X)$ of locally rectifiable currents on X, of degree p or dimension n - p, are very natural for computing \mathbb{Z} -cohomology. However they are not closed under exterior differentiation. There are several ways to remedy this defect (see Examples A.4, A.5 and A.8 below).

Examples of locally rectifiable currents, $T \subset \mathcal{R}^p_{loc}(X)$, can be constructed as follows. Choose an oriented submanifold S of X, which has an orientable neighborhood in X, and fix an orientation on the neighborhood. Then $T(\psi) \equiv \int_S \overline{\psi}$ defines $T \in \mathcal{R}^p_{loc}(X)$, where ψ is an arbitrary \mathcal{O}_X twisted n - p form on X with compact support and where $\overline{\psi}$ denotes the corresponding (untwisted) n - p-form on a neighborhood of S.

Example A.4 (Twisted Singular chains). Let $\tilde{C}^p \subset \mathcal{R}^p_{loc}$ denote the subsheaf of germs of currents arising from integration of (twisted) forms over finite sums of simplicies with oriented neighborhoods as discussed above. Then, using exterior differentiation or the usual boundary operator,

$$0 \to \mathbb{Z} \to \widetilde{\mathcal{C}}^0 \to \widetilde{\mathcal{C}}^1 \to \ldots \to \widetilde{\mathcal{C}}^n \to 0$$

is an acyclic resolution of \mathbb{Z} , and hence can be used to compute integer cohomology.

Example A.5 (Integral currents). The space $I_{loc}^{p}(X)$ of (locally) integral currents of degree p on X is defined to be $\{T \in \mathcal{R}_{loc}^{p}(X) : dT \in \mathcal{R}_{loc}^{p+1}(X)\}$. Let \mathcal{I}^{p} denote the sheaf of germs of

(locally) integral currents with d taken to be exterior differentiation. Then

$$0 \to \mathbb{Z} \to \mathcal{I}^0 \to \mathcal{I}^1 \to \ldots \to \mathcal{I}^n \to 0$$

is an acyclic resolution of \mathbb{Z} , and hence can be used to compute integer cohomology.

Example A.6 (\mathcal{O}_X -twisted currents). Let \mathcal{D}'_k denote the sheaf of germs of k dimensional \mathcal{O}_X -twisted currents on X. Then $0 \to \widetilde{\mathbb{R}}_X \to \mathcal{D}'_n \to \mathcal{D}'_{n-1} \to \ldots \to \mathcal{D}'_0 \to 0$ is an acyclic resolution of $\widetilde{\mathbb{R}}_X$, and hence, can be used to compute $H^p(X, \widetilde{\mathbb{R}}_X)$ or real homology $H_k(X, \mathbb{R}) := H^p_{cpt}(X, \widetilde{\mathbb{R}}_X)$, where p + k = n.

Example A.7 (Singular chains). Via integration of forms, each smooth singular k-chain determines a k-dimensional \mathcal{O}_X -twisted current. Let $\overline{\mathcal{C}}_k \subset \mathcal{D}'_k$ denote the subsheaf of the sheaf of germs of twisted currents corresponding to integration over singular chains. Then

$$0 \to \widetilde{\mathbb{Z}}_X \to \overline{\mathcal{C}}_n \to \overline{\mathcal{C}}_{n-1} \to \ldots \to \overline{\mathcal{C}}_0 \to 0$$

is an acyclic resolution of $\widetilde{\mathbb{Z}}_X$ and hence can be used to compute $H^p(X, \widetilde{\mathbb{Z}}_X)$ and to compute $H^p_{cpt}(X, \widetilde{\mathbb{Z}}_X)$. It can be shown that this $H^p_{cpt}(X, \widetilde{\mathbb{Z}}_X)$ computed from $\overline{\mathcal{C}}^*$ agrees with the usual homology $H_k(X, \mathbb{Z})$ (k = n - p) of the complex of smooth singular chains. That is, homology is just (compactly supported) cohomology with the coefficient sheaf $\widetilde{\mathbb{Z}}_X$, the twisted integers.

Example A.8 (Integrally flat currents). This is one of the examples of central importance in this paper and so will be treated in more detail. Let $\mathcal{F}_{loc}^{p}(X)$ denote the space of locally integrally flat degree *p* currents on *X*. We take as definition $\mathcal{F}_{loc}^{p}(X) := \mathcal{R}_{loc}^{p}(X) + d\mathcal{R}_{loc}^{p-1}(X)$, i.e., all currents that can be written as A + dB with $A \in \mathcal{R}_{loc}^{p}(X)$ and $B \in \mathcal{R}_{loc}^{p-1}(X)$ where $\mathcal{R}_{loc}^{p}(X)$ denotes the space of locally rectifiable degree *p* currents. In results where the degree of a locally integrally flat current can be arbitrary, we use the less encumbered notation $\mathcal{F}_{loc}(U)$.

The spaces $\{\mathcal{F}_{loc}(U) : U^{open} \subset X\}$ form a presheaf of abelian groups. One can form the associated sheaf \mathcal{F}_{loc} of germs, and consider the natural map from $\mathcal{F}_{loc}(U)$ to $\Gamma(U, \mathcal{F}_{loc})$. This map is injective because the support axiom is satisfied.

Support axiom

Let \mathcal{F} be a presheaf. If $T \in \mathcal{F}(X)$ restricts to be zero in a neighborhood of each point of X, then T = 0.

Note. For a general presheaf \mathcal{F} this axiom is equivalent to the concept of support being well defined. The support of $T \in \mathcal{F}(X)$ is defined to be the complement of the set of points $x \in X$ such that $T \mid_{U} = 0$ for some neighborhood U of x.

This map is surjective because the following axiom is satisfied.

Local to global axiom

Suppose $T_{\alpha} \in \mathcal{F}(U_{\alpha})$ is given, where $\{U_{\alpha}\}$ is a locally finite open cover of X. Let $U_{\alpha\beta} = U_{\alpha} \cap U_{\beta}$. If $T_{\alpha\beta} = T_{\alpha} |_{U_{\alpha\beta}} - T_{\beta} |_{U_{\alpha\beta}}$ vanishes, then there exists a global $T \in \mathcal{F}(X)$ such that $T |_{U_{\alpha}} = T_{\alpha}$.

For a given presheaf, if both of these conditions/axioms are satisfied, then the presheaf is said to be a **sheaf**. A sheaf is said to be **soft** if for each closed set $C \subset X$ and each section of the sheaf on C

there exists an extension to all of X. That is, for each section on a neighborhood of C there exists a section on X which agrees with the given section on a (smaller) neighborhood of C. Soft sheaves are always acyclic, for basically the same reason fine sheaves are acyclic. Namely, the decompositions provided by a partition of unity exist (even though these may not arise from a partition of unity).

Theorem A.9. The presheaf $\{\mathcal{F}_{loc}(U)\}$ of locally integrally flat currents is a sheaf and this sheaf \mathcal{F}_{loc} is soft.

Proof. Since each $\mathcal{F}_{loc}(U)$ is a subset of $\mathcal{D}'(U)$ and the support axiom is satisfied for the presheaf $\mathcal{D}'(U)$, the support axiom is automatic for $\mathcal{F}_{loc}(U)$.

To prove the local to global property we first describe the proof for $\mathcal{R}_{loc}(U)$. Given $A_{\alpha} \in \mathcal{R}_{loc}(U_{\alpha})$ with $A_{\alpha} = A_{\beta}$ on $U_{\alpha\beta}$ choose a partition of unity $\{\chi_{\alpha}\}$ for $\{U_{\alpha}\}$ with each χ_{α} a characteristic function of a Borel set. Then, since $\chi_{\alpha}A_{\alpha}$ is also a rectifiable current on U_{α} (but vanishing near ∂U_{α}), we may consider $\chi_{\alpha}A_{\alpha} \in \mathcal{R}_{loc}(X)$ extended by zero to all of X. Set $A = \sum \chi_{\beta}A_{\beta}$ and note that $A \mid_{U_{\alpha}} = A_{\alpha}$.

The proof of the local to global property for $\mathcal{F}_{loc}(U)$ can be outlined as follows. For a more complete proof, see [9], Lemma 3.1. Suppose that we are given $T_{\alpha} = A_{\alpha} + dB_{\alpha}$ with $A_{\alpha}, B_{\alpha} \in \mathcal{R}_{loc}(U_{\alpha})$ and $T_{\alpha} = T_{\beta}$ on $U_{\alpha\beta}$, i.e., $A_{\alpha} - A_{\beta} = d(B_{\beta} - B_{\alpha})$. Suppose we could set $T = \sum \chi_{\alpha} T_{\alpha}$, and verify that

$$\chi_{\alpha}T_{\alpha} = \chi_{\alpha}A_{\alpha} + \chi_{\alpha}dB_{\alpha} = \chi_{\alpha}A_{\alpha} - (d\chi_{\alpha})B_{\alpha} + d(\chi_{\alpha}B_{\alpha}).$$
(A.10)

More precisely we must show that $\chi_{\alpha}A_{\alpha} - d\chi_{\alpha}B_{\alpha}$ and $\chi_{\alpha}B_{\alpha}$ define locally rectifiable currents satisfying Equation (A.10). This is not always true because $(d\chi_{\alpha})B_{\alpha}$ and/or $\chi_{\alpha}dB_{\alpha}$ may not be defined. However, by Federer's theory of slicing, we may choose a slight perturbation of χ_{α} so that $d\chi_{\alpha}B_{\alpha}$ is a well-defined rectifiable current and so that $\chi_{\alpha}dB_{\alpha}$ is a well-defined current with the equation $d(\chi_{\alpha}B_{\alpha}) = (d\chi_{\alpha})B_{\alpha} + \chi_{\alpha}dB_{\alpha}$ satisfied.

The proof that \mathcal{F}_{loc} is soft is easier. First consider the analogous result for locally rectifiable currents. \mathcal{R}_{loc} is soft because given a closed set $C \subset X$ and $A \in \mathcal{R}_{loc}(U)$ where U is a neighborhood of C we may choose χ to be the characteristic function of $\overline{V} \subset U$ where V is an open neighborhood of C and then set $\widehat{A} = \chi A \in \mathcal{R}_{loc}(X)$ to be the desired extension. Similarly, given $T = A + dB \in \mathcal{F}_{loc}(U)$ the current $\widehat{T} = \chi A + d(\chi B)$ provides the required extension.

Since \mathcal{F}_{loc} is soft we have the following:

Corollary A.11. The sheaf \mathcal{F}_{loc} is acyclic. In particular, for each locally finite open cover $\mathcal{U} = \{U_{\alpha}\}$ of X, the cohomology $H^{1}(\mathcal{U}, \mathcal{F}_{loc}) = 0$. That is, given $S_{\alpha\beta} \in \mathcal{F}_{loc}^{p}(U_{\alpha\beta})$ satisfying

$$S_{\alpha\beta} + S_{\beta\gamma} + S_{\gamma\alpha} = 0$$
 on $U_{\alpha\beta\gamma}$

there exists $S_{\alpha} \in \mathcal{F}_{loc}^{p}(U_{\alpha})$ such that

$$S_{\alpha\beta} = S_{\alpha} - S_{\beta}$$
 on $U_{\alpha\beta}$

Theorem A.12.

$$0 \to \mathbb{Z} \to \mathcal{F}^0_{\text{loc}} \to \mathcal{F}^1_{\text{loc}} \to \ldots \to \mathcal{F}^n_{\text{loc}} \to 0$$

is exact.

Proof. Exactness on the left is equivalent to $0 \to \mathbb{Z} \to \mathcal{R}^0_{loc} \to \mathcal{R}^1_{loc}$ being exact since $\mathcal{F}^0_{loc} = \mathcal{R}^0_{loc}$. This follows immediately from the standard fact that $0 \to \mathbb{R} \to \mathcal{D}'^0 \to \mathcal{D}'^1$ is exact, i.e.,

locally a *d*-closed generalized function is represented by a constant function. Let *U* be an open ball in \mathbb{R}^n . Suppose T = A + dB where $A \in \mathcal{R}^p_{loc}(U)$, $B \in \mathcal{R}^{p-1}_{loc}(U)$, and $p \ge 1$. If *T* is *d*-closed, then dA = 0. Using the standard cone construction (and Federer's theory) there exists $\widehat{A} \in \mathcal{R}^{p-1}_{loc}(U)$ such that $d\widehat{A} = A$ on *U*. Therefore, $d(\widehat{A} + B) = T$ where $\widehat{A} + B \in \mathcal{R}^{p-1}_{loc}(U) \subset \mathcal{F}^{p-1}_{loc}(U)$.

Corollary A.13.

$$H^{p}(X,\mathbb{Z}) = \frac{\left\{T \in \mathcal{F}_{\text{loc}}^{p}(X) : dT = 0\right\}}{d\mathcal{F}_{\text{loc}}^{p-1}(X)}$$

That is, the integrally flat currents can be used to compute \mathbb{Z} cohomology.

Lemma A.14. Given $T \in \mathcal{F}_{loc}(X)$ and a neighborhood U of spt T there exist A, $B \in \mathcal{R}_{loc}(X)$ satisfying T = A + dB and with spt $A \subset U$, spt $B \subset U$.

Remark. In particular, Lemma A.14 implies that if $T \in \mathcal{F}_{cpt}(X)$, then there exist $A, B \in \mathcal{R}_{cpt}(X)$ with T = A + dB. Consequently, Federer's definition of $\mathcal{F}_{cpt}(X)$ (where $\mathcal{F}_{cpt}(X) = \mathcal{R}_{cpt}(X) + d\mathcal{R}_{cpt}(X)$) agrees with the definition given in this appendix (where $\mathcal{F}_{cpt}(X)$ is the space of compactly supported sections of \mathcal{F}_{loc}).

Federer defines $\mathcal{F}_{loc}(X)$ to be the space of those currents on X which locally agree with a current of the type A + dB where $A, B \in \mathcal{R}_{cpt}(X)$. As a consequence of the discussion above, one can easily show that the definition of $\mathcal{F}_{loc}(X)$ given in this appendix (namely $\mathcal{R}_{loc}(X) + d\mathcal{R}_{loc}(X)$) agrees with the definition in [5].

Proof of Lemma. Choose $A, B \in \mathcal{R}_{loc}(X)$ with T = A + dB. On $X \sim \operatorname{spt} T$, both B and dB = -A are locally rectifiable. By Federer's theory of slicing, there exists a neighborhood V of spt T with $\overline{V} \subset U$ such that on $X \sim \operatorname{spt} T$, the slice $(d\chi)B \in \mathcal{R}_{loc}(X \sim \operatorname{spt} T)$ exists and $d(\chi B) = (d\chi)B + \chi dB$ on $X \sim \operatorname{spt} T$. Here χ denotes the characteristic function of \overline{V} . Consequently,

$$T = A + dB = \chi A + \chi dB = \chi A - (d\chi)B + d(\chi B),$$

where $\chi A - (d\chi)B$ and $d(\chi B)$ are locally rectifiable on X with support in U.

Example A.15 (Integrally flat \mathcal{O}_X -twisted currents). This example is almost identical to Example A.8. Let $\mathcal{R}_k^{\text{loc}}(X)$ denote the space of locally rectifiable \mathcal{O}_X -twisted currents of dimension k, and let $\mathcal{F}_k^{\text{loc}} := \mathcal{R}_k^{\text{loc}}(X) + d\mathcal{R}_k^{\text{loc}}(X)$ denote the space of locally integrally flat \mathcal{O}_X -twisted currents of dimension k. Then

$$0 \to \widetilde{\mathbb{Z}}_X \to \mathcal{F}_n^{\mathrm{loc}} \to \mathcal{F}_{n-1}^{\mathrm{loc}} \to \ldots \to \mathcal{F}_0^{\mathrm{loc}} \to 0$$

is an acyclic resolution of $\widetilde{\mathbb{Z}}_X$. Hence, the complex $\{\mathcal{F}_k^{\text{loc}}, d\}$ can be used to compute $H^p(X, \widetilde{\mathbb{Z}}_X)$, or integer homology $H_k(X, \mathbb{Z}) = H^p_{\text{cpt}}(X, \widetilde{\mathbb{Z}}_X)$, where p + k = n.

Example A.16 (Integrally flat \mathcal{O}_V -**twisted currents).** Let $V \to X$ be a real bundle with orientation sheaf \mathcal{O}_V and let $\widetilde{\mathbb{Z}}_V := \mathbb{Z} \otimes_{\mathbb{Z}_2} \mathcal{O}_V$. Let $\widetilde{\mathcal{F}}_{loc}^p(X)$ denote the space of degree p locally integrally flat \mathcal{O}_V -twisted currents on X. Then

$$0 \to \widetilde{\mathbb{Z}}_V \to \widetilde{\mathcal{F}}^0_{\text{loc}} \to \widetilde{\mathcal{F}}^1_{\text{loc}} \to \ldots \to \widetilde{\mathcal{F}}^n_{\text{loc}} \to 0$$

is an acyclic resolution of $\widetilde{\mathbb{Z}}_V$. Hence, the complex $\{\widetilde{\mathcal{F}}_{loc}^p, d\}$ can be used to compute $H^p(X, \widetilde{\mathbb{Z}}_V)$.

Example A.17 (Mod q integrally flat currents). Cohomology with \mathbb{Z}_q coefficients can be computed using mod q currents.

Definition A.18. The space $\mathcal{F}_{loc}^{mod q}(X)$ of mod q locally integrally flat currents on X is defined to be the quotient $\mathcal{F}_{loc}^{p}(X) / q \mathcal{F}_{loc}^{p}(X)$.

To avoid an excess of subscripts and superscripts we always drop the superscript p for the degree and the subscript k for the dimension when describing mod q currents.

Remark. Federer takes the quotient of $\mathcal{F}_{loc}^{p}(X)$ by the closure of $q\mathcal{F}_{loc}^{p}(X)$ in the flat topology on $\mathcal{F}_{loc}^{p}(X)$ in order to prove compactness theorems for mod q currents. For our purposes — computing cohomology — this closure is an unnecessary complication. In addition, Fred Almgren (private communication) has proven that $q\mathcal{F}_{loc}^{p}(X)$ is already closed in $\mathcal{F}_{loc}^{p}(X)$.

Theorem A.19. The presheaf $\{\mathcal{F}_{loc}^{mod q}(U)\}$ is a sheaf and this sheaf $\mathcal{F}_{loc}^{mod q}$ is soft.

Proof. First we must verify the support axiom. Suppose $\overline{T} \in \mathcal{F}_{loc}^{mod q}(X)$ is given (and represented by $T \in \mathcal{F}_{loc}(X)$). Further suppose $\overline{T} \mid_{U_{\alpha}} = 0$ for each α , i.e., there exists $S_{\alpha} \in \mathcal{F}_{loc}(U_{\alpha})$ such that $T \mid_{U_{\alpha}} = qS_{\alpha}$ for each α . Now $qS_{\alpha} = qS_{\beta}$ on $U_{\alpha\beta}$ implies $S_{\alpha} = S_{\beta}$ on $U_{\alpha\beta}$. Since $\{\mathcal{F}_{loc}(U)\}$ satisfies the local to global axiom, there exists $S \in \mathcal{F}_{loc}(X)$ such that $S \mid_{U_{\alpha}} = S_{\alpha}$. Consequently T = qS on X so that $\overline{T} = 0$.

Second we must verify the local to global axiom for $\{\mathcal{F}_{loc}^{mod\,q}(U)\}$. Suppose $\overline{T}_{\alpha} \in \mathcal{F}_{loc}^{mod\,q}(\mathcal{U}_{\alpha})$ are given with $\overline{T}_{\alpha} = \overline{T}_{\beta}$ on $U_{\alpha\beta}$. Choosing representatives $T_{\alpha} \in \mathcal{F}_{loc}(U_{\alpha})$ for \overline{T}_{α} , the equation $\overline{T}_{\alpha} = \overline{T}_{\beta}$ on $U_{\alpha\beta}$ says that there exists $S_{\alpha\beta} \in \mathcal{F}_{loc}(U_{\alpha\beta})$ such that $T_{\alpha} - T_{\beta} = qS_{\alpha\beta}$ on $U_{\alpha\beta}$. The cocycle condition for $S_{\alpha\beta}$ is satisfied since it is satisfied for $qS_{\alpha\beta}$. Recall that $H^1(\mathcal{U}, \mathcal{F}_{loc}) = 0$. This implies that there exists $S_{\alpha} \in \mathcal{F}_{loc}(\mathcal{U}_{\alpha})$ such that $S_{\alpha} - S_{\beta} = S_{\alpha\beta}$. Consequently,

$$T = T_{\alpha} - q S_{\alpha} \in \mathcal{F}_{\text{loc}}(X)$$

is well defined independent of α and provides the global representation of a class $\overline{T} \in \mathcal{F}_{loc}^{mod q}(X)$ with $\overline{T}|_{U_{\alpha}} = \overline{T}_{\alpha}$ for each α .

Finally, we must show that the sheaf $\mathcal{F}_{loc}^{mod q}$ is soft. Suppose $\overline{T} \in \mathcal{F}_{loc}^{mod q}(U)$ is given (represented by $T \in \mathcal{F}_{loc}(U)$) and U is an open neighborhood of a closed set C. Since \mathcal{F}_{loc} is soft, there exists $S \in \mathcal{F}_{loc}(X)$ such that $S \mid_{V} = T \mid_{V}$ for some neighborhood $V \subset U$ of C. Now $\overline{S} \in \mathcal{F}_{loc}^{mod q}(X)$ and $\overline{S} = \overline{T}$ on the neighborhood V of C.

Proposition A.20. Given $\overline{T} \in \mathcal{F}_{loc}^{mod q}(X)$ and a neighborhood U of $A := \operatorname{spt} \overline{T}$ there exists a representative $T \in \mathcal{F}_{loc}(X)$ of \overline{T} with $\operatorname{spt} T \subset U$.

Proof. Let $T \in \mathcal{F}_{loc}(X)$ denote an arbitrary representative of \overline{T} . Restricted to $X \sim A$, $T \mid_{X \sim A} = q\widehat{S}$ for some $\widehat{S} \in \mathcal{F}_{loc}(X \sim A)$. Since \mathcal{F}_{loc} is soft there exists a global section $S \in \mathcal{F}_{loc}(X)$ which agrees with \widehat{S} on a neighborhood of the closed set $X \sim U$. Now $T - qS \in \mathcal{F}_{loc}(X)$ is another representative of \overline{T} and T - qS vanishes on a neighborhood of $X \sim U$.

Note that the sheaf \mathbb{Z}_q is a subsheaf of $\mathcal{F}_{loc}^{mod q}$ when the degree is zero (i.e., when dimension is *n*).

Lemma A.21. Suppose $\overline{T} \in \mathcal{F}_{loc}^{mod q}(U)$ is of degree zero on the open unit ball U. If $d\overline{T} = 0$, then \overline{T} is represented by a constant integer valued function.

Proof. The equation $d\overline{T} = 0$ means that dT = qS with $S \in \mathcal{F}_{loc}^{1}(U)$ and $T \in \mathcal{F}_{loc}^{0}(U)$ a representative for \overline{T} . The current qS and hence S is d-closed. Therefore, there exists $R \in \mathcal{F}_{loc}^{0}(U)$ satisfying dR = S (because this is true for locally rectifiable currents). Now $T - qR \in \mathcal{F}_{loc}^{0}(U)$ also represents \overline{T} and is d-closed. Therefore, T - qR is a constant integer valued function.

This proves that $0 \to \mathbb{Z}_q \to \mathcal{F}^{\mathrm{mod}\,q}_{\mathrm{loc,deg}\,0} \xrightarrow{d} \mathcal{F}^{\mathrm{mod}\,q}_{\mathrm{loc,deg}\,1}$ is exact.

Lemma A.22. Let $p \ge 1$. Given $\overline{T} \in \mathcal{F}_{\text{loc,deg }p}^{\text{mod }q}(U)$ on the unit ball U satisfying $d\overline{T} = 0$ there exists $\overline{S} \in \mathcal{F}_{\text{loc,deg }p-1}^{\text{mod }q}(U)$ with $d\overline{S} = \overline{T}$.

Proof. There exist $T \in \mathcal{F}_{loc}^{p}(U)$ and $S \in \mathcal{F}_{loc}^{p+1}(U)$ such that dT = qS. Thus, dS = 0. Now S = A + dB where $A, B \in \mathcal{R}_{loc}(U)$. Thus, dA = 0. Since degree $A = p + 1 \ge 1$ there exists $R \in \mathcal{R}_{loc}^{p}(U)$ such that A = dR. Therefore, S = dB for some $B \in \mathcal{R}_{loc}^{p}(U)$. Consequently $\widehat{T} = T - qB \in \mathcal{F}_{loc}^{p}(U)$ is another representative of \overline{T} with $d\widehat{T} = 0$. Finally solve $dS = \widehat{T}$ on U with $S \in \mathcal{F}_{loc}^{p}(U)$.

In summary, the mod q currents $\mathcal{F}_{loc}^{mod q}(X)$ may be used to compute cohomology with \mathbb{Z}_q coefficients.

Theorem A.23.

$$H^{p}\left(X, \mathbb{Z}_{q}\right) = \frac{\left\{\overline{T} \in \mathcal{F}_{\mathrm{loc}, \deg p}^{\mathrm{mod} q}(X) : d\overline{T} = 0\right\}}{d \mathcal{F}_{\mathrm{loc}, \deg p-1}^{\mathrm{mod} q}(X)}$$

Example A.24 (Mod q integrally flat \mathcal{O}_X -twisted currents). This example is almost identical to Example A.17. Let $\mathcal{F}_{\text{mod }q}^{\text{loc}}(X) := \mathcal{F}_k^{\text{loc}}(X) / q \mathcal{F}_k^{\text{loc}}(X)$ be the space of mod q locally integrally flat \mathcal{O}_X -twisted currents on X and let $\widetilde{\mathbb{Z}}_q := \mathcal{O}_X \otimes_{\mathbb{Z}_2} \mathbb{Z}_q$. Then the complex $\{\mathcal{F}_{\text{mod }q}^{\text{loc}}(X), d\}$ can be used to compute $H^p(X, \widetilde{\mathbb{Z}}_q)$, or $H_k(X, \mathbb{Z}_q) = H_{\text{cpl}}^p(X, \widetilde{\mathbb{Z}}_q)$ where p + k = n.

Example A.25 (The mod 2 integrally flat case). In this example we wish to compute \mathbb{Z}_2 cohomology $H^p(X, \mathbb{Z}_2)$. This can be done in several equivalent ways. First, restating Theorem A.23 in the case q = 2, $H^p(X, \mathbb{Z}_2)$ can be computed from the complex $\mathcal{F}_{loc}^{mod 2}(X) := \mathcal{F}_{loc}^*(X) / 2\mathcal{F}_{loc}^*(X)$ of mod 2 locally integrally flat currents on X.

Next suppose that V is a real bundle with orientation sheaf \mathcal{O}_V and let $\widetilde{\mathcal{F}}_{loc}^p(X)$ denote the degree p locally integrally flat \mathcal{O}_V -twisted currents on X.

Lemma A.26.

 $\mathcal{F}^{p}_{\text{loc}}(X) \ \big/ \ 2\mathcal{F}^{p}_{\text{loc}}(X) \ \cong \ \widetilde{\mathcal{F}}^{p}_{\text{loc}}(X) \ \big/ \ 2\widetilde{\mathcal{F}}^{p}_{\text{loc}}(X)$

Therefore, $H^p(X, \mathbb{Z}_2)$ can also be computed using the complex of mod 2 locally integrally flat \mathcal{O}_V -twisted currents.

Proof of Lemma A.26. First we define a mapping $\varphi : \widetilde{\mathcal{F}}_{loc}^{p}(X) \to \mathcal{F}_{loc}^{p}(X) / 2\mathcal{F}_{loc}^{p}(X)$. Let $T \in \widetilde{\mathcal{F}}_{loc}^{p}(X)$. Choose a locally finite open cover $\{U_{\alpha}\}$ of X and fix ordered frames e_{α} for V over U_{α} .

For each pair (U_{α}, e_{α}) let $T_{\alpha} \in \mathcal{F}_{loc}^{p}(U_{\alpha})$ be defined by $T = [e_{\alpha}] \otimes T_{\alpha}$ on U_{α} . Note that $T_{\alpha} = \pm T_{\beta}$ on $U_{a} \cap U_{b}$. Define $S_{\alpha\beta} \in \mathcal{F}_{loc}(U_{\alpha} \cap U_{\beta})$ by

$$S_{\alpha\beta} = \begin{cases} 0 & \text{if } T_{\alpha} = T_{\beta} \text{ on } U_{\alpha} \cap U_{\beta}, \\ T_{\alpha} & \text{if } T_{\alpha} = -T_{\beta} \text{ on } U_{\alpha} \cap U_{\beta}. \end{cases}$$

Then

$$T_{\alpha} - T_{\beta} = 2S_{\alpha\beta} \quad \text{on } U_{\alpha} \cap U_{\beta} .$$
 (A.27)

Let $\overline{T_{\alpha}} \in \mathcal{F}_{\text{loc}}^{\text{mod 2}}(U_{\alpha})$ be the mod 2 class of T_{α} . Then, by Equation (A.27), $\overline{T_{\alpha}} = \overline{T_{\beta}}$ on $U_{\alpha} \cap U_{\beta}$. Since the presheaf $\{\mathcal{F}_{\text{loc}}^{\text{mod 2}}(U)\}$ satisfies the Local to Global Axiom (see Theorem A.19), there is a well-defined element $\varphi(T) \in \mathcal{F}_{\text{loc}}^{\text{mod 2}}(X)$ so that $\varphi(T) \mid_{U_{\alpha}} = \overline{T_{\alpha}}$. Note that $\varphi(T)$ is well defined independent of the choices of locally finite open cover $\{U_{\alpha}\}$ and frames e_{α} for V over U_{α} .

In summary, we have defined a map $\varphi : \widetilde{\mathcal{F}}_{loc}^{p}(X) \to \mathcal{F}_{loc}^{p}(X) / 2\mathcal{F}_{loc}^{p}(X)$. Now, since $\varphi(2T) \mid_{U_{\alpha}} = \overline{2T_{\alpha}} = 0$, the induced map $\widehat{\varphi} : \widetilde{\mathcal{F}}_{loc}^{p}(X) / 2\widetilde{\mathcal{F}}_{loc}^{p}(X) \to \mathcal{F}_{loc}^{p}(X) / 2\mathcal{F}_{loc}^{p}(X)$ is well defined. Finally, to show that $\widehat{\varphi}$ is an isomorphism, we can use the same procedure to construct an inverse for $\widehat{\varphi}$.

References

- [1] Banchoff, T. and McCrory, C. Whitney duality and singularities of projections, L.N.M., 597, 68-82, (1976).
- [2] Borel, A. Topics in the homology theory of fibre bundles, L.N.M., 36, 1-95, (1967).
- [3] Bott, R. and Chern, S.S. Hermitian vector bundles and the equidistribution of the zeroes of their holomorphic sections, *Acta Math.*, **114**, 71–112, (1965).
- [4] Eells, J. A generalization of the Gauss-Bonnet Theorem, Trans. A.M.S., 92, 142–153, (1959).
- [5] Federer, H. Geometric Measure Theory. Springer-Verlag, New York, 1969.
- [6] Fu, J. and McCrory, C. Stiefel–Whitney classes and the conormal cycle of a singular variety, Trans. A.M.S., 349 (2), 809–835, (1997).
- [7] Fulton, W. Intersection Theory. Ergobnisse der Mathematik und ihrer Grenzgebeite 3, Folge, Band 2, Springer-Verlag, Berlin-Heidelberg, 1984.
- [8] Godement, R. Topologie Algébrique et Théorie des Faisceaux. Hermann, Paris, 1964.
- [9] Hardt, R.M. Slicing and intersection theory for chains associated with real analytic varieties, Acta Math., 129, 75–136, (1972).
- [10] Harvey, F.R. and Lawson, H.B. A theory of characteristic currents associated with a singular connection, *Astérisque*, 213, 1–268, (1993).
- [11] Harvey, F.R. and Lawson, H.B. A theory of characteristic currents associated with a singular connection Announcement, B.A.M.S., 31, 54–63, (1994).
- [12] Harvey, F.R. and Lawson, H.B. Geometric residue theorems, Am. J. Math., 117, 829-873, (1995).
- [13] Harvey, F.R. and Lawson, H.B. Morse Theory and Stokes's Theorem, (to appear), 1-32.
- [14] Harvey, F.R. and Semmes, S. Zero divisors of atomic functions, Ann. Math., 135, 567-600, (1992).
- [15] Halperin, S. and Toledo, D. Stiefel-Whitney homology classes, Ann. Math. (S2), 96, 511-525, (1972).
- [16] Milnor, J.W. and Stasheff, J.D. Characteristic Classes, Ann. Math. Studies, 76, Princeton University Press, Princeton, NJ.
- [17] Pontrjagin, L.S. Vector fields on manifolds, Mat. Sbornik N.S., 24, 129-162, (1949).
- [18] Pontrjagin, L.S. Characteristic cycles on differentiable manifolds, Mat. Sbornik N.S., 21, 233–284, (1947); A.M.S. Translation, 32, (1950).
- [19] Porteous, I.R. Simple singularities of maps Proceedings of Liverpool Singularities Symposium, I, Lecture Notes in Mathematics, 192, 286–307, (1971).
- [20] Ronga, F. Le calcul de la classe de cohomologie duale a $\overline{\Sigma}^k$ Proceedings of Liverpool Singularities Symposium, I, Lecture Notes in Mathematics, **192**, 313–315, (1971).
- [21] Steenrod, N.E. The Topology of Fibre Bundles. Princeton University Press, Princeton, NJ, 1951.

Reese Harvey and John Zweck

- [22] Stiefel, E. Richtungsfelder und Fernparallelismus in n-dimensionalen Mannigfaltigkeiten, Comment. Math. Helv., 8, 305–353, (1934-1936).
- [23] Thom, R. Les singularitiés des applications différentiables, Ann. Inst. Fourier, Grenoble, 6, 43-87, (1955-1956).
- [24] Whitney, H. Sphere-spaces, Proc. Natl. Acad. Sci., 21, 64-68, (1935).
- [25] Zweck, J. Compactification problems in the theory of characteristic currents associated with a singular connection, Thesis, Rice University, Houston, TX, 1993.
- [26] Zweck, J. Euler and Pontrjagin currents of a section of a compactified real bundle, *Diff. Geom. Applns.*, **5**, 277–309, (1995).

Received Novermber 28, 1995 Revision received September 26, 1996

Rice University, Houston, Texas

University of Nevada, Reno, Nevada