

Einstein Constants

and

Differential Topology

Claude LeBrun

Stony Brook University

MPI-Oberseminar,
Max-Planck-Institut für Mathematik,
Bonn, 5. Juni, 2025.

Let (M^n, g) be a Riemannian n -manifold,

Let (M^n, g) be a Riemannian n -manifold, $p \in M$.

Let (M^n, g) be a Riemannian n -manifold, $p \in M$.
Metric defines locally shortest curves, called *geodesics*.

Let (M^n, g) be a Riemannian n -manifold, $p \in M$.
Metric defines locally shortest curves, called *geodesics*.
Following geodesics from p defines a map

$$\exp : T_p M \dashrightarrow M$$

Let (M^n, g) be a Riemannian n -manifold, $p \in M$.
Metric defines locally shortest curves, called *geodesics*.
Following geodesics from p defines a map


$$\exp : T_p M \longrightarrow M$$

which is a diffeomorphism on a neighborhood of 0:

Let (M^n, g) be a Riemannian n -manifold, $p \in M$.
Metric defines locally shortest curves, called *geodesics*.
Following geodesics from p defines a map

$$\exp : T_p M \longrightarrow M$$


which is a diffeomorphism on a neighborhood of 0:



Let (M^n, g) be a Riemannian n -manifold, $p \in M$.
 Metric defines locally shortest curves, called *geodesics*.
 Following geodesics from p defines a map

$$\exp : T_p M \longrightarrow M$$

which is a diffeomorphism on a neighborhood of 0:



Now choosing $T_p M \xrightarrow{\cong} \mathbb{R}^n$ via some orthonormal basis gives us special coordinates on M .

In these “geodesic normal” coordinates the metric volume measure is given by

$$d\mu_g = d\mu_{\text{Euclidean}},$$

In these “geodesic normal” coordinates the metric volume measure is given by

$$d\mu_g = \left[1 - \frac{1}{6} R_{ij} x^i x^j + \frac{1}{120} R_{ijkl} x^i x^j x^k x^l - \frac{1}{420} R_{ijklm} x^i x^j x^k x^l x^m + \dots \right] d\mu_{\text{Euclidean}},$$

In these “geodesic normal” coordinates the metric volume measure is given by

$$d\mu_g = \left[1 - \frac{1}{6} r_{jk} x^j x^k + \right] d\mu_{\text{Euclidean}},$$

In these “geodesic normal” coordinates the metric volume measure is given by

$$d\mu_g = \left[1 - \frac{1}{6} r_{jk} x^j x^k + O(|x|^3) \right] d\mu_{\text{Euclidean}},$$

In these “geodesic normal” coordinates the metric volume measure is given by

$$d\mu_g = \left[1 - \frac{1}{6} r_{jk} x^j x^k + O(|x|^3) \right] d\mu_{\text{Euclidean}},$$

where r is the *Ricci tensor*

In these “geodesic normal” coordinates the metric volume measure is given by

$$d\mu_g = \left[1 - \frac{1}{6} r_{jk} x^j x^k + O(|x|^3) \right] d\mu_{\text{Euclidean}},$$

where r is the *Ricci tensor* $r_{jk} = \mathcal{R}^i_{jik}$.

In these “geodesic normal” coordinates the metric volume measure is given by

$$d\mu_g = \left[1 - \frac{1}{6} r_{jk} x^j x^k + O(|x|^3) \right] d\mu_{\text{Euclidean}},$$

where r is the *Ricci tensor* $r_{jk} = \mathcal{R}^i_{jik}$.

The *Ricci curvature*

In these “geodesic normal” coordinates the metric volume measure is given by

$$d\mu_g = \left[1 - \frac{1}{6} r_{jk} x^j x^k + O(|x|^3) \right] d\mu_{\text{Euclidean}},$$

where r is the *Ricci tensor* $r_{jk} = \mathcal{R}^i_{jik}$.

The *Ricci curvature* is by definition the function on the unit tangent bundle

In these “geodesic normal” coordinates the metric volume measure is given by

$$d\mu_g = \left[1 - \frac{1}{6} r_{jk} x^j x^k + O(|x|^3) \right] d\mu_{\text{Euclidean}},$$

where r is the *Ricci tensor* $r_{jk} = \mathcal{R}^i_{jik}$.

The *Ricci curvature* is by definition the function on the unit tangent bundle

$$STM = \{v \in TM \mid g(v, v) = 1\}$$

In these “geodesic normal” coordinates the metric volume measure is given by

$$d\mu_g = \left[1 - \frac{1}{6} r_{jk} x^j x^k + O(|x|^3) \right] d\mu_{\text{Euclidean}},$$

where r is the *Ricci tensor* $r_{jk} = \mathcal{R}^i_{jik}$.

The *Ricci curvature* is by definition the function on the unit tangent bundle

$$STM = \{v \in TM \mid g(v, v) = 1\}$$

given by

$$v \longmapsto r(v, v).$$

Definition. *A Riemannian metric g is said to be Einstein if it has constant Ricci curvature*

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.


“...the greatest blunder of my life!”

— A. Einstein, to G. Gamow

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.



Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

As punishment ...

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Mathematicians call λ the Einstein constant.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Mathematicians call λ the Einstein constant.

Has same sign as the *scalar curvature*

$$s = r_j^j = \mathcal{R}^{ij}_{ij}.$$

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$


for some constant $\lambda \in \mathbb{R}$.

Mathematicians call λ the Einstein constant.

Has same sign as the *scalar curvature*

$$s = r_j^j = \mathcal{R}^{ij}_{ij}.$$

$$\frac{\text{vol}_g(B_\varepsilon(p))}{c_n \varepsilon^n} = 1 - s \frac{\varepsilon^2}{6(n+2)} + O(\varepsilon^4)$$



Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Generalizes constant sectional curvature condition, but weaker.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Generalizes constant sectional curvature condition, but weaker — unless $\dim M = 2$ or 3 .

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Generalizes constant sectional curvature condition, but weaker.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Generalizes constant sectional curvature condition, but weaker.

Determined system:

same number of equations as unknowns.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Generalizes constant sectional curvature condition, but weaker.

Determined system:

same number of equations as unknowns.

g_{jk} : $\frac{n(n+1)}{2}$ components.

r_{jk} : $\frac{n(n+1)}{2}$ components.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Generalizes constant sectional curvature condition, but weaker.

Determined system:

same number of equations as unknowns.

g_{jk} : $\frac{n(n+1)}{2}$ components.

r_{jk} : $\frac{n(n+1)}{2}$ components.

$\mathcal{R}^j_{k\ell m}$: $\frac{n^2(n^2-1)}{12}$ components.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Generalizes constant sectional curvature condition, but weaker.

Determined system:

same number of equations as unknowns.

Elliptic non-linear PDE after gauge fixing.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Generalizes constant sectional curvature condition, but weaker.

Determined system:

same number of equations as unknowns.

Elliptic non-linear PDE after gauge fixing.

$$\Delta x^j = 0 \implies r_{jk} = \frac{1}{2} \Delta g_{jk} + \text{lots.}$$

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Definition. An Einstein manifold (M, g) is a smooth compact manifold M

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Definition. An Einstein manifold (M, g) is a smooth compact manifold M (without boundary),

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

Definition. An Einstein manifold (M, g) is a smooth compact manifold M (without boundary), equipped with a smooth Einstein metric g .

Definition. A Riemannian metric g is said to be Einstein if it has constant Ricci curvature — i.e.

$$r = \lambda g$$

for some constant $\lambda \in \mathbb{R}$.

For the purposes of this talk...

Definition. An Einstein manifold (M, g) is a smooth compact manifold M (without boundary), equipped with a smooth Einstein metric g .

Ergebnisse der Mathematik und ihrer Grenzgebiete

3. Folge · Band 10

A Series of Modern Surveys in Mathematics

Arthur L. Besse

Einstein Manifolds



Springer-Verlag Berlin Heidelberg GmbH



Marcel Berger



Besse en Chandesse



L'Auvergne

Acknowledgements

Pour rassembler les éléments un peu disparates qui constituent ce livre, j'ai dû faire appel à de nombreux amis, heureusement bien plus savants que moi. Ce sont, entre autres, Geneviève Averous, Lionel Bérard-Bergery, Marcel Berger, Jean-Pierre Bourguignon, Andrei Derdzinski, Dennis M. DeTurck, Paul Gauduchon, Nigel J. Hitchin, Josette Houillot, Hermann Karcher, Jerry L. Kazdan, Norihito Koiso, Jacques Lafontaine, Pierre Pansu, Albert Polombo, John A. Thorpe, Liane Valère.

Les institutions suivantes m'ont prêté leur concours matériel, et je les en remercie: l'UER de mathématiques de Paris 7, le Centre de Mathématiques de l'Ecole Polytechnique, Unités Associées du CNRS, l'UER de mathématiques de Chambéry et le Conseil Général de Savoie.

Enfin, qu'il me soit permis de saluer ici mon prédécesseur et homonyme Jean Besse, de Zürich, qui s'est illustré dans la théorie des fonctions d'une variable complexe (voir par exemple [Bse]).

Vôtre,

A handwritten signature in black ink, consisting of a stylized, overlapping loop structure with a horizontal line extending to the right.

Arthur Besse

Le Faux, le 15 septembre 1986

Ergebnisse der Mathematik und ihrer Grenzgebiete

3. Folge · Band 10

A Series of Modern Surveys in Mathematics

Arthur L. Besse

Einstein Manifolds



Springer-Verlag Berlin Heidelberg GmbH

The book's Introduction concludes as follows:

The book's Introduction concludes as follows:

“Can Einstein metrics with [Einstein] constants of opposite signs exist on the same manifold?”

The book's Introduction concludes as follows:

“Can Einstein metrics with [Einstein] constants of opposite signs exist on the same manifold?”

If this is impossible, it would add weight to the remark... that positive and negative Einstein metrics belong to essentially distinct families.”

The book's Introduction concludes as follows:

“Can Einstein metrics with [Einstein] constants of opposite signs exist on the same manifold?”

If this is impossible, it would add weight to the remark... that positive and negative Einstein metrics belong to essentially distinct families.”

— A.L. Besse, *Einstein Manifolds*, p. 19

“Can Einstein metrics with [Einstein] constants of opposite signs exist on the same manifold?”

— A.L. Besse, *Einstein Manifolds*, p. 19

“Can Einstein metrics with [Einstein] constants of opposite signs exist on the same manifold?”

— A.L. Besse, *Einstein Manifolds*, p. 19

Became a folk-conjecture that the answer is **No**.

“Can Einstein metrics with [Einstein] constants of opposite signs exist on the same manifold?”

— A.L. Besse, *Einstein Manifolds*, p. 19

Became a folk-conjecture that the answer is **No**.

But it turns out that the answer is actually **Yes!**

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Specific examples:

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Specific examples:

$$M = \underbrace{Y \times \cdots \times Y}_k$$

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Specific examples:

$$M = \underbrace{Y \times \cdots \times Y}_k$$

where

$$Y^4 = \mathbb{CP}_2 \# 8\overline{\mathbb{CP}}_2.$$


Conventions:

$\overline{\mathbb{CP}}_2$ = reverse oriented $\mathbb{CP}_2$.

Conventions:

$\overline{\mathbb{CP}}_2$ = reverse oriented $\mathbb{CP}_2$.


Connected sum #:



Conventions:

$\overline{\mathbb{CP}}_2$ = reverse oriented $\mathbb{CP}_2$.


Connected sum #:



Conventions:

$\overline{\mathbb{CP}}_2$ = reverse oriented $\mathbb{CP}_2$.


Connected sum #:



Conventions:

$\overline{\mathbb{CP}}_2$ = reverse oriented $\mathbb{CP}_2$.


Connected sum #:



Conventions:

$\overline{\mathbb{CP}}_2$ = reverse oriented $\mathbb{CP}_2$.


Connected sum #:



Conventions:

$\overline{\mathbb{CP}}_2$ = reverse oriented $\mathbb{CP}_2$.


Connected sum #:



Conventions:

$\overline{\mathbb{CP}}_2$ = reverse oriented $\mathbb{CP}_2$.

Connected sum #:



Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Specific examples:

$$M = \underbrace{Y \times \cdots \times Y}_k$$

where

$$Y^4 = \mathbb{CP}_2 \# 8 \overline{\mathbb{CP}_2}.$$

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Specific examples:

$$M = \underbrace{Y \times \cdots \times Y}_k$$

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Specific examples:

$$M = \underbrace{Y \times \cdots \times Y}_k$$

“Key triviality:”

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Specific examples:

$$M = \underbrace{Y \times \cdots \times Y}_k$$

“Key triviality:”

Cartesian product

$$(M_1, g_1) \times (M_2, g_2) = (M_1 \times M_2, g_1 \oplus g_2)$$

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Specific examples:

$$M = \underbrace{Y \times \cdots \times Y}_k$$

“Key triviality:”

Cartesian product

$$(M_1, g_1) \times (M_2, g_2) = (M_1 \times M_2, g_1 \oplus g_2)$$

of two Einstein manifolds

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Specific examples:

$$M = \underbrace{Y \times \cdots \times Y}_k$$

“Key triviality:”

Cartesian product

$$(M_1, g_1) \times (M_2, g_2) = (M_1 \times M_2, g_1 \oplus g_2)$$

of two Einstein manifolds is Einstein

$\iff$ they have the same Einstein constant λ .

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Ingredients:

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Ingredients:

- geography of complex surfaces;

Theorem (Catanesi-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Ingredients:

- geography of complex surfaces;
- complex deformation theory;

Theorem (Catanesi-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Ingredients:

- geography of complex surfaces;
- complex deformation theory;
- theory of Kähler-Einstein metrics;

Theorem (Catanesi-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*


Ingredients:

- geography of complex surfaces;
- complex deformation theory;
- theory of Kähler-Einstein metrics;
- h -cobordisms of 4-manifolds; and


Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Ingredients:

- geography of complex surfaces;
- complex deformation theory;
- theory of Kähler-Einstein metrics;
- h -cobordisms of 4-manifolds; and
- Smale's h -cobordism theorem.




Cobordism




h -Cobordism


if $X \hookrightarrow W$, $Y \hookrightarrow W$ both homotopy equivalences



Smale: Suppose that X^n is h -cobordant to Y^n . If $\pi_1 = 0$ and $n > 4$, then X is diffeomorphic to Y .




Wall: Suppose that X^4 homotopy equivalent to Y^4 .
 If $\pi_1 = 0$, then X is h -cobordant to Y .




Wall: Suppose that X^4 homotopy equivalent to Y^4 .
 If $\pi_1 = 0$, then X is h -cobordant to Y .

But Smale doesn't apply when $n = 4$!




However, we still do get ...



However, we still do get ...


Lemma. *If X^4 and Y^4 are homotopy equivalent and simply connected, then $X \times X$ is actually diffeomorphic to $Y \times Y$.*



Indeed, if W is h -cobordism $X \sim Y$, then


$$(X \times W) \cup_{X \times Y} (W \times Y)$$

is an h -cobordism $(X \times X) \sim (Y \times Y)$.




Lemma. *If X^4 and Y^4 are homotopy equivalent and simply connected, then*

$$\underbrace{X \times \cdots \times X}_k \approx_{\text{diff}} \underbrace{Y \times \cdots \times Y}_k$$



Lemma. *If X^4 and Y^4 are homotopy equivalent and simply connected, then*

$$\underbrace{X \times \cdots \times X}_k \approx_{\text{diff}} \underbrace{Y \times \cdots \times Y}_k \quad \forall k \geq 2.$$



Lemma. *If X^4 and Y^4 are simply connected, non-spin, with $\chi(X) = \chi(Y)$, $\tau(X) = \tau(Y)$, then*

$$\underbrace{X \times \cdots \times X}_k \approx_{\text{diff}} \underbrace{Y \times \cdots \times Y}_k \quad \forall k \geq 2.$$

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Barlow 1985: Found compact complex surface (X^4, J) homotopy equivalent to $\mathbb{CP}_2 \# 8 \overline{\mathbb{CP}_2}$ with $c_1 \leq 0$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Barlow 1985: Found compact complex surface (X^4, J) homotopy equivalent to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$ with $c_1 \leq 0$.

C-L 1997: Barlow surface can be deformed to (X^4, J') with $c_1 < 0$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Barlow 1985: Found compact complex surface (X^4, J) homotopy equivalent to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$ with $c_1 \leq 0$.

C-L 1997: Barlow surface can be deformed to (X^4, J') with $c_1 < 0$. Aubin '76/Yau '77 $\implies X^4$ admits Einstein metric g^- with $\lambda < 0$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Barlow 1985: Found compact complex surface (X^4, J) homotopy equivalent to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$ with $c_1 \leq 0$.


C-L 1997: Barlow surface can be deformed to (X^4, J') with $c_1 < 0$. Aubin '76/Yau '77 $\implies X^4$ admits Kähler-Einstein metric g^- with $\lambda < 0$.

$(M^n, g):$

holonomy


$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy




$(M^n, g):$

holonomy



$(M^n, g):$


holonomy $\subset \mathbf{O}(n)$



Kähler metrics:


(M^{2m}, g) :

holonomy




Kähler metrics:

(M^{2m}, g) Kähler $\iff$ holonomy $\subset \mathbf{U}(m)$



Kähler metrics:

(M^{2m}, g) Kähler $\iff$ holonomy $\subset \mathbf{U}(m)$



$$\mathbf{U}(m) := \mathbf{O}(2m) \cap \mathbf{GL}(m, \mathbb{C})$$

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Barlow 1985: Found compact complex surface (X^4, J) homotopy equivalent to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$ with $c_1 \leq 0$.

C-L 1997: Barlow surface can be deformed to (X^4, J') with $c_1 < 0$. Aubin '76/Yau '77 $\implies X^4$ admits Kähler-Einstein metric g^- with $\lambda < 0$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Barlow 1985: Found compact complex surface (X^4, J) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$ with $c_1 \leq 0$.

C-L 1997: Barlow surface can be deformed to (X^4, J') with $c_1 < 0$. Aubin '76/Yau '77 $\implies X^4$ admits Kähler-Einstein metric g^- with $\lambda < 0$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Barlow 1985: Found compact complex surface (X^4, J) homeomorphic to $\mathbb{CP}_2 \# 8 \overline{\mathbb{CP}_2}$ with $c_1 \leq 0$.

Freedman 1982.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Barlow 1985: Found compact complex surface (X^4, J) homeomorphic to $\mathbb{CP}_2 \# 8 \overline{\mathbb{CP}_2}$ with $c_1 \leq 0$.

Freedman 1982.

Seiberg-Witten theory 1995. NOT diffeomorphic.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

$\exists \lambda < 0$ Einstein (X^4, g^-) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

$\exists \lambda < 0$ Einstein (X^4, g^-) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

Tian-Yau 1987: There are complex surfaces (Y^4, J)
with $c_1 > 0$

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

$\exists \lambda < 0$ Einstein (X^4, g^-) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

Tian-Yau 1987: There are complex surfaces (Y^4, J)
with $c_1 > 0$

“Del Pezzo surfaces”

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

$\exists \lambda < 0$ Einstein (X^4, g^-) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

Tian-Yau 1987: There are complex surfaces (Y^4, J) with $c_1 > 0$

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

$\exists \lambda < 0$ Einstein (X^4, g^-) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

Tian-Yau 1987: There are complex surfaces (Y^4, J) with $c_1 > 0$ which are diffeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

$\exists \lambda < 0$ Einstein (X^4, g^-) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

Tian-Yau 1987: There are complex surfaces (Y^4, J) with $c_1 > 0$ which are diffeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$ and admit Kähler-Einstein metrics with $\lambda > 0$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

$\exists \lambda < 0$ Einstein (X^4, g^-) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

$\exists \lambda > 0$ Einstein (Y^4, g^+) diffeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

$\exists \lambda < 0$ Einstein (X^4, g^-) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

$\exists \lambda > 0$ Einstein (Y^4, g^+) diffeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

$(\underbrace{X \times \cdots \times X}_k, g^- \oplus \cdots \oplus g^-)$ Einstein with $\lambda < 0$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

$\exists \lambda < 0$ Einstein (X^4, g^-) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

$\exists \lambda > 0$ Einstein (Y^4, g^+) diffeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

$(\underbrace{X \times \cdots \times X}_k, g^- \oplus \cdots \oplus g^-)$ Einstein with $\lambda < 0$.

$(\underbrace{Y \times \cdots \times Y}_k, g^+ \oplus \cdots \oplus g^+)$ Einstein with $\lambda > 0$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

$\exists \lambda < 0$ Einstein (X^4, g^-) homeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

$\exists \lambda > 0$ Einstein (Y^4, g^+) diffeomorphic to $\mathbb{CP}_2 \# 8\overline{\mathbb{CP}_2}$.

$(\underbrace{X \times \cdots \times X}_k, g^- \oplus \cdots \oplus g^-)$ Einstein with $\lambda < 0$.

$(\underbrace{Y \times \cdots \times Y}_k, g^+ \oplus \cdots \oplus g^+)$ Einstein with $\lambda > 0$.

$\underbrace{X \times \cdots \times X}_k \approx_{\text{diff}} \underbrace{Y \times \cdots \times Y}_k$ by h -cobordism thm.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Theorem (Răşdeaconu-Şuvaina 2009). *For every $k \geq 2$, there are at least $\binom{k+3}{3}$ distinct smooth closed simply-connected $4k$ -manifolds that admit both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Theorem (Răşdeaconu-Şuvaina 2009). *For every $k \geq 2$, there are at least $\binom{k+3}{3}$ distinct smooth closed simply-connected $4k$ -manifolds that admit both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Similar story for complex surfaces homeomorphic to $\mathbb{CP}_2 \# k \overline{\mathbb{CP}_2}$, $k = 5, 6, 7, 8$.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Theorem (Răşdeaconu-Şuvaina 2009). *For every $k \geq 2$, there are at least $\binom{k+3}{3}$ distinct smooth closed simply-connected $4k$ -manifolds that admit both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Similar story for complex surfaces homeomorphic to $\mathbb{CP}_2 \# k \overline{\mathbb{CP}_2}$, $k = 5, 6, 7, 8$.

Builds on breakthrough results of Jongil Park, Yongnam Lee, Heesang Park, and Dongsoo Shin.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Theorem (Răşdeaconu-Şuvaina 2009). *For every $k \geq 2$, there are at least $\binom{k+3}{3}$ distinct smooth closed simply-connected $4k$ -manifolds that admit both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

Theorem (Răşdeaconu-Şuvaina 2009). *For every $k \geq 2$, there are at least $\binom{k+3}{3}$ distinct smooth closed simply-connected $4k$ -manifolds that admit both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

These Einstein metrics are all of special holonomy.

Theorem (Catanese-L 1997). *For each $k \geq 2$, there is a closed simply-connected $4k$ -manifold M that admits both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*


Theorem (Răşdeaconu-Şuvaina 2009). *For every $k \geq 2$, there are at least $\binom{k+3}{3}$ distinct smooth closed simply-connected $4k$ -manifolds that admit both $\lambda > 0$ and $\lambda < 0$ Einstein metrics.*

These Einstein metrics are all of special holonomy.

Kähler, but for wildly different complex structures.


$(M^n, g):$

holonomy $\subset \mathbf{O}(n)$



$(M^n, g):$

holonomy $\subset \mathbf{O}(n)$



What are the possible holonomy groups?

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold.*

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither*

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds*

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space,*

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

[illegible]

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

[illegible]

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$

$$\mathbf{SU}(m) := \mathbf{O}(2n) \cap \mathbf{SL}(m, \mathbb{C})$$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	$\mathbb{H}$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	$\mathbb{H}$

$$\mathbf{Sp}(k) := \mathbf{O}(4k) \cap \mathbf{GL}(k, \mathbb{H})$$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	$\mathbb{H}$
$4k$	$\mathbf{Sp}(k)\mathbf{Sp}(1)$	Quaternion-Kähler	$\mathbb{H}$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	$\mathbb{H}$
$4k$	$\mathbf{Sp}(k)\mathbf{Sp}(1)$	Quaternion-Kähler	$\mathbb{H}$

$$\mathbf{Sp}(k)\mathbf{Sp}(1) := [\mathbf{Sp}(k) \times \mathbf{Sp}(1)]/\mathbb{Z}_2$$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	$\mathbb{H}$
$4k$	$\mathbf{Sp}(k)\mathbf{Sp}(1)$	Quaternion-Kähler	$\mathbb{H}$
7	$\mathbf{G}_2$	Exceptional I	$\mathbb{O}$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	$\mathbb{H}$
$4k$	$\mathbf{Sp}(k)\mathbf{Sp}(1)$	Quaternion-Kähler	$\mathbb{H}$
7	$\mathbf{G}_2$	Exceptional I	$\mathbb{O}$

$$\mathbf{G}_2 := \text{Aut}(\mathbb{O}) \subset \mathbf{SO}(7)$$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	$\mathbb{H}$
$4k$	$\mathbf{Sp}(k)\mathbf{Sp}(1)$	Quaternion-Kähler	$\mathbb{H}$
7	$\mathbf{G}_2$	Exceptional I	$\mathbb{O}$
8	$\mathbf{Spin}(7)$	Exceptional II	$\mathbb{O}$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	$\mathbb{H}$
$4k$	$\mathbf{Sp}(k)\mathbf{Sp}(1)$	Quaternion-Kähler	$\mathbb{H}$
7	$\mathbf{G}_2$	Exceptional I	$\mathbb{O}$
8	$\mathbf{Spin}(7)$	Exceptional II	$\mathbb{O}$

$$\mathbf{Spin}(7) := \langle S^7 \subset \mathbb{O}^\times \rangle \subset \mathbf{SO}(8)$$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	$\mathbb{H}$
$4k$	$\mathbf{Sp}(k)\mathbf{Sp}(1)$	Quaternion-Kähler	$\mathbb{H}$
7	$\mathbf{G}_2$	Exceptional I	$\mathbb{O}$
8	$\mathbf{Spin}(7)$	Exceptional II	$\mathbb{O}$

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	relevant field
n	$\mathbf{SO}(n)$	generic	$\mathbb{R}$
$2m$	$\mathbf{U}(m)$	Kähler	$\mathbb{C}$
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	$\mathbb{C}$
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	$\mathbb{H}$
$4k$	$\mathbf{Sp}(k)\mathbf{Sp}(1)$	Quaternion-Kähler	$\mathbb{H}$
7	$\mathbf{G}_2$	Exceptional I	$\mathbb{O}$
8	$\mathbf{Spin}(7)$	Exceptional II	$\mathbb{O}$

Simons, Aleksevskii, Calabi, Hitchin, Bryant, Joyce...

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	Einstein?
n	$\mathbf{SO}(n)$	generic	?
$2m$	$\mathbf{U}(m)$	Kähler	?
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	✓
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	✓
$4k$	$\mathbf{Sp}(k)\mathbf{Sp}(1)$	Quaternion-Kähler	✓
7	$\mathbf{G}_2$	Exceptional I	✓
8	$\mathbf{Spin}(7)$	Exceptional II	✓

Theorem (Berger 1955...). *Let (M^n, g) be a complete, simply-connected Riemannian manifold. If (M, g) is neither a product of lower-dimensional manifolds nor a symmetric space, then its holonomy group appears on the following table:*

dimension n	$\text{Hol}(M^n, g)$	geometry	Ricci-flat?
n	$\mathbf{SO}(n)$	generic	?
$2m$	$\mathbf{U}(m)$	Kähler	✗
$2m$	$\mathbf{SU}(m)$	Calabi-Yau	✓
$4k$	$\mathbf{Sp}(k)$	Hyper-Kähler	✓
$4k$	$\mathbf{Sp}(k)\mathbf{Sp}(1)$	Quaternion-Kähler	✗
7	$\mathbf{G}_2$	Exceptional I	✓
8	$\mathbf{Spin}(7)$	Exceptional II	✓

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

Challenge to the community:

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

Challenge to the community: find more!

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

Challenge to the community: find more!

Known examples arise from special holonomy!

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

Known examples have $n \equiv 0 \pmod{4}$.

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

Known examples have $n \equiv 0 \pmod{4}$.

What about odd dimensions?

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

Known examples have $n \equiv 0 \pmod{4}$.

What about odd dimensions?

What about $n \equiv 2 \pmod{4}$?

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

Known examples have $n \equiv 0 \pmod{4}$.

What about odd dimensions?

What about $n \equiv 2 \pmod{4}$?

What about coexistence of $\lambda = 0$ and $\lambda \neq 0$?

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

Known examples have $n \equiv 0 \pmod{4}$.

What about odd dimensions?

What about $n \equiv 2 \pmod{4}$?

What about coexistence of $\lambda = 0$ and $\lambda \neq 0$?

What about coexistence of $\lambda = 0$ and $\lambda > 0$?

We have seen that $\exists$ smooth closed M^n that admit Einstein metrics for both $\lambda > 0$ and $\lambda < 0$.

But the known examples remain rare and peculiar.

Known examples have $n \equiv 0 \pmod{4}$.

What about odd dimensions?

What about $n \equiv 2 \pmod{4}$?

What about coexistence of $\lambda = 0$ and $\lambda \neq 0$?

What about coexistence of $\lambda = 0$ and $\lambda > 0$?

The latter seems to be especially delicate!

What about odd dimensions?

What about odd dimensions?

G_2 is the only irreducible holonomy option.

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat.

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat. ($\lambda = 0$)

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat. ($\lambda = 0$)

Any such manifold has $|\pi_1(M)| \leq \infty$.

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat. ($\lambda = 0$)

Any such manifold has $|\pi_1(M)| \leq \infty$.

Gromov-Lawson: $\widetilde{M}^7$ admits metrics with $s > 0$.

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat. ($\lambda = 0$)

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat. ($\lambda = 0$)

Constructions of Einstein 7-manifolds of with $\lambda \neq 0$?

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat. ($\lambda = 0$)

Constructions of Einstein 7-manifolds of with $\lambda \neq 0$?

$\exists$ vast multiverse of Einstein (M^7, g) with $\lambda > 0$.

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat. ($\lambda = 0$)

Constructions of Einstein 7-manifolds of with $\lambda \neq 0$?

$\exists$ vast multiverse of Einstein (M^7, g) with $\lambda > 0$.

Sasaki-Einstein 7-manifolds (M^7, g) :

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat. ($\lambda = 0$)

Constructions of Einstein 7-manifolds with $\lambda \neq 0$?

$\exists$ vast multiverse of Einstein (M^7, g) with $\lambda > 0$.

Sasaki-Einstein 7-manifolds (M^7, g) :

Metric cone $(M^7 \times \mathbb{R}^+, dr^2 + r^2 g)$ holonomy $\mathbf{SU}(4)$.

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat. ($\lambda = 0$)

Constructions of Einstein 7-manifolds with $\lambda \neq 0$?

$\exists$ vast multiverse of Einstein (M^7, g) with $\lambda > 0$.

Sasaki-Einstein 7-manifolds (M^7, g) :

Metric cone $(M^7 \times \mathbb{R}^+, dr^2 + r^2 g)$ is Calabi-Yau.

What about odd dimensions?

$\mathbf{G}_2$ is the only irreducible holonomy option.

Holonomy $\mathbf{G}_2 \implies (M^7, g)$ Ricci-flat. ($\lambda = 0$)

Constructions of Einstein 7-manifolds with $\lambda \neq 0$?

$\exists$ vast multiverse of Einstein (M^7, g) with $\lambda > 0$.

Sasaki-Einstein 7-manifolds (M^7, g) :

Metric cone $(M^7 \times \mathbb{R}^+, dr^2 + r^2 g)$ is Calabi-Yau.

Can such metrics coexist?

Alas, no!

Alas, no!

Theorem. *No smooth compact M^7*

Alas, no!

Theorem. *No smooth compact M^7 can admit both a Sasaki-Einstein metric g_1*

Alas, no!

Theorem. *No smooth compact M^7 can admit both a Sasaki-Einstein metric g_1 and a metric g_2*

Alas, no!

Theorem. *No smooth compact M^7 can admit both a Sasaki-Einstein metric g_1 and a metric g_2 with holonomy $\subset \mathbf{G}_2$.*

Alas, no!

Theorem. *No smooth compact M^7 can admit both a Sasaki-Einstein metric g_1 and a metric g_2 with holonomy $\subset \mathbf{G}_2$.*

Key:

Alas, no!

Theorem. *No smooth compact M^7 can admit both a Sasaki-Einstein metric g_1 and a metric g_2 with holonomy $\subset \mathbf{G}_2$.*

Key:

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 ,*

Alas, no!

Theorem. *No smooth compact M^7 can admit both a Sasaki-Einstein metric g_1 and a metric g_2 with holonomy $\subset \mathbf{G}_2$.*

Key:

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

Alas, no!

Theorem. *No smooth compact M^7 can admit both a Sasaki-Einstein metric g_1 and a metric g_2 with holonomy $\subset \mathbf{G}_2$.*

Key:

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

To make this plausible,

Alas, no!

Theorem. *No smooth compact M^7 can admit both a Sasaki-Einstein metric g_1 and a metric g_2 with holonomy $\subset \mathbf{G}_2$.*

Key:

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

To make this plausible, will first illustrate assertion for prototypical examples due to S. Kobayashi '63.

Prototypical examples:

Prototypical examples:

$$\begin{array}{ccc} S^1 & \rightarrow & M^7 \\ & & \downarrow \\ & & X^6 \end{array}$$

Prototypical examples:

$$\begin{array}{ccc} S^1 & \rightarrow & M^7 \\ & & \downarrow \\ & & X^6 \end{array}$$

(X, h) Kähler-Einstein, $\lambda = 8$.

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \\ X^6 \end{array}$$

(X, h) Kähler-Einstein, $\lambda = 8$.

M unit circle bundle in $K^* = \Lambda^3 T^{1,0} X$.

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

(X, h) Kähler-Einstein, $\lambda = 8$.

M unit circle bundle in $K^* = \Lambda^3 T^{1,0} X$.

ϖ Riemannian submersion.

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

(X, h) Kähler-Einstein, $\lambda = 8$.

M unit circle bundle in $K^* = \Lambda^3 T^{1,0} X$.

ϖ Riemannian submersion.

Horizontal distribution: Chern connection for h .

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

(X, h) Kähler-Einstein, $\lambda = 8$.

M unit circle bundle in $K^* = \Lambda^3 T^{1,0} X$.

ϖ Riemannian submersion.

Horizontal distribution: Chern connection for h .

Fiber circles totally geodesic, of length $\pi/2$.

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

(X, h) Kähler-Einstein, $\lambda = 8$.

M unit circle bundle in $K^* = \Lambda^3 T^{1,0} X$.

ϖ Riemannian submersion.

Horizontal distribution: Chern connection for h .

Fiber circles totally geodesic, of length $\pi/2$.

(M, g) Einstein, $\lambda = 6$.

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

(X, h) Kähler-Einstein, $\lambda = 8$.

M unit circle bundle in $K^* = \Lambda^3 T^{1,0} X$.

ϖ Riemannian submersion.

Horizontal distribution: Chern connection for h .

Fiber circles totally geodesic, of length $\pi/2$.

(M, g) Einstein, $\lambda = 6$. “Sasaki-Einstein.”

Prototypical examples:

$$\begin{array}{ccc} S^1 & \rightarrow & M^7 \\ & \downarrow \varpi & \\ & X^6 & \end{array}$$

Prototypical examples:

$$\begin{array}{ccc} S^1 & \rightarrow & M^7 \\ & & \downarrow \varpi \\ & & X^6 \end{array}$$

Gysin sequence:

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^k(M) \rightarrow H^{k-1}(X) \xrightarrow{\mathbb{L}} H^{k+1}(X) \xrightarrow{\varpi^*} H^{k+1}(M) \rightarrow \cdots$$

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^k(M) \rightarrow H^{k-1}(X) \xrightarrow{\mathbf{L}} H^{k+1}(X) \xrightarrow{\varpi^*} H^{k+1}(M) \rightarrow \cdots$$

where $\mathbf{L} = \text{cup with Euler class } c_1(M) \text{ of } K^* \rightarrow X$.

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^3(M) \rightarrow H^2(X) \xrightarrow{\mathbf{L}} H^4(X) \xrightarrow{\varpi^*} H^4(M) \rightarrow \cdots$$

where $\mathbf{L} = \text{cup with Euler class } c_1(M) \text{ of } K^* \rightarrow X$.

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^3(M) \rightarrow H^2(X) \xrightarrow{\mathbf{L}} H^4(X) \xrightarrow{\varpi^*} H^4(M) \rightarrow \cdots$$

where $\mathbf{L} = \text{cup with Euler class } c_1(M) \text{ of } K^* \rightarrow X$.

But X is Fano! So $\mathbf{L} = \text{Lefschetz operator of } (X, h)$.

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^3(M) \rightarrow H^2(X) \xrightarrow{\mathbf{L}} H^4(X) \xrightarrow{\varpi^*} H^4(M) \rightarrow \cdots$$

where $\mathbf{L} = \text{cup with Euler class } c_1(M) \text{ of } K^* \rightarrow X$.

But X is Fano! So $\mathbf{L} = \text{Lefschetz operator of } (X, h)$.

Hard Lefschetz theorem for (X, h) :

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^3(M) \rightarrow H^2(X) \xrightarrow{\mathbf{L}} H^4(X) \xrightarrow{\varpi^*} H^4(M) \rightarrow \cdots$$

where $\mathbf{L} = \text{cup with Euler class } c_1(M) \text{ of } K^* \rightarrow X$.

But X is Fano! So $\mathbf{L} = \text{Lefschetz operator of } (X, h)$.

Hard Lefschetz theorem for (X, h) :

$$\mathbf{L} : H^2(X, \mathbb{R}) \xrightarrow{\cong} H^4(X, \mathbb{R})$$

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^3(M) \rightarrow H^2(X) \xrightarrow{\cong} H^4(X) \xrightarrow{\varpi^*} H^4(M) \rightarrow \cdots$$

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^3(M) \rightarrow H^2(X) \xrightarrow{\cong} H^4(X) \xrightarrow{\varpi^*} H^4(M) \rightarrow \cdots$$

Hence

$$\varpi^* : H^4(X, \mathbb{R}) \xrightarrow{\cong} H^4(M, \mathbb{R})$$

vanishes!

Prototypical examples:

$$\begin{array}{c} S^1 \rightarrow M^7 \\ \downarrow \varpi \\ X^6 \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^3(M) \rightarrow H^2(X) \xrightarrow{\cong} H^4(X) \xrightarrow{\varpi^*} H^4(M) \rightarrow \cdots$$

Hence

$$\varpi^* : H^4(X, \mathbb{R}) \xrightarrow{\cong} H^4(M, \mathbb{R})$$

vanishes!

But $p_1(M) = \varpi^* p_1(X)$, because

$$TM \cong \mathbb{R} \oplus \varpi^* TX.$$

Prototypical examples:

$$\begin{array}{ccc} S^1 & \rightarrow & M^7 \\ & & \downarrow \varpi \\ & & X^6 \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^3(M) \rightarrow H^2(X) \xrightarrow{\cong} H^4(X) \xrightarrow{\varpi^*} H^4(M) \rightarrow \cdots$$

Hence

$$\varpi^* : H^4(X, \mathbb{R}) \xrightarrow{\cong} H^4(M, \mathbb{R})$$

vanishes!

But $p_1(M) = \varpi^* p_1(X)$, because

$$TM \cong \mathbb{R} \oplus \varpi^* TX.$$

So $[p_1(M)]^{\mathbb{R}} \in H^4(M, \mathbb{R})$ vanishes.

Prototypical examples:

$$\begin{array}{ccc} S^1 & \rightarrow & M^7 \\ & \downarrow \varpi & \\ & X^6 & \end{array}$$

Gysin sequence:

$$\cdots \rightarrow H^3(M) \rightarrow H^2(X) \xrightarrow{\mathbb{L}} H^4(X) \xrightarrow{\varpi^*} H^4(M) \rightarrow \cdots$$

Hence

$$\varpi^* : H^4(X, \mathbb{R}) \xrightarrow{\cong} H^4(M, \mathbb{R})$$

vanishes!

But $p_1(M) = \varpi^* p_1(X)$, because

$$TM \cong \mathbb{R} \oplus \varpi^* TX.$$

So $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion element.

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

To prove general case, replace $H^*(X, \mathbb{R})$ with the basic cohomology $H_B^*(M, \mathfrak{F})$ of M relative to the Reeb foliation $\mathfrak{F}$.

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

To prove general case, replace $H^*(X, \mathbb{R})$ with the basic cohomology $H_B^*(M, \mathfrak{F})$ of M relative to the Reeb foliation $\mathfrak{F}$.

Replace Hard Lefschetz on $H^*(X, \mathbb{R})$ with transverse version on $H_B^*(M, \mathfrak{F})$ due to El Kacimi-Alaoui.

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

By contrast, if (M^7, g_2) has holonomy $\mathbf{G}_2$,

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

By contrast, if (M^7, g_2) has holonomy $\mathbf{G}_2$, then $p_1(M) \in H^4(M, \mathbb{Z})$ can't be a torsion class:

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

By contrast, if (M^7, g_2) has holonomy $\mathbf{G}_2$, then $p_1(M) \in H^4(M, \mathbb{Z})$ can't be a torsion class:

$$\langle p_1(M) \cup [\varphi], [M] \rangle = -\frac{1}{8\pi^2} \int_M |\mathcal{R}|^2 d\mu_{g_2} < 0,$$

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

By contrast, if (M^7, g_2) has holonomy $\mathbf{G}_2$, then $p_1(M) \in H^4(M, \mathbb{Z})$ can't be a torsion class:

$$\langle p_1(M) \cup [\varphi], [M] \rangle = -\frac{1}{8\pi^2} \int_M |\mathcal{R}|^2 d\mu_{g_2} < 0,$$

where $\mathcal{R}$ is Riemann curvature tensor and φ is the parallel 3-form determining g_2 .

Proposition. *If smooth compact M^7 carries a Sasaki-Einstein metric g_1 , then $p_1(M) \in H^4(M, \mathbb{Z})$ is a torsion class.*

By contrast, if (M^7, g_2) has holonomy $\mathbf{G}_2$, then $p_1(M) \in H^4(M, \mathbb{Z})$ can't be a torsion class:

$$\langle p_1(M) \cup [\varphi], [M] \rangle = -\frac{1}{8\pi^2} \int_M |\mathcal{R}|^2 d\mu_{g_2} < 0,$$

where $\mathcal{R}$ is Riemann curvature tensor and φ is the parallel 3-form determining g_2 .

Theorem. *No smooth compact M^7 can admit both a Sasaki-Einstein metric g_1 and a metric g_2 with holonomy $\subset \mathbf{G}_2$.*

What about other dimensions?

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 0 \pmod{4}$, then M spin, with $\hat{A}(M) \neq 0$:

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 0 \pmod{4}$, then M spin, with $\hat{A}(M) \neq 0$:

$$\hat{A}(M) = \left\{ \right.$$

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 0 \pmod{4}$, then M spin, with $\hat{A}(M) \neq 0$:

$$\hat{A}(M) = \begin{cases} 2, & \text{if Hol} = \mathbf{SU}(2m) \text{ and } n = 2m, \\ \end{cases}$$

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 0 \pmod{4}$, then M spin, with $\hat{A}(M) \neq 0$:

$$\hat{A}(M) = \begin{cases} 2, & \text{if Hol} = \mathbf{SU}(2m) \text{ and } n = 2m, \\ k + 1, & \text{if Hol} = \mathbf{Sp}(k) \text{ and } n = 4k, \end{cases}$$

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 0 \pmod{4}$, then M spin, with $\hat{A}(M) \neq 0$:

$$\hat{A}(M) = \begin{cases} 2, & \text{if Hol} = \mathbf{SU}(2m) \text{ and } n = 2m, \\ k + 1, & \text{if Hol} = \mathbf{Sp}(k) \text{ and } n = 4k, \\ 1, & \text{if Hol} = \mathbf{Spin}(7) \text{ and } n = 8. \end{cases}$$

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 0 \pmod{4}$, then M spin, with $\hat{A}(M) \neq 0$:

$$\hat{A}(M) = \begin{cases} 2, & \text{if Hol} = \mathbf{SU}(2m) \text{ and } n = 2m, \\ k + 1, & \text{if Hol} = \mathbf{Sp}(k) \text{ and } n = 4k, \\ 1, & \text{if Hol} = \mathbf{Spin}(7) \text{ and } n = 8. \end{cases}$$

$\Rightarrow \nexists$ metrics with $s > 0$.

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 0 \pmod{4}$, then M spin, with $\hat{A}(M) \neq 0$:

$$\hat{A}(M) = \begin{cases} 2, & \text{if Hol} = \mathbf{SU}(2m) \text{ and } n = 2m, \\ k + 1, & \text{if Hol} = \mathbf{Sp}(k) \text{ and } n = 4k, \\ 1, & \text{if Hol} = \mathbf{Spin}(7) \text{ and } n = 8. \end{cases}$$

$\Rightarrow \nexists$ metrics with $s > 0$.

$\Rightarrow \nexists$ Einstein metrics with $\lambda > 0$.

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 2 \pmod{8}$, then M spin, with $\alpha(M) \neq 0$.

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 2 \pmod{8}$, then M spin, with $\alpha(M) \neq 0$.

Here $\text{Hol} = \mathbf{SU}(4\ell + 1)$ in dimension $n = 8\ell + 2$.

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 2 \pmod{8}$, then M spin, with $\alpha(M) \neq 0$.

Here $\text{Hol} = \mathbf{SU}(4\ell + 1)$ in dimension $n = 8\ell + 2$.

$\alpha(M) \in \mathbb{Z}_2$ means $\dim \ker$ Dirac operator, mod 2.

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 2 \pmod{8}$, then M spin, with $\alpha(M) \neq 0$.

Here $\text{Hol} = \mathbf{SU}(4\ell + 1)$ in dimension $n = 8\ell + 2$.

$\alpha(M) \in \mathbb{Z}_2$ means $\dim \ker$ Dirac operator, mod 2.

$\implies \nexists$ metrics with $s > 0$.

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

If $n \equiv 2 \pmod{8}$, then M spin, with $\alpha(M) \neq 0$.

Here $\text{Hol} = \mathbf{SU}(4\ell + 1)$ in dimension $n = 8\ell + 2$.

$\alpha(M) \in \mathbb{Z}_2$ means $\dim \ker$ Dirac operator, mod 2.

$\implies \nexists$ metrics with $s > 0$.

$\implies \nexists$ Einstein metrics with $\lambda > 0$.

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

This leaves us with Calabi-Yau manifolds of real dimension $8\ell + 6$.

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

This leaves us with Calabi-Yau manifolds of real dimension $8\ell + 6$.

Gromov-Lawson, Stolz: Always have $s > 0$ metrics on simply-connected manifolds of these dimensions!

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

This leaves us with Calabi-Yau manifolds of real dimension $8\ell + 6$.

Gromov-Lawson, Stolz: Always have $s > 0$ metrics on simply-connected manifolds of these dimensions!

For example, every Calabi-Yau 3-fold admits $s > 0$ metrics.

What about other dimensions?

Can Ricci-flat metrics of special holonomy ever co-exist with $\lambda > 0$ Einstein?

If Ricci-flat metric is holonomy irreducible, impossible in most allowed dimensions!

This leaves us with Calabi-Yau manifolds of real dimension $8\ell + 6$.

Gromov-Lawson, Stolz: Always have $s > 0$ metrics on simply-connected manifolds of these dimensions!

For example, every Calabi-Yau 3-fold admits $s > 0$ metrics. What about $\lambda > 0$ Einstein?

Coexistence Problem in Dimension 6:

Coexistence Problem in Dimension 6:

Is there a smooth closed M^6 that admits both a Kähler-Einstein metric g_1 with $\lambda > 0$ and a Calabi-Yau metric g_0 ?

Coexistence Problem in Dimension 6:

Is there a smooth closed M^6 that admits both a Kähler-Einstein metric g_1 with $\lambda > 0$ and a Calabi-Yau metric g_0 ?

Completely open!

Coexistence Problem in Dimension 6:

Is there a smooth closed M^6 that admits both a Kähler-Einstein metric g_1 with $\lambda > 0$ and a Calabi-Yau metric g_0 ?

Completely open!

Only four candidates for (M, J_1) :

Coexistence Problem in Dimension 6:

Is there a smooth closed M^6 that admits both a Kähler-Einstein metric g_1 with $\lambda > 0$ and a Calabi-Yau metric g_0 ?

Completely open!

Only four candidates for (M, J_1) :

- sextic hypersurface in $\mathbb{CP}(1, 1, 1, 2, 3)$;
- double cover of $\mathbb{CP}_3$ branched over quartic;
- cubic hypersurface in $\mathbb{CP}_4$;
- transverse intersection of two quadrics in $\mathbb{CP}_5$.

Coexistence Problem in Dimension 6:

Is there a smooth closed M^6 that admits both a Kähler-Einstein metric g_1 with $\lambda > 0$ and a Calabi-Yau metric g_0 ?

Completely open!

Only four candidates for (M, J_1) :

- sextic hypersurface in $\mathbb{CP}(1, 1, 1, 2, 3)$;
- double cover of $\mathbb{CP}_3$ branched over quartic;
- cubic hypersurface in $\mathbb{CP}_4$;
- transverse intersection of two quadrics in $\mathbb{CP}_5$.

These all admit K-E metrics with $\lambda > 0$.

Coexistence Problem in Dimension 6:

Is there a smooth closed M^6 that admits both a Kähler-Einstein metric g_1 with $\lambda > 0$ and a Calabi-Yau metric g_0 ?

Completely open!

Only four candidates for (M, J_1) :


- sextic hypersurface in $\mathbb{CP}(1, 1, 1, 2, 3)$;
- double cover of $\mathbb{CP}_3$ branched over quartic;
- cubic hypersurface in $\mathbb{CP}_4$;
- transverse intersection of two quadrics in $\mathbb{CP}_5$.

These all admit K-E metrics with $\lambda > 0$.

But no one has found Calabi-Yau partners for these Fano manifolds!


Vielen Dank für diese Einladung!

Vielen Dank für diese Einladung!



MPI Mathematik

Thanks for the invitation!



MPI Mathematik

Ich freue mich sehr, hier zu sein!

Ich freue mich sehr, hier zu sein!



It's a real pleasure to be here!



It's a real pleasure to be here!

