

## On the complexity of some birational transformations

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2006 J. Phys. A: Math. Gen. 39 3641

(<http://iopscience.iop.org/0305-4470/39/14/009>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

### Download details:

IP Address: 140.182.75.117

This content was downloaded on 15/09/2016 at 12:13

Please note that [terms and conditions apply](#).

You may also be interested in:

[How to detect the integrability of discrete systems](#)

B Grammaticos, R G Halburd, A Ramani et al.

[A classification of four-state spin edge Potts models](#)

J-Ch Anglès d'Auriac, J-M Maillard and C M Viallet

[On the algebraic structure of rational discrete dynamical systems](#)

C-M Viallet

[Algebraic entropy of an extended Hietarinta–Viallet equation](#)

Masataka Kanki, Takafumi Mase and Tetsuji Tokihiro

[The discrete Painlevé I equations:transcendental integrability and asymptotic solutions](#)

M Bernardo, T T

Truong and G Rollet

[Algebraic entropy for algebraic maps](#)

A N W Hone, Orlando Ragnisco and Federico Zullo

[Integrable mappings from matrix transformations and their singularity properties](#)

A Ramani, B Grammaticos, J -M Maillard et al.

# On the complexity of some birational transformations

J-Ch Anglès d'Auriac<sup>1</sup>, J-M Maillard<sup>2,3</sup> and C M Viallet<sup>2,3</sup>

<sup>1</sup> Centre de Recherches sur les Très Basses Températures, BP 166, 38042 Grenoble, France

<sup>2</sup> LPTMC, Case 121, 4 Place Jussieu, 75252 Paris Cedex 05, France

<sup>3</sup> LPTHE, Boîte 126, 4 Place Jussieu, 75252 Paris Cedex 05, France

E-mail: [dauriac@grenoble.cnrs.fr](mailto:dauriac@grenoble.cnrs.fr), [maillard@lptmc.jussieu.fr](mailto:maillard@lptmc.jussieu.fr) and [viallet@lpthe.jussieu.fr](mailto:viallet@lpthe.jussieu.fr)

Received 6 December 2005

Published 22 March 2006

Online at [stacks.iop.org/JPhysA/39/3641](http://stacks.iop.org/JPhysA/39/3641)

## Abstract

Using three different approaches, we analyse the complexity of various birational maps constructed from simple operations (inversions) on square matrices of arbitrary size. The first approach comprises the study of the images of lines, and relies mainly on univariate polynomial algebra, the second approach is a singularity analysis and the third method is more numerical, using integer arithmetics. These three methods have their own domain of application, but they give corroborating results, and lead us to a conjecture on the algebraic entropy of a class of maps constructed from matrix inversions.

PACS numbers: 02.40.Xx, 05.50.+q, 05.10.–a

## 1. Presentation

We investigate some properties of birational realizations of Coxeter groups on projective spaces of various dimensions. There are many motivations to examine these realizations. They appear naturally in the study of ‘spin models’ of statistical mechanics, in relation with the symmetries of the Yang–Baxter (alias star-triangle equations) [1, 2] as well as through their connection with association schemes and knot polynomials [3–8]. Among the basic ingredients in all these applications are various inversions of matrices (matrix inverse, block inverses, element by element inverse).

The typical group we consider is a Coxeter group with two involutive generators  $I$  and  $J$ , and no extra relations between  $I$  and  $J$ . They generate a discrete group  $\Gamma$ , the infinite dihedral group, isomorphic to the semi-direct product  $\mathbb{Z} \rtimes \mathbb{Z}_2$ . The action we analyse is the one of the infinite parts of  $\Gamma$ , i.e. to say the iterates of  $K = I \circ J$ .

Studying the iterates of  $K$  is the same thing as studying the rational (in fact bi-rational) discrete dynamical system. Such an investigation is a part of the algebraic dynamics, and we focus on the analysis of the complexity of the iterates [9–22].

We perform this analysis for a definite class of transformations, defined from elementary operations on matrices of size  $q \times q$ , the entries of the matrices being the natural coordinates of complex projective spaces  $\mathbb{C}\mathbb{P}_n$ . Depending on the specific form of the matrices, the dimension  $n$  will take different values ( $n \leq q^2 - 1$ ).

Another motivation for our choice is that arbitrary algebraic transformations are not invertible. Being by construction rational and invertible, matrix inversions constitute a factory of almost everywhere invertible transformations of any degree, and with any number of variables. They provide us with a variety of explicit birational dynamics. The specific choice we made here for the form of the matrices (see also [23, 24]) is motivated by their use in lattice statistical mechanics, and the richness of the structures of the systems we construct.

We explain, exemplify and confront three different approaches to the problem. We also present a conjecture for the value of the algebraic entropy for a family of transformations of interest to statistical mechanics.

The paper is organized as follows. We state in section 2 the problem of calculating the complexity of a birational transformation acting on a projective space, and define the basic objects of interest, in particular the algebraic entropy or equivalently the rate of growth of the degrees of the iterates of a map. We introduce four families of maps, which will be used for explicit calculations. In section 3 we indicate how to surmise the generating function of the sequence of degree of iterates of a map from its first terms. This provides a first method of calculation of the complexity. In section 4, we calculate exactly the sequence of degrees by an analysis of the singularity structure for one of the families of maps. In section 5, we describe an arithmetic approach, where we examine the action of iterates on rational points (integer homogeneous coordinates), and simply measure the growth of the size of the coordinates. This yields approximate values of the complexity. We conclude with a conjecture.

## 2. The problem

Let  $K$  be a birational transformation of complex projective space  $\mathbb{C}\mathbb{P}_n$ . If we write  $K$  in terms of homogeneous coordinates, it appears as a polynomial transformation given by  $n + 1$  homogeneous polynomials of the same degree  $d$ . With the rule that we should factorize out any common factor,  $d$  is well defined in a given system of coordinates. Of course it is *not* invariant by changes of coordinates. We may construct the sequence  $\{d_n\}$  of the degrees of the iterates  $K^n$  of  $K$ .

We will use [15], as a measure of the complexity of  $K$ , the growth of the sequence  $d_n$ : in the absence of factorizations of the polynomials the sequence would just be

$$d_n = d_1^n = d^n. \quad (1)$$

What happens is that if some factorizations appear, they induce a drop of the degree, so that we only have an upper bound

$$d_n \leq d^n. \quad (2)$$

The drop may even be so important that the growth of  $d_n$  becomes polynomial or is bounded. A measure of the growth is the algebraic entropy [15]<sup>4</sup>

$$\epsilon = \lim_{n \rightarrow \infty} \frac{1}{n} \log d_n, \quad (3)$$

or the rate of growth

$$\lambda = \exp(\epsilon). \quad (4)$$

<sup>4</sup> The existence of the limit is a consequence of the elementary relation  $d_{n+m} \leq d_n \cdot d_m$ .

Both the entropy  $\epsilon$  and the rate of growth  $\lambda$  are invariant by any birational change of coordinates. They are canonically associated with the map  $K$ . Our aim is to calculate them for definite classes of maps, which we now describe.

Suppose  $M$  is a  $q \times q$  matrix (the ‘ $q$ ’ is reminiscent of the  $q$ -state Potts model of statistical mechanics), and consider the two simple rational involutions  $I$  and  $J$ : the involution  $I$  is the matrix inverse up to a factor (i.e. when written polynomially it amounts to replacing each entry by its cofactor). The involution  $J$  is the element by element inverse (also called a Hadamard inverse, which replaces each entry  $M_{ij}$  by its inverse  $1/M_{ij}$ ). The two involutions  $I$  and  $J$  do not commute, and their composition  $K = I \circ J$  is generically of infinite order.

The map  $K$  acts naturally on  $\mathbb{C}\mathbb{P}_{q^2-1}$ . It is however possible to define various reductions to smaller projective spaces in the following way [25]. For a given size of square matrices, we define a pattern as a set of equalities between entries of the matrix. The set of all patterns is the set of all partitions of the entries of the matrix. An example of a pattern is ‘all diagonal entries equal, all off-diagonal entries equal’. This corresponds to the partition of the entries in two parts (diagonal + off-diagonal). Clearly any pattern is preserved by the action of  $J$ . We call admissible a pattern which is also stable by  $I$  (or equivalently  $K$ ).

All admissible patterns have been classified for  $q = 4$  and some of them for  $q = 5$  in [23–26]. It has been also shown that  $\lambda$  can vary considerably from one admissible pattern to another. For example for  $5 \times 5$  cyclic and symmetric matrices one has  $\lambda = 1$  (polynomial growth), whereas with the cyclic matrices one gets  $\lambda = (7 + 3\sqrt{5})/2$ .

We will focus on four fundamental admissible patterns, which exist whatever the size  $q$  of the matrices is. The first one is the pattern ( $S$ ) of symmetric matrices. The second one ( $C$ ) is the pattern of the cyclic matrices defined by  $M_{i,j} = M_{i+1,j+1}$  (with indices taken modulo  $q$ ). The third one is the pattern of matrices which are at the same time cyclic and symmetric ( $CS$ ). The last one is the general pattern ( $G$ ), without equality conditions between the entries. From the results obtained on these different patterns, we conjecture that, contrary to intuition and although their number of variables differ enormously,  $\lambda$  is the same for cyclic ( $S$ ), symmetric ( $S$ ) and general ( $G$ ) patterns.

### 3. A first approach: generating functions

From the sequence of degrees  $\{d_n\}$ , it is possible to construct a generating function

$$f(u) = \sum_{n=0}^{\infty} d_n u^n. \tag{5}$$

Since the degrees are bounded by (2), the series (5) always has a non-zero radius of convergence  $\rho$ . Actually

$$\rho = \frac{1}{\lambda}, \tag{6}$$

with  $\lambda$  being the rate of growth defined above.

The calculation method is the following: calculate explicitly the first terms of the series, and try to guess the values of the generating function. The method is sensible if the generating function is rational.

The striking fact is that indeed the generating function  $f(u)$  happens to be a rational fraction with integer coefficients in most cases. The consequence is that a finite number of terms of the series determine it completely. For reversible maps (i.e. when there exists a similarity relation between the map and its inverse), we have not found any counterexample to this rule. There are however non-reversible maps for which the generating function is not

**Table 1.** Generating functions for the cyclic symmetric (*CS*) patterns. The formulae for values of  $q$  tagged with a  $(\star)$  can be proved.  $n_{\max}$  is the maximum number of iteration performed,  $m$  refers to the Padé approximation,  $\lambda$  is the rate of growth and  $\lambda_{\text{num}}$  is the numerical complexity calculated in section 5.

| $q$          | $f_q(u)$   | $n_{\max}$ | $m$ | $\lambda$ | $\lambda_{\text{num}}$ |
|--------------|--|------------|-----|-----------|------------------------|
| 4 $(\star)$  | $\frac{(1+u)^2}{(1-u)^2}$  | $\infty$   | 4   | 1         |                        |
| 5 $(\star)$  | $\frac{(1+u+2u^2)^2}{(1-u)^3(1+u+u^2)}$                          | 14         | 9   | 1         | 1.00                   |
| 6            | $\frac{(1+2u)^2}{(1-u)(1-4u)}$                                   | 15         | 4   | 4         | 4.00                   |
| 7 $(\star)$  | $\frac{(1+u+3u^2)^2}{(1-u)(1+u+u^2)(1-7u+u^2)}$                  | 12         | 9   | 6.854 10  | 6.85                   |
| 8            | $\frac{(1+u)(1+2u-u^2)}{(1-u)(1-11u+7u^2-u^3)}$                  | 11         | 7   | 10.331 85 | 10.33                  |
| 9 $(\star)$  | $\frac{(1+u+3u^2-3u^3)^2}{(1-u)(1-13u+2u^2+u^3+12u^4-8u^5+u^6)}$ | 11         | 13  | 12.832 69 | 12.83                  |
| 10           | $\frac{(1+3u)^2}{(1-u)(1-18u+u^2)}$                              | 9          | 5   | 17.944 27 | 17.94                  |
| 11 $(\star)$ | $\frac{(1+u+5u^2)^2}{(1-u)(1+u+u^2)(1-23u+u^2)}$                 | 7          | 9   | 22.956 44 | 22.95                  |
| 12           | $\frac{(1+4u-3u^2)(1+2u-u^2)}{(1-u)(1-27u+31u^2-9u^3)}$          | 8          | 8   | 25.812 54 | 25.81                  |
| 13 $(\star)$ | $\frac{(1+u+6u^2)^2}{(1-u)(1+u+u^2)(1-34u+u^2)}$                 |            | 9   | 33.970 56 | 33.97                  |

rational [22]. Another consequence of the rationality of  $f$  is that  $\lambda$  is an algebraic integer, and we have no counterexample yet to that.

For practical purposes, it is necessary to push the calculation of the degree of the iterates as far as possible. Instead of evaluating the full iterate, it is sufficient to consider the image of a generic line  $l$  with a running point

$$l(t) = [a_0 + b_0t, a_1 + b_1t, \dots, a_n + b_nt], \quad (7)$$

where  $a_i, b_i$  are arbitrary coefficients, and to evaluate the images of  $l(t)$  by  $K^n$ . The degree  $d_n$  is read off from this image. The calculation may furthermore be improved by using integer coefficients in (7) and calculating (formal calculation software are quite efficient at that) over polynomial with coefficients in  $\mathbb{Z}/\mathbb{Z}_p$  with  $p$  being a sufficiently large prime integer. Taking different values of  $p$  and of the coefficients  $a_i, b_i$  helps eliminate the accidental simplifications which may occur.

Suppose we have the degree  $d_n$  for the first values of  $n$ , say  $n = 1 \dots n_{\max}$ . We may fit the series with a Padé approximant  $F$ , with numerator (respectively denominator) of degree  $N$  (respectively  $M$ ), such that

$$N + M = n_{\max} - 1 \quad (8)$$

$N$  running from 0 to  $n_{\max} - 1$ . Our experience is that, if  $n_{\max}$  is large enough, the rational fraction  $F$  we find simplifies drastically, and stabilizes for some central values of  $N$  (i.e. the numerator and denominator are respectively of degree smaller than  $M$  and  $N$ ). This usually means that the exact generating function has been reached.

Note that the expansion of the non-optimal  $[N, M]$  Padé approximants yields non-integer, or negative coefficients in the expansion of  $F$ , in contradiction with these coefficients being a degree. Table 1 displays the 'exact' expression we have inferred for the generating function for various values of  $q$  for the (*CS*) pattern, as well as the value of  $m = N + M$  and the value of  $n_{\max}$ .

When  $n_{\max}$  is larger than  $m$ , we have a prediction on the next values of the degree, and this gives confidence that the result is exact.

In table 1, we also give the inverse of the modulus of the smallest zero of the denominator, as well as a numerical value computed as explained in section 5.

**4. A second approach: singularity analysis**

In this section we prove that the rate of growth of the patterns (CS) for prime  $q$  is a quadratic integer, by showing that the sequence of degrees verifies a linear recurrence relation of length 2 with integer coefficients. This implies that the generating function of the degrees is a rational fraction and corroborates a part of the results given in table 1.

*4.1. Some notations*

Let  $M$  be a cyclic symmetric matrix of size  $q \times q$ . The matrix  $M$  may be written in terms of the basic cycle of order  $q$ :

$$\sigma = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 \\ 1 & 0 & 0 & \dots & 0 \end{pmatrix}$$

$$M_{\text{even}} = x_0 + x_1(\sigma + \sigma^{q-1}) + \dots + x_{p-1}\sigma^{q/2}, \quad p = \frac{q}{2} + 1$$

$$M_{\text{odd}} = x_0 + x_1(\sigma + \sigma^{q-1}) + \dots + x_{p-1}(\sigma^{(q-1)/2} + \sigma^{(q+1)/2}), \quad p = \frac{q+1}{2}$$

when  $q$  is even and odd, respectively.

The parameter space is a projective space  $\mathbb{C}P_{p-1}$  of dimension  $p - 1$ , with  $p = q/2 + 1$  if  $q$  is even and  $p = (q + 1)/2$  if  $q$  is odd. We use homogeneous coordinates  $[x_0, \dots, x_{p-1}]$ .

We will study the two elementary transformations  $I$  and  $J$  acting on  $M$ . Both are rational involutions (and are thus birational transformations).

The Hadamard inverse  $J$  may be written polynomially in terms of the homogeneous coordinates

$$J : [x_0, \dots, x_{p-1}] \longrightarrow \left[ \prod_{k \neq 0} x_k, \prod_{k \neq 1} x_k, \dots, \prod_{k \neq p-1} x_k \right]. \tag{9}$$

The matrix inverse  $I$ , up to a factor, transforms cyclic matrices into cyclic matrices, and symmetric matrices into symmetric matrices. It thus acts on cyclic symmetric matrices.

For cyclic symmetric matrices, the matrix inverse  $I$  is related to the Hadamard inverse  $J$ , by a similarity transformation:

$$I = C^{-1} \circ J \circ C. \tag{10}$$

The transformation  $C$  acts linearly on the  $p$  homogeneous coordinates. Denoting  $\omega$  the  $q$ th root of unity,  $C$  is given by the  $p \times p$  matrix with entries:

$$C_{r,0} = 1, \quad C_{r,s} = \omega^{rs} + \frac{1}{\omega^{rs}} \quad r = 0, \dots, p - 1, \quad s = 1, \dots, p - 1 \tag{11}$$

for  $q$  odd and

$$C_{r,0} = 1, \quad C_{r,s} = \omega^{rs} + \frac{1}{\omega^{rs}} \quad r = 0, \dots, p - 1, \quad s = 1, \dots, p - 2 \tag{12}$$

$$C_{r,p-1} = (-1)^r$$

for  $q$  even.

The matrix  $C$  verifies  $C^2 = \mathbf{1}$ .




Figure 1. Successive images.

#### 4.2. Sequences of surfaces and degrees

Consider now a sequence of hypersurfaces in  $\mathbb{C}\mathbb{P}^{p-1}$ , obtained by applying successively  $I$ ,  $J$ , then  $I$  and so on, starting with a generic hyperplane  $S_0$  (see figure 1). Each surface  $S_n$  has a polynomial equation, of degree  $d_n$ , which we also denote as  $S_n$ . Since for non-singular points,

$$x \in S_{2n} \iff J(x) \in S_{2n-1}, \quad (13)$$

$S_{2n}$  can be obtained from  $S_{2n-1}$  by substituting the coordinates of  $x$  with the homogeneous polynomial expressions of the coordinates of  $J(x)$  in  $S_{2n-1}(x)$ . Note that, since  $J$  is an involution,  $S_{2n-1}$  may be obtained from  $S_{2n}$  in the same manner.

What happens at the level of the equations is that  $S_{2n-1}(J(x))$  may factorize. One of the factors is  $S_{2n}(x)$ . The only other possible factors are powers of the coordinates of  $x = (x_0, \dots, x_{p-1})$  as explained in the lemma below. The relation

$$S_{2n-1}(J(x)) = S_{2n}(x) \cdot \prod_{k=0}^{p-1} x_k^{\alpha_{2n-1}^{(k)}} \quad (14)$$

defines the exponents  $\alpha_{2n-1}^{(k)}$ .

#### 4.3. A lemma

The previous relation is crucial. Its proof is elementary and goes as follows.

Suppose  $B$  is a birational involution. When written in terms of the homogeneous coordinates,  $B^2$  appears as the multiplication by some common polynomial factor of all the coordinates, that is to say the identity transformation in projective space.

$$B(B(x)) = [\kappa_B(x) \cdot x_0, \kappa_B(x) \cdot x_2, \dots, \kappa_B(x) \cdot x_{p-1}] \quad (15)$$

with  $\kappa_B(x) =$  some polynomial.

We then have, if two algebraic hypersurfaces  $S$  and  $S'$  are the proper images of each other by involution  $B$ :

$$S(B(x)) = S'(x) \cdot R(x) \quad (16)$$

$$S'(B(x)) = S(x) \cdot T(x) \quad (17)$$

with  $R$  and  $T$  being some polynomial expressions of the coordinates. We then have, using (16) and (17):

$$\kappa_B(x)^{\deg(S)} \cdot S(x) = S(x) \cdot R(B(x)) \cdot T(x), \tag{18}$$

i.e. to say

$$\kappa_B(x)^{\deg(S)} = R(B(x)) \cdot T(x). \tag{19}$$

Equation (19) shows that the only factors in the right-hand side of equation (17) are the equations of  $S$ , and polynomial expressions  $T(x)$  which divide  $\kappa_B(x)$ , possibly raised to some power.

In the specific example  $B = J$ , and  $S = S_{2n-1}$ , using

$$\kappa_J(x) = \prod_{i=0}^{p-1} x_i^{p-2} \tag{20}$$

we obtain

$$S_{2n-1}(J(x)) = S_{2n}(x) \cdot \prod_{i=0}^{p-1} x_i^{\rho_i}, \tag{21}$$

with  $x_i(x)$  being the  $i$ th coordinate of  $t$  and  $\rho_i$  some integer power.

This ends the proof of formula (14).

#### 4.4. Recurrence relation

Similar to equation (14), we have

$$S_{2n}(J(x)) = S_{2n-1}(x) \cdot \prod_{i=0}^{p-1} x_i^{\alpha_{2n}^{(i)}}, \tag{22}$$

with the constraint

$$\kappa_J(x)^{d_{2n}} = \prod_{i=0}^{p-1} x_i^{\alpha_{2n-1}^{(i)}}(x) \cdot \prod_{j=0}^{p-1} x_j^{\alpha_{2n}^{(j)}}(J(x)). \tag{23}$$

We also have the corresponding equations for the action of  $I$ .

$$S_{2n}(I(x)) = S_{2n+1}(x) \cdot \prod_{k=0}^{p-1} X_k^{\beta_{2n}^{(k)}} \tag{24}$$

$$S_{2n+1}(I(x)) = S_{2n}(x) \cdot \prod_{i=0}^{p-1} X_i^{\beta_{2n+1}^{(i)}}, \tag{25}$$

where  $X_i$  is the  $i$ th coordinate of  $Cx$ .

To make relations more uniform, we introduce a slight change of notation: define the sequences  $\{u_n^i\}$  and  $\{v_n^i\}$  with the convention that

$$\alpha_{2k+1}^i = u_{2k+1}^i, \quad \alpha_{2k}^i = v_{2k}^i, \tag{26}$$

$$\beta_{2k+1}^i = v_{2k+1}^i, \quad \beta_{2k}^i = u_{2k}^i. \tag{27}$$

At step  $n$  we have  $2p + 1$  variables ( $d_n, u_n^i$  and  $v_n^i$ ).



A first equation simply expresses the factorization:

$$d_n = (p - 1)d_{n-1} - \sum_{i=0}^{p-1} u_{n-1}^i. \quad (28)$$

Another set of equations is obtained by expressing that both  $I$  and  $J$  are involutions:

$$(p - 2)d_n = v_{n-1}^i + \sum_{j \neq i} u_n^j, \quad i = 0 \dots p - 1. \quad (29)$$

It is easy to obtain from equations (28) and 29:

$$v_n^i = (p - 2)d_{n-1} + u_{n-1}^i - \sum_{j=0}^{p-1} u_{n-1}^j, \quad i = 0 \dots p - 1. \quad (30)$$

#### 4.5. Singularity structure

We need  $p$  additional equations to complete the previous system. They will be given, under some constraints, by the analysis of the singularity structure. The basic idea is that the numbers  $\alpha_n^i, \beta_n^i$  (or equivalently  $u_n^i, v_n^i$ ) have a geometrical meaning: they are the multiplicity of some specific points on the surface  $S_n$ .

The singularity structure of  $J$  is very simple. A singular point is a point whose image is undetermined: this happens when all polynomial expressions giving the image (9) vanish simultaneously. Any point with more than two vanishing coordinates is singular for  $J$ .

We will look at the action of the pair  $I, J$  on the hypersurfaces composing the factor  $\kappa_J$  of equation (23). Those are just the  $n$  hyperplanes  $\Pi_k, k = 0 \dots p - 1$  of equation

$$\Pi_k : \quad \{x_k = 0\}. \quad (31)$$

All intersections of these hyperplanes are made out of singular points of  $J$ . Some points are in a sense *maximally* singular. They are the intersections of all but one of the planes  $\Pi_r$ , i.e. all but one of their coordinates vanish. There are  $p$  such points

$$P_k = [0, \dots, 0, 1, 0, \dots, 0], \quad k = 0 \dots p - 1 \quad (32)$$

with 1 in  $(k + 1)$  th position.

The singularity structure of  $I$  is the same as that of  $J$ , up to the linear change of coordinates  $C$ . There are in particular  $p$  distinguished singular points  $Q_k, k = 1 \dots p$  of  $I$ :

$$Q_k = C^{-1}P_k, \quad k = 0 \dots p - 1. \quad (33)$$

To complete the set of equations (28), (29), (30), we need to explore in some more details the singularity structure of the maps. What matters is the interplay between  $I$  and  $J$ .

The map  $J$  sends the hyperplane  $\Pi_k$  (31) onto the point  $P_k$  (32). The subsequent images depend on what  $q$  is.

The situation is tractable when  $q$  is a prime number, in which case the subsequent images of  $\Pi_k$  *always go back to the point  $P_k$  after a finite number of steps*, actually one or three steps. There, we meet a singularity, *and the equation of  $\Pi_k$  factorizes*. We will examine the case where  $q$  is a prime number,  $q = 2p - 1$ .

The coordinate  $x_0$  plays a special role and the point  $P_0$  behaves differently from the other points  $P_s, s = 1 \dots p - 1$ .

Whatever  $q$ , the transformation of the hyperplane  $\Pi_0$  reads

$$\Pi_0 \xrightarrow{J} P_0 \xrightarrow{I} P_0 \xrightarrow{J} \Pi_0. \quad (34)$$

We use the following convention concerning the arrows: when a variety is sent by the map onto a variety of same codimension we use the plain arrow  $\longrightarrow$ . When the codimension of the image is lower (blow-down) we use the symbol  $\succrightarrow$ , and when it is larger (blow-up) we use the squiggly arrow  $\rightsquigarrow$ . A blow-up for the birational mapping  $K$  corresponds to a blow-down for its inverse  $K^{-1}$ .

The action of  $I$  and  $J$  on the hyperplane  $\Pi_s$  reads

$$\Pi_s \xrightarrow{J} P_s \xrightarrow{I} R_s \xrightarrow{J} R_s \xrightarrow{I} P_s \rightsquigarrow \Pi_s. \tag{35}$$

The points  $R_s$  have coordinates  $[\pm 1, \pm 1, \dots, \pm 1]$ . For example for  $q = 5$ ,  $R_2 = [+1, +1, -1]$  and  $R_3 = [+1, -1, +1]$ , while, for  $q = 7$ ,  $R_2 = [+1, -1, -1, +1]$ ,  $R_3 = [-1, -1, +1, +1]$  and  $R_4 = [-1, +1, -1, +1]$ .

The pattern is similar for the points  $Q_k$ . It is obtained from the previous one by the linear change of coordinates defined by  $C$ . The planes  $\Pi_k$  are replaced by the planes  $\Pi'_k = C^{-1}\Pi_k$  and the points  $R_s$  are replaced by the points  $R'_s = C^{-1}R_s$ .

When  $q$  is not a prime number, the pattern is different: the successive actions of  $I$  and  $J$  lead to singular points other than the  $P_k$ 's and  $Q_k$ 's. In appendix A the case  $q = 9$  is studied as an example.

Relations (34), (35) allow us to relate the multiplicities of the singular points  $P_k$  on different surfaces  $S_n$ . Since  $P_0 \rightarrow P_0$  in (equation 34) we have

$$u_n^0 = v_{n-1}^0 \tag{36}$$

and since  $P_s \rightarrow P_s$  in (equation 35) we obtain

$$u_n^s = v_{n-3}^s, \quad s = 1 \dots p - 1. \tag{37}$$

#### 4.6. End of the proof

The previous analysis shows that when  $q$  is prime, the factors  $x_i$  (respectively  $X_i$ )  $1 \leq i < p$  appear with the same exponent. In other words, for  $q$  a prime number, the points  $P_1, P_2, \dots, P_{p-1}$  play an equivalent role; they will have the *same multiplicities* on each  $S_n$ ; and we will use  $u_n^1$  to denote their common value.

Using (36), (37) together with (28) and (30) we get

$$\begin{aligned} d_n &= (p - 1)d_{n-1} - u_{n-1}^0 - (p - 1)u_{n-1}^1, \\ u_n^0 &= (p - 2)d_{n-2} - (p - 1)u_{n-2}^1, \\ u_n^1 &= (p - 2)d_{n-4} - u_{n-4}^0 - (p - 2)u_{n-4}^1. \end{aligned} \tag{38}$$

The rate of growth of the  $d_n$ 's is the inverse of the modulus of the smallest eigenvalues of the  $12 \times 12$  matrix given by the above linear system. The outcome is that the rate of growth of  $K$  is the inverse of the smaller root of

$$x^2 + (2 - (p - 1)^2)x + 1 = 0. \tag{39}$$

To obtain the full expression of the generating functions, we need to specify the initial values of  $d_n, u_n^0$  and  $u_n^1$ . They can easily be calculated with the help of formal calculation software. The results are summarized in table 2.

Note that when  $q$  is not prime, we may still write a set of recursions similar to (38). The system is not complete, and cannot be obtained from the analysis presented in section 4 (see appendix A).

**Table 2.** Initial values of  $d_n$ ,  $u_n^0$  and  $u_n^1$  for  $0 \leq n < 4$ .

| $n$ | $d_n$                         | $u_n^0$            | $u_n^1$ |
|-----|-------------------------------|--------------------|---------|
| 0   | 1                             | 0                  | 0       |
| 1   | $p - 1$                       | 0                  | 0       |
| 2   | $(p - 1)^2$                   | $p - 2$            | 0       |
| 3   | $p^3 - 3p^2 + 2p + 1$         | $(p - 1)(p - 2)$   | 0       |
| 4   | $(p - 1)(p^3 - 3p^2 + p + 3)$ | $(p - 1)^2(p - 2)$ | $p - 2$ |

## 5. Arithmetical approach: complexity through number of digits

The third approach consists in calculating the image of integer points, and evaluating the growth of the size of the coordinates, through the number of digits. It means that we do not try to calculate the iterates formally. This method was already experimented in [13].

Obviously the integer coordinates become extremely large, as large as  $10^{6000}$  and we used the GMP library to implement the program [27]. At each step of the calculation we factor out the greatest common divisor of the components. We assume that the existence of a common factor between all the coordinates is due to a factorization of the underlying polynomial. This assumption is valid, at least after the first step where an accidental factorization could occur. The degree of the polynomial is estimated as the number of bits used to store a typical entry (i.e.,  $\log_2(M_{ij})$ ). The algorithm proceeds as follows: (i) construct a random matrix of integers respecting the equalities of the pattern under consideration, (ii) replace each term by its cofactor, (iii) divide every term by the greatest common factor of all of them, (iv) replace each term by the product of all others, (v) divide every term by the greatest common factor of all of them, (vi) record the number of digits used to store the matrix elements. Note that one can exchange (ii) and (iv) without altering the results. The procedure is iterated for as many steps as possible, and possibly several runs with different initial matrices are performed. Note that for pattern involving only very few variables, it can be efficient to write directly the recursion relation over the variables.

The results are summarized in table 3, giving the value of the complexity for various values of  $q$  and for the four patterns introduced above. For cyclic matrices and general  $q$  it has been shown in [15] that the rate of growth for  $K = I \circ J$  is a quadratic integer which is the inverse of the smaller root of

$$x^2 + (2 - (q - 2)^2)x + 1. \quad (40)$$

In table 3 an empty cell means that we have not been able to compute the corresponding  $\lambda$ . This is due to the fast growth of the coordinates, preventing us to perform a sufficient number of numerical iterations. The number of digits displayed is just an indication of the estimated accuracy of our numerical result.

## 6. Conclusion

The three different approaches we have used give corroborating results. The first two methods are indeed measuring the same object, i.e. the rate of growth of the degree of iterates. The third method is evaluating something which may *a priori* be different, but appears to coincide with the previous one. We see by comparing the last two columns of table 3 that  $\lambda_G$  happens to be extremely close to  $\lambda_S$ , as well as to  $\lambda_C$ . This allows us to state the two conjectures:

**Table 3.** Complexities of  $K = I \circ J$  for various values of  $q$ , for patterns (CS), (C), (S) and (G). The numerical and analytical results are displayed. The number in brackets is the number of iterations of  $K$  we have been able to calculate.

| $q$ |            | Cyclic symmetric<br>$\lambda_{CS}$ | Cyclic<br>$\lambda_C$ | Symmetric<br>$\lambda_S$ | General<br>$\lambda_G$ |
|-----|------------|------------------------------------|-----------------------|--------------------------|------------------------|
| 5   | Numerical  | 1.000                              | 6.85 [7]              | 6.86 [7]                 | 6.86 [6]               |
|     | Analytical | 1                                  | 6.854                 |                          |                        |
| 6   | Numerical  | 4.00 [10]                          | 13.93 [5]             | 13.88 [5]                | 13.9 [4]               |
|     | Analytical | 4                                  | 13.928                |                          |                        |
| 7   | Numerical  | 6.854 [7]                          | 22.96 [4]             | 22.97 [4]                | 22.97 [4]              |
|     | Analytical | 6.854                              | 22.956                |                          |                        |
| 8   | Numerical  | 10.33 [6]                          | 33.97 [4]             | 33.97 [3]                | 34.1 [3]               |
|     | Analytical |                                    | 33.970                |                          |                        |
| 9   | Numerical  | 12.83 [5]                          | 47.0 [3]              | 47.0 [3]                 | 47.0 [3]               |
|     | Analytical |                                    | 46.978                |                          |                        |
| 10  | Numerical  | 17.9 [4]                           | 62.0 [3]              | 62.0 [3]                 | 62.0 [2]               |
|     | Analytical |                                    | 61.984                |                          |                        |
| 11  | Numerical  | 22.96 [4]                          | 79.0 [3]              | 79.1 [2]                 | 80.7 [2]               |
|     | Analytical | 22.956                             | 78.987                |                          |                        |
| 12  | Numerical  | 25.8 [4]                           | 98.0 [3]              | 99.1 [2]                 | 100.3 [2]              |
|     | Analytical |                                    | 97.990                |                          |                        |
| 13  | Numerical  | 33.97 [3]                          | 130.3 [3]             | 121.6 [2]                | 121.5 [1]              |
|     | Analytical | 33.970                             | 118.992               |                          |                        |
| 14  | Numerical  | 39.1 [2]                           | 142.8 [2]             | 144.5 [2]                | 144.2 [1]              |
|     | Analytical |                                    | 141.993               |                          |                        |
| 15  | Numerical  | 42.19 [2]                          | 167. [2]              | 170. [2]                 |                        |
|     | Analytical |                                    | 166.99                |                          |                        |
| 16  | Numerical  | 49.10 [2]                          | 194. [2]              |                          |                        |
|     | Analytical |                                    | 193.995               |                          |                        |
| 17  | Numerical  | 61.6 [2]                           | 224. [2]              |                          |                        |
|     | Analytical | 61.984                             | 222.995               |                          |                        |

**Proposition 1.** *Evaluating the rate of growth of the ‘size’ of iterates on rational points yields the same value as the rate of growth of the degree of the iterates.*

**Proposition 2.** *The rate of growth of the degree of the iterates of the transformation  $K = I \circ J$  for the general matrices (pattern (G)), for symmetric matrices (pattern (S)), and for cyclic matrices (pattern (C)) are the same. Their common value is the inverse of the modulus of the smaller root of  $x^2 - (q^2 - 4q + 2)x + 1 = 0$ .*

Such a statement means that although the number of parameters of patterns (G) and (S) is much bigger than the one of pattern (C), the latter captures the entirety of the complexity of the product of inversions  $K = I \circ J$ . This might be related to the structure of bialgebra of the set of square matrices equipped with ordinary matrix product and Hadamard product. Phrased differently, the skeleton formed by the cyclic matrices encodes the structure of the whole bialgebra. This deserves further investigations which are beyond the scope of this paper.

**Appendix. The cyclic symmetric case for  $q = 9$**

We consider in this appendix the case  $q = 9$ . Since  $q$  is not a prime number, our result of section 4 does not apply.

**Table A1.** The initial values of  $d_n, u_n^0, u_n^1$  and  $u_n^2$  for  $0 \leq n \leq 4$ .

| $n$ | $d_n$ | $u_n^0$ | $u_n^1$ | $u_n^2$ |
|-----|-------|---------|---------|---------|
| 0   | 1     | 0       | 0       | 0       |
| 1   | 4     | 0       | 0       | 0       |
| 2   | 16    | 3       | 0       | 2       |
| 3   | 59    | 12      | 0       | 8       |
| 4   | 216   | 46      | 3       | 32      |

The number of homogeneous variables is  $p = (q + 1)/2 = 5$ . We use the same notation as in the text for the hyperplane  $\Pi_k$  and the point  $P_i$ . In addition we define the three points  $Q_1 = (1, 1, -1, -1, 1)$ ,  $Q_2 = (1, 1, 1, -1, -1)$  and  $Q_4 = (1, -1, 1, -1, 1)$ . We also introduce the codimension-two variety  $\Pi_{0,3}$  defined by the two equations  $x_0 = 0$  and  $x_3 = 0$ . The singularity structure is

$$\begin{aligned} \Pi_0 &\xrightarrow{J} P_0 \xrightarrow{I} P_0 \rightsquigarrow \Pi_0 \\ \Pi_1 &\xrightarrow{J} P_1 \xrightarrow{I} Q_1 \xrightarrow{J} Q_1 \xrightarrow{I} P_1 \rightsquigarrow \Pi_1 \\ \Pi_2 &\xrightarrow{J} P_2 \xrightarrow{I} Q_2 \xrightarrow{J} Q_2 \xrightarrow{I} P_2 \rightsquigarrow \Pi_2 \\ \Pi_3 &\xrightarrow{J} P_3 \xrightarrow{I} Q_3 \rightsquigarrow \Pi_{0,3} \\ \Pi_4 &\xrightarrow{J} P_4 \xrightarrow{I} Q_4 \xrightarrow{J} Q_4 \xrightarrow{I} P_4 \rightsquigarrow \Pi_4 \end{aligned}$$

the subsequent iterates of  $\Pi_{0,3}$  are non-singular. We see that there will be six sets of exponents:  $u_n^0$  and  $v_n^0$  related to  $x_0$ ;  $u_n^1$  and  $v_n^1$  related to  $x_1, x_2$  and  $x_4$ ; and finally  $u_n^2$  and  $v_n^2$  related to  $x_3$ . The equations expressing the degree drop due to the factorization, and the fact that  $I$  and  $J$  are involutions, are

$$\begin{aligned} d_{n+1} &= 4d_n - u_n^0 - 3u_n^1 - u_n^2, \\ v_{n+1}^0 &= 3d_n - 3u_n^1 - u_n^2, \\ v_{n+1}^1 &= 3d_n - u_n^0 - 2u_n^1 - u_n^2, \\ v_{n+1}^2 &= 3d_n - u_n^0 - 3u_n^1. \end{aligned}$$

Moreover, the singularity structure shown above yields

$$u_{n+1}^0 = v_n^0 \quad u_{n+1}^1 = v_{n-2}^1.$$

It is clear that an equation is missing to close the system:

$$\begin{aligned} d_{n+1} &= 4d_n - u_n^0 - 3u_n^1 - u_n^2 \\ u_{n+1}^0 &= 3d_{n-1} - 3u_{n-1}^1 - u_{n-1}^2 \\ u_{n+1}^1 &= 3d_{n-3} - u_{n-3}^0 - 2u_{n-3}^1 - u_{n-3}^2. \end{aligned}$$

If we suppose that there exists a recursion relation of the form

$$u_{n+1}^2 = ad_{n-q} + bu_{n-q}^0 + cu_{n-q}^1 + du_{n-q}^2 + e,$$

where the  $a, b, c, d, e$ , as well as the shift  $q$  are integer constants. The hypothesis  $q = 1$  yields

$$u_{n+1}^2 = 2d_{n-1} - 3u_{n-1}^1. \tag{A.1}$$

Introducing, with obvious notations, the generating functions

$$d(s) = \sum d_n s^n, \quad u_i(s) = \sum u_n^i s^n, \quad i = 1, 2, 3,$$

one easily finds (see table A1)

$$d(s) = 1 + \frac{(4 - s^2 - s^6)s}{P(s)}, \quad u_0(s) = \frac{(2s^2 - 3)(1 + s^2)s^4}{P(s)},$$

$$u_1(s) = \frac{(2s^2 - 3)s^4}{P(s)}, \quad u_2(s) = \frac{(3s^4 - 2s^2 - 2)s^2}{P(s)},$$

with

$$P(s) = (1 - s) \cdot (1 - 3s - 2s^2 - s^3 + 2s^4 + 2s^5 - s^6)$$

from which

$$f_9(u) = \frac{(1 + u + 3u^2 - 3u^3)^2}{(1 - u)(1 - 13u + 2u^2 + u^3 + 12u^4 - 8u^5 + u^6)}.$$

## References

- [1] Bellon M P, Maillard J-M and Viallet C-M 1991 Integrable Coxeter groups *Phys. Lett. A* **159** 221–32
- [2] Bellon M P, Maillard J-M and Viallet C-M 1991 Infinite discrete symmetry group for the Yang-Baxter equations: spin models *Phys. Lett. A* **157** 343–53
- [3] Jaeger F 1992 Strongly regular graphs and spin models for the Kauffman polynomial *Geom. Ded.* **44** 23–52
- [4] Jaeger F 1994 Classification of small spin models *Kyushu J. Math* **48** 185–200
- [5] de la Harpe P 1994 Spin models for link polynomials, strongly regular graphs and Jaeger’s Higman-Sims model *Pac. J. Math.* **162** 57–96
- [6] Jones V F R 1999 *Planar Algebras (Preprint math.QA/9909027)*
- [7] Kauffman L H 1999 Combinatorics and topology—François Jaeger’s work in knot theory *Ann. Inst. Fourier* **49** 927–53
- [8] Matsumoto M 1999 A generalization of Jaeger-Nomura’s Bose Mesner algebra associated to type II matrices *Ann. Inst. Fourier* **49** 1027–35
- [9] Arnold V I 1990 Dynamics of complexity of intersections *Bol. Soc. Bras. Mat.* **21** 1–10
- [10] Bellon M P, Maillard J-M, Rollet G and Viallet C-M 1992 Deformations of dynamics associated to the chiral Potts model *Int. J. Mod. Phys. B* **6** 3575–84
- [11] Falqui G and Viallet C-M 1993 Singularity, complexity, and quasi-integrability of rational mappings *Commun. Math. Phys.* **154** 111–25
- [12] Abarenkova N, Anglès d’Auriac J-C, Boukraa S, Hassani S and Maillard J-M 1999 Rational dynamical zeta function for birational transformations *Physica A* **264** 264–93
- [13] Abarenkova N, Anglès d’Auriac J-C, Boukraa S and Maillard J-M 1999 Growth-complexity spectrum of some discrete dynamical systems *Physica D* **130** 27–42
- [14] Russakovskii A and Shiffman B 1997 Value distribution of sequences of rational mappings and complex dynamics *Indiana Univ. Math. J.* **46** 897–932
- [15] Bellon M and Viallet C-M 1999 Algebraic entropy *Commun. Math. Phys.* **204** 425–37
- [16] Boninfant A M and Fornaess J E 2000 Growth of degree for iterates of rational maps in several variables *Indiana Univ. Math. J.* **49** 751–78
- [17] Takenawa T 2001 Discrete dynamical systems associated with root systems of indefinite type *Commun. Math. Phys.* **224** 657–81
- [18] Takenawa T 2001 A geometric approach to singularity confinement and algebraic entropy *J. Phys. A: Math. Gen.* **34** L95–L102
- [19] Diller J and Favre C 2001 Dynamics of bimeromorphic maps of surfaces *Am. J. Math.* **123** 1135–69
- [20] Cantat S and Favre C 2003 Symétries birationnelles des surfaces feuilletées *J. Reine Angew. Math.* **561** 199–235 (Preprint [math.CV/0206209](https://arxiv.org/abs/math.CV/0206209))
- [21] Bedford E and Kim K 2004 On the degree growth of birational mappings in higher dimension *J. Geom. Anal.* **14** 567–96 (Preprint [math.DS/0406621](https://arxiv.org/abs/math.DS/0406621))
- [22] Hasselblatt B and Propp J 2005 *Monomial maps and algebraic entropy Ergodic Theor. Dyn. Syst.* submitted
- [23] Anglès d’Auriac J-C, Maillard J-M and Viallet C-M 2002 A classification of four-state spin edge Potts models *J. Phys. A: Math. Gen.* **35** 9251–72 (Preprint [cond-mat/0209557](https://arxiv.org/abs/cond-mat/0209557))

- 
- [24] Korepanov I 2003 Exact solution for a matrix dynamical system with usual and Hadamard inverses *Preprint nlin.SI/0303007*
  - [25] Bellon M P, Maillard J-M and Viallet C-M 1991 Rational mappings, arborescent iterations, and the symmetries of integrability *Phys. Rev. Lett.* **67** 1373–6
  - [26] Bellon M P, Maillard J-M and Viallet C-M 1991 Higher dimensional mappings *Phys. Lett. A* **159** 233–44
  - [27] Gnu MultiPrecision, Arbitrary precision arithmetics software, see <http://www.swox.com/gmp/>