

RATIONAL CURVES ON HYPERSURFACES OF LOW DEGREE II

JOE HARRIS AND JASON STARR

ABSTRACT. This is the second in a sequence of papers on the geometry of spaces of rational curves of degree e on a general hypersurface $X \subset \mathbb{P}^n$ of degree d . In [11] it is proved that if $d < \frac{n+1}{2}$ then for each e the space of rational curves is irreducible, reduced and has the expected dimension. In this paper it is proved that if $d^2 + d + 1 \leq n$, then for each e the space of rational curves is a *rationally connected* variety; in particular it has negative Kodaira dimension.

CONTENTS

1. Statement of results	1
2. Deformation ample	5
3. Deformation theory of stable maps	11
4. Conditions on families of stable maps	26
5. The induction argument	32
6. Twistable lines on hypersurfaces	43
7. Base case of the induction for hypersurfaces	47
8. Proof of the main theorem	57
References	57

1. STATEMENT OF RESULTS

This is the second in a sequence of papers on the geometry of spaces of rational curves on a general hypersurface $X \subset \mathbb{P}^n$. Let $d = \deg(X)$. In [11] it is proved that if $d < \frac{n+1}{2}$ then for each e the space of rational curves of degree e is irreducible, reduced and has the expected dimension. The main result of this paper is the following theorem.

Theorem 1.1. *If $X \subset \mathbb{P}^n$ is a general hypersurface of degree d and if $n \geq d^2 + d + 1$ then for each integer $e \geq 1$ the stack $\overline{\mathcal{M}}_{0,0}(X, e)$ is rationally connected. More precisely, there exists a morphism $f : \mathbb{P}^1 \rightarrow ((\overline{\mathcal{M}}_{0,0}(X, e))_{\text{fine}})_{\text{sm}}$ such that $f^*T_{\overline{\mathcal{M}}_{0,0}(X, e)}$ is ample.*

Remark 1.2. (i) The scheme $\overline{\mathcal{M}}_{0,0}(X, e)$ is the coarse moduli space of $\overline{\mathcal{M}}_{0,0}(X, e)$, the open subset $(\overline{\mathcal{M}}_{0,0}(X, e))_{\text{fine}}$ is the fine moduli locus, and the open subset $((\overline{\mathcal{M}}_{0,0}(X, e))_{\text{fine}})_{\text{sm}}$ is the smooth locus of the fine moduli locus.

(ii) For the cases $d = 1, 2$, a related and stronger theorem is proved in [15, Thm. 3]; namely the coarse moduli space $\overline{\mathcal{M}}_{0,0}(X, e)$ is *rational*. The proof relies on the fact that X is a homogeneous space and does not extend to the case $d \geq 3$. Moreover, it is not clear from [15] that there exists a very free rational curve in the smooth locus of the fine moduli locus of the coarse moduli space.

The space $\overline{\mathcal{M}}_{0,0}(X, e)$ is the *Kontsevich moduli stack of stable maps*, which will be recalled below; it is a Deligne-Mumford stack containing the parameter space of smooth rational curves in X of degree e as an open substack. A variety is *rationally connected* if any two closed points are contained in the image of a morphism from \mathbb{P}^1 to the variety. Rationally connected varieties have negative Kodaira dimension, hence the schemes $\overline{\mathcal{M}}_{0,0}(X, e)$ have negative Kodaira dimension.

The motivation behind Theorem 1.1 is a conjectural relationship between rational connectedness of $\overline{\mathcal{M}}_{0,0}(X, e)$ and a theorem of Lang about rational points of varieties defined over the function field of a surface.

Theorem 1.3 (Lang, [18]). *Let K be the function field of a surface over \mathbb{C} and let $X \subset \mathbb{P}_K^n$ be a hypersurface of degree d . If $d^2 \leq n$, then $X(K) \neq \emptyset$.*

There is a naive parameter count that suggests that if $d^2 \leq n$ then for $e \gg 0$ the stack $\overline{\mathcal{M}}_{0,0}(X, e)$ is rationally connected, and if $d^2 \geq n + 2$ then for $e \gg 0$ the stack $\overline{\mathcal{M}}_{0,0}(X, e)$ is of general type; i.e., the Kodaira dimension of $\overline{\mathcal{M}}_{0,0}(X, e)$ is determined by the same inequality as in Lang's theorem. This suggests Lang's theorem is related to rational connectedness of spaces of rational curves. In a personal communication, A. J. de Jong has outlined an approach for proving that a K -variety X has a K -point if the stacks $\overline{\mathcal{M}}_{0,0}(X \otimes_K \overline{K}, e)$ are rationally connected, and a certain *Brauer obstruction* vanishes. Of course Lang's original proof is simple and direct. But de Jong's approach could apply to classes of varieties where Lang's proof does not apply, i.e., to varieties that are not hypersurfaces in projective space.

The naive parameter count is not rigorous. As it seems impossible to make it rigorous, it is not recalled here. In this paper a different strategy is developed for proving the stacks $\overline{\mathcal{M}}_{0,0}(X, e)$ are rationally connected, and this strategy is applied to hypersurfaces in projective space. This strategy should also apply to other varieties, hence it is formulated in greater generality than strictly needed for the case of hypersurfaces.

1.1. The Kontsevich moduli space. The most natural parameter space for rational curves of degree e on X is the open subscheme of the Hilbert scheme parametrizing smooth rational curves of degree e on X , $R_X^e \subset \text{Hilb}_X^{et+1}$. For $e > 1$, R_X^e is not proper. To study the global geometry of R_X^e , e.g. to determine its Kodaira dimension, it is necessary to embed it as an open subset of a proper scheme. The simplest choice is to take the closure $\overline{R_X^e}$ inside Hilb_X^{et+1} . This is a poor choice for two reasons: First, there is no simple characterization of the closed subschemes of X that correspond to points in $\overline{R_X^e}$. Second, the deformation theory of a closed subscheme of X is difficult to work with.

There is a better choice; one where the points of the closure have a simple geometric meaning, and where the deformation theory is easier to work with. This choice is $\overline{\mathcal{M}}_{0,0}(X, e)$, or more generally $\overline{\mathcal{M}}_{0,r}(X, e)$, the *Kontsevich moduli space of degree e , r -pointed, genus 0 stable maps to X* . This space has one disadvantage over $\overline{R_X^e}$; namely $\overline{\mathcal{M}}_{0,r}(X, e)$ is a *Deligne-Mumford stack* rather than a scheme. However, the coarse moduli space $\overline{M}_{0,r}(X, e)$ is a projective scheme, cf. [2]. For the reader unfamiliar with stacks, most occurrences of $\overline{\mathcal{M}}_{0,r}(X, e)$ can safely be replaced by $(\overline{M}_{0,r}(X, e))_{\text{fine}}$, the *fine moduli locus* of $\overline{M}_{0,r}(X, e)$.

To be precise, $\overline{\mathcal{M}}_{0,r}(X, e)$ is the stack whose objects are triples

$$\zeta = ((p : \Sigma \rightarrow B, \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X),$$

consisting of a flat, proper family of curves $p : \Sigma \rightarrow B$, a collection of r disjoint sections $\sigma_i : B \rightarrow \Sigma, i = 1, \dots, r$ with image in the smooth locus of p , and a morphism $g : \Sigma \rightarrow X$ such that for each

geometric point $b \in B$, the fiber Σ_b is a connected, at-worst-nodal curve of arithmetic genus 0, the morphism $g_b : \Sigma_b \rightarrow X \otimes_{\mathbb{C}} \kappa(b)$ has no infinitesimal automorphism fixing the marked points $\sigma_i(b)$, and the degree of $(g_b)^* \mathcal{O}_X(1)$ is e , cf. [9], [5].

A refinement of this stack used in this paper is the *Behrend-Manin stack*, $\overline{\mathcal{M}}(X, \tau)$, associated to a genus 0 *stable A-graph* τ , cf. [5]. A genus 0 stable *A-graph* is a *tree with tails* with a degree associated to each vertex that satisfies a certain stability condition. A tree is a graph that contains no cycles. A tail, or a *half-edge*, is an edge that originates on a vertex but does not terminate on a vertex (e.g. what one would get if one were to “cut in half” an edge of a usual graph). A *degree function* is an assignment of a nonnegative integer to each vertex of the graph. And the *stability condition* is that no vertex both has degree 0 and valence less than 3 (tails count toward the valence of a vertex).

Given an at-worst-nodal curve of genus 0, Σ , a collection of marked points on Σ , $\sigma_1, \dots, \sigma_r$, and a morphism $g : \Sigma \rightarrow X$, there is an associated *A-graph* defined as follows. The tree is the dual graph of Σ ; there is one vertex for each irreducible component of Σ and one edge for each node of Σ . For each marked point σ_i of Σ there is a tail attached to the obvious vertex. And the degree of a vertex is simply the degree of $g^* \mathcal{O}(1)$ on the corresponding irreducible component. The Behrend-Manin stack $\overline{\mathcal{M}}(X, \tau)$ (essentially) parametrizes the closure of the locally closed substack of $\overline{\mathcal{M}}_{0,r}(X, e)$ of stable maps whose associated *A-graph* equals τ . For the precise definition, cf. [5].

The boundary of the stack $\overline{\mathcal{M}}(X, \tau)$ is a union of stacks $\overline{\mathcal{M}}(X, \sigma)$ where σ ranges over stable *A-graphs* such that there is a *contraction* from σ to τ , i.e., the graph σ is “more degenerate” than the graph τ , cf. [5]. Hence the boundary of every Behrend-Manin stack can be understood inductively starting from the “most degenerate” graphs τ , for which $\overline{\mathcal{M}}(X, \tau)$ is a fiber product over X of the space of pointed lines on X . So questions about $\overline{\mathcal{M}}(X, \tau)$ that can be studied by specializing to points in the boundary eventually reduce to questions about the space of pointed lines on X . Moreover, the deformation theory of a point in $\overline{\mathcal{M}}(X, \tau)$ is straightforward; it will be recalled in Section 3.

1.2. Sketch of the proof. The proof of Theorem 1.1 uses a theorem of Kollár.

Theorem 1.4 (Kollár, Thm. IV.3.7 [16]). *Let V be an irreducible, projective variety, and let $V_{sm} \subset V$ denote the smooth locus of V . If there exists a very free morphism $f : \mathbb{P}^1 \rightarrow V_{sm}$, i.e. a morphism such that $f^* T_V$ is an ample vector bundle, then V is rationally connected.*

The reader is warned that “very free” is the first of a multitude of definitions with “very” similar names: free, very free, deformation ample, very stable, unobstructed, twisting, very twisting, twistable, very twistable, positive, very positive, inducting pair, inductable, modification, typical, and c -generating linear system. A morphism $f : \mathbb{P}^1 \rightarrow V$ is *free* if $f^* T_V$ is a vector bundle that is generated by global sections. The goal is to prove there exists a very free morphism to $\overline{\mathcal{M}}_{0,0}(X, e)$ for all e . It is difficult to directly construct a very free morphism. However, existence of a very free morphism can be studied by specializing to the boundary of $\overline{\mathcal{M}}_{0,0}(X, e)$, and, using an induction argument, can ultimately be reduced to a question about pointed lines on X . The induction argument does not produce a very free morphism to the boundary of $\overline{\mathcal{M}}_{0,0}(X, e)$, but rather a reducible rational curve B in the boundary such that the restriction of $T_{\overline{\mathcal{M}}_{0,0}(X, e)}$ to B is *deformation ample*: for a deformation of B to an irreducible rational curve B' , the restriction of $T_{\overline{\mathcal{M}}_{0,0}(X, e)}$ to B' is *ample*, cf. Section 2 (a curve B' is called a *deformation* of B if both B and B' have a common generalization). The rational curve B' is a very free rational curve which proves that $\overline{\mathcal{M}}_{0,0}(X, e)$ is rationally connected.

The induction argument constructs a reducible rational curve B which itself parametrizes stable maps from reducible rational curves to X . Each of these stable maps is the union of a map of degree $e - 1$, $g : \Sigma \rightarrow X$, and a line $L \subset X$ that intersect in a node $\sigma \in \Sigma \cap L$. Since the union is stable, the 1-pointed map (Σ, σ, g) is stable. To simplify the deformation theory, it is assumed that (Σ, σ, g) is *very stable*: the unmarked map (Σ, g) is stable (this assumption must be justified!). To guarantee that B is in the smooth locus of $\overline{\mathcal{M}}_{0,0}(X, e)$, it is assumed that (Σ, g) is *unobstructed*: $\mathbb{E}xt_{\mathcal{O}_\Sigma}^1(L_g, \mathcal{O}_\Sigma) = \{0\}$.

The condition that $T_{\overline{\mathcal{M}}_{0,0}(X,e)}|_B$ is deformation ample can be translated into a condition on the family of pointed lines, (L, σ) , together with a condition on the family of pointed maps of degree $e - 1$, (Σ, σ, g) ; these conditions are defined in Section 4. The condition on the family of pointed lines is that it is *very twisting*. The family is *twisting* (resp. *very twisting*) if the associated morphism $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ pulls back the vertical tangent bundle of the projection $\text{pr} : \overline{\mathcal{M}}_{0,1}(X, 1) \rightarrow \overline{\mathcal{M}}_{0,0}(X, 1)$ to a bundle which is generated by global sections, and pulls back the vertical tangent bundle of the “evaluation at the marked point” morphism $\text{ev} : \overline{\mathcal{M}}_{0,1}(X, 1) \rightarrow X$ to a bundle which is generated by global sections (resp. deformation ample). Composing ζ with ev gives a map $h : B \rightarrow X$. Such a map which arises from a twisting (resp. very twisting) family is called *twistable* (resp. *very twistable*). The property of being twistable (resp. very twistable) is an open condition on the family of all maps.

The condition on the family of pointed maps of degree $e - 1$ is that it is very positive. The family is *positive* (resp. *very positive*) if the associated morphism $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e - 1)$ pulls back the vertical tangent bundle of the projection $\text{pr} : \overline{\mathcal{M}}_{0,1}(X, 1) \rightarrow \overline{\mathcal{M}}_{0,0}(X, 1)$ to a bundle which is generated by global sections (resp. ample) and the pullback by $\text{pr} \circ \zeta$ of $T_{\overline{\mathcal{M}}_{0,0}(X,1)}$ is deformation ample. The main observation is this: For a positive (resp. very positive) family over a smooth rational curve, the morphism $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e - 1)$ is free (resp. very free).

An *inducting pair* consists of a very twisting family of pointed lines over B and a very positive family of pointed maps of degree $e - 1$ over B which intersect along the marked points. The induction step proves that if an inducting pair exists for degree e , then an inducting pair exists for degree $e + 1$. It is sometimes useful to “forget” the very twisting family of pointed lines, and only “remember” that the family of marked points is a very twistable map. A very positive family of pointed maps of degree $e - 1$ such that the family of marked points is a very twistable map is an *inductable* family.

The induction step begins with an inducting pair for degree e and produces an inductable family for degree $e + 1$. In particular, the inductable family is very positive and there is a deformation to a very free morphism from an irreducible curve to $\overline{\mathcal{M}}_{0,1}(X, e)$ whose projection to $\overline{\mathcal{M}}_{0,0}(X, e)$ is a very free morphism. The family of unmarked maps over B is obtained by gluing the very twisting family of lines and the very positive family of maps of degree $e - 1$ along the curve of marked points σ . To get a family of marked maps, let σ' be an irreducible curve in the total space Σ of the very positive family which is linearly equivalent to σ . The curve σ' fails to be a family of marked points precisely at the finitely many points in B over which the curve σ' intersects the curve σ . The solution is to blow up each of these finitely many points on Σ ; this produces a family of marked stable maps of degree e . Unfortunately blowing up destroys the “very positivity” of the family. Very positivity is restored by making a *modification* at each of the finitely many points of B over which Σ is blown up. The modification attaches a \mathbb{P}^1 to B at the specified point and extends the family over this \mathbb{P}^1 so that the stabilized family of unmarked maps is a constant family, but such that the

marked points in this constant family vary. After modification, the family of marked stable maps is inductable. Incidentally, it is the process of modification that requires using reducible curves in $\overline{\mathcal{M}}_{0,0}(X, e)$ and working with the property of deformation ample bundles.

The last section establishes the base case of the induction argument: existence of a family of pointed lines that is both very twisting and very positive. The total space of a very positive family of lines is a scroll Σ . The whole argument reduces to producing a pair of a scroll Σ and a hypersurface X containing Σ such that Σ corresponds to a very positive family of lines on X . This reduces to a computation of the dimension of certain linear systems on the scroll Σ , *c-generating linear systems*. This is a straightforward computation in the Cox homogeneous coordinate ring of the scroll. The hypothesis that $d^2 + d + 1 \leq n$ is used in this last computation.

1.3. Technical Hypotheses. In the proof of the induction step, some technical hypotheses are used.

Hypothesis 1.5. For each contraction of genus 0 stable A -graphs, $\phi : \sigma \rightarrow \tau$, the image of the morphism of Behrend-Manin stacks $\overline{\mathcal{M}}(X, \sigma) \rightarrow \overline{\mathcal{M}}(X, \tau)$ has codimension $\dim(X, \tau) - \dim(X, \sigma)$ in $\overline{\mathcal{M}}(X, \tau)$.

By [11, Prop. 7.4], if $d < \frac{n+1}{2}$ and if $X \subset \mathbb{P}^n$ is a general hypersurface of degree d , each stack $\overline{\mathcal{M}}(X, \sigma)$ has the expected dimension; thus Hypothesis 1.5 holds for X .

Hypothesis 1.6. A general fiber of the evaluation map $\text{ev} : \overline{\mathcal{M}}_{0,1}(X, 1) \rightarrow X$ is irreducible.

For a pair (X, p) consisting of a hypersurface $X \subset \mathbb{P}^n$ of degree d and a point $p \in X$, the associated fiber of ev is a subvariety $Z \subset \mathbb{P}^{n-1}$ which is a complete intersection of a sequence of hypersurfaces Y_1, \dots, Y_d in \mathbb{P}^n with $\deg(Y_i) = i$: the defining equation of Y_i is the degree i homogeneous part of the Taylor expansion of the defining equation of X about the point p . If the pair (X, p) is general, the sequence of hypersurfaces Y_1, \dots, Y_d is general. By the Bertini theorem [14, Thms. 4.10, 6.10] the intersection $Y_1 \cap \dots \cap Y_d$ is smooth and connected if $d \leq n - 2$.

Hypothesis 1.7. For each integer $e \geq 0$, the locus in $\overline{\mathcal{M}}_{0,1}(X, e)$ parametrizing stable maps with nontrivial automorphism group has codimension at least 2.

Of course any stable map with nontrivial automorphism group has an irreducible component which is a multiple cover of its image. In light of [11, Prop. 7.4], a simple parameter count shows that if $d \leq \frac{n+1}{2}$ and if $X \subset \mathbb{P}^n$ is a general hypersurface of degree d , Hypothesis 1.7 is satisfied.

1.4. Conventions. Unless stated otherwise, schemes are of finite type and separated over $\text{Spec } \mathbb{C}$. Absolute fiber products of schemes will be fiber products over $\text{Spec } \mathbb{C}$. Absolute fiber products of stacks will be 2-fibered products over $\text{Spec } \mathbb{C}$.

1.5. Acknowledgments. We are very grateful to A. J. de Jong and Steven Kleiman for many useful conversations. We are also grateful to the referees for their many useful comments.

2. DEFORMATION AMPLE

Let T be a scheme.

Definition 2.1. A family of prestable curves of genus g over T is a proper, flat morphism $\pi : B \rightarrow T$ such that every geometric fiber of π is a connected, at-worst-nodal curve of arithmetic genus g .

Notation 2.2. Let $\pi : B \rightarrow T$ be a morphism of schemes and let E be a quasi-coherent sheaf on B . Denote by $\mu_E : \pi^* \pi_* E \rightarrow E$ the morphism left adjoint to the identity morphism $\pi_* E \rightarrow \pi_* E$. The sheaf E is π -relatively generated by global sections if μ_E is surjective.

Let $\pi : B \rightarrow T$ be a family of prestable curves of genus 0 and let E be a coherent sheaf on B . Let $\sigma : T \rightarrow B$ be a section of π and let $\mathcal{I} \subset \mathcal{O}_B$ be the ideal sheaf of $\sigma(T)$.

Lemma 2.3.

- (i) If E is π -relatively generated by global sections, then $R^1\pi_*E = \{0\}$.
- (ii) If E is π -relatively generated by global sections, then $R^1\pi_*(\mathcal{I} \cdot E) = \{0\}$.
- (iii) Let E' and E'' be coherent sheaves on B and let

$$0 \longrightarrow E' \longrightarrow E \longrightarrow E'' \longrightarrow 0$$

be a short exact sequence of coherent sheaves. If E' and E'' are π -relatively generated by global sections, then E is π -relatively generated by global sections.

Proof. All three statements can be proved locally over T . Hence it suffices to consider the case that T is affine.

(i): Since E is π -relatively generated by global sections and T is affine, E is generated by global sections, i.e., there is a short exact sequence of coherent sheaves,

$$0 \longrightarrow K \longrightarrow \mathcal{O}_B^{\oplus N} \longrightarrow E \longrightarrow 0.$$

Since π is of relative dimension 1, $R^2\pi_*K = \{0\}$. Since the fibers of B are connected of arithmetic genus 0, $R^1\pi_*\mathcal{O}_B = \{0\}$. In the long exact sequence of higher direct images associated to the short exact sequence above, $R^1\pi_*E$ fits between $R^1\pi_*\mathcal{O}_B^{\oplus N}$ and $R^2\pi_*K$; hence $R^1\pi_*E = \{0\}$.

(ii): There is a short exact sequence of coherent sheaves,

$$0 \longrightarrow \mathcal{I} \cdot E \longrightarrow E \longrightarrow E \otimes_{\mathcal{O}_B} \mathcal{O}_{\sigma(T)} \longrightarrow 0$$

giving rise to a long exact sequence of cohomology groups,

$$\pi_*E \longrightarrow \pi_*(E \otimes_{\mathcal{O}_B} \mathcal{O}_{\sigma(T)}) \longrightarrow R^1\pi_*(\mathcal{I} \cdot E) \longrightarrow R^1\pi_*E.$$

By the last paragraph, $R^1\pi_*E = \{0\}$. Since E is generated by global sections, $\pi_*E \rightarrow \pi_*(E \otimes_{\mathcal{O}_B} \mathcal{O}_{\sigma(T)})$ is surjective. Therefore $R^1\pi_*(\mathcal{I} \cdot E) = \{0\}$.

(iii): By (i), $h^1(B, E') = 0$; therefore every global section of E'' is the image of a global section of E . So the global sections of E generate E'' . And the global sections of E' generate E' . Therefore E is generated by global sections. \square

Let B be a prestable curve of genus 0, and let E be a locally free sheaf of positive rank on B . A *smoothing* of the pair (B, E) over a discrete valuation ring R is a pair $(\mathcal{B}, \mathcal{E})$ consisting of a family $\mathcal{B} \rightarrow \text{Spec } R$ of prestable curves of genus 0 and a locally free sheaf \mathcal{E} such that the generic fiber of \mathcal{B} is a smooth curve, such that the closed fiber of \mathcal{B} is isomorphic to B , and such that the restriction of \mathcal{E} to the closed fiber is isomorphic to E . What conditions on (B, E) guarantee that for every smoothing $(\mathcal{B}, \mathcal{E})$, the restriction of \mathcal{E} to the generic fiber is an ample locally free sheaf? Certainly if E is ample, this is true. But E need not be ample for this condition to hold: e.g. if E is an invertible sheaf such that the total degree of E is positive, then for every smoothing the restriction of \mathcal{E} to the generic fiber is ample. Although it is not the most general criterion, the following criterion is used in the rest of the paper.

Definition 2.4. Let B be a connected, proper, at-worst-nodal curve of arithmetic genus 0. A locally free sheaf E on B with positive rank is *deformation ample* if

- (i) E is generated by global sections, and
- (ii) $h^1(B, E(K_B)) = 0$, where $\mathcal{O}_B(K_B)$ is the dualizing sheaf of B .

Remark 2.5.

- (i) Conditions (i) and (ii) above are independent.
- (ii) If E is invertible, then E is deformation ample iff the restriction of E to every irreducible component has nonnegative degree and the restriction to at least one irreducible component has positive degree, cf. Lemma 2.11.
- (iii) One can determine whether E is deformation ample in terms of the splitting type of the restriction of E to each irreducible component together with the patching isomorphisms at the nodes of B .

Let T be a scheme, let $\pi : B \rightarrow T$ be a family of prestable curves of genus 0, and let E be a locally free sheaf of positive rank on B .

Definition 2.6. The sheaf E is π -relatively deformation ample (or simply *deformation ample* if π is understood) if

- (i) E is π -relatively generated by global sections, and
- (ii) $R^1\pi_*(E(K_\pi)) = \{0\}$, where $\mathcal{O}_B(K_\pi)$ is the relative dualizing sheaf of π .

Let $\pi : B \rightarrow T$ be a proper, flat family of connected, at-worst-nodal curves of arithmetic genus 0, let E be a locally free sheaf on B of positive rank, and let $f : T' \rightarrow T$ be a morphism of schemes. Denote the fiber product as in the diagram,

$$\begin{array}{ccc} B' & \xrightarrow{g} & B \\ \pi' \downarrow & & \downarrow \pi \\ T' & \xrightarrow{f} & T \end{array}$$

Denote by E' the pullback g^*E .

Lemma 2.7. *If E is π -relatively deformation ample, then E' is π' -relatively deformation ample. If f is surjective, the converse also holds.*

Proof. (\Rightarrow): For the main direction, by [10, Sec. 8.5.2, Prop. 8.9.1], it suffices to consider the case when T and T' are Noetherian affine schemes.

There is a canonical map of $\mathcal{O}_{T'}$ -modules, $\nu : f^*\pi_*E \rightarrow (\pi')_*g^*E$, which fits into a commutative diagram,

$$\begin{array}{ccc} (\pi')^*f^*\pi_*E & \xrightarrow{=} & g^*\pi^*\pi_*E \\ (\pi')^*\nu \downarrow & & \downarrow g^*\mu_E \\ (\pi')^*(\pi')_*E' & \xrightarrow{\mu_{E'}} & E' \end{array}$$

Since μ_E is surjective, $g^*\mu_E$ is surjective. Hence also $\mu_{E'}$ is surjective, i.e., E' is π' -relatively generated by global sections.

Since π has relative dimension 1, $R^2\pi_*E(K_\pi) = \{0\}$. By [13, Thm. III.12.11(b)], for every closed point $t \in T$, $h^1(B_t, E(K_\pi)|_{B_t}) = 0$. By [13, Prop. III.9.3], for every closed point $t' \in T'$, $h^1(B'_{t'}, E'(K_{\pi'})|_{B'_{t'}}) = 0$. So by [13, Thm. III.12.11(a)] and Nakayama's lemma, $R^1\pi'_*(E'(K_{\pi'})) = \{0\}$. Hence E' is π' -relatively deformation ample.

(\Leftarrow): Now suppose that f is surjective and that E' is π' -relatively deformation ample. As above, it suffices to consider the case when T and T' are Noetherian affine schemes. As above, for every closed point $t' \in T'$, $h^1(B'_{t'}, E'(K_{\pi'})|_{B'_{t'}}) = 0$. Since $T' \rightarrow T$ is surjective, by [13, Prop. III.9.3] for every closed point $t \in T$, $h^1(B_t, E(K_\pi)|_{B_t}) = 0$. So by [13, Thm III.12.11(a)] and Nakayama's lemma, $R^1\pi_*(E(K_\pi)) = \{0\}$.

It remains to prove that E is π -relatively generated by global sections. For every closed point $t \in T$, there is a closed point $t' \in T'$ mapping to t . Since $E'|_{B_{t'}}$ is generated by global sections, also $E|_{B_t}$ is generated by global sections. By Lemma 2.3, $h^1(B_t, E|_{B_t}) = 0$. By [13, Thm. III.12.11(a)] and Nakayama's lemma, $R^1\pi_*(E) = \{0\}$.

The claim is that for any coherent \mathcal{O}_T -module \mathcal{F} , $R^1\pi_*(\pi^*\mathcal{F} \otimes E) = \{0\}$. This is local on T . Locally on T , \mathcal{F} is the cokernel of $\mathcal{O}_T^{\oplus N}$ for some N , so $\pi^*\mathcal{F} \otimes E$ is the cokernel of $E^{\oplus N}$. Since π has relative dimension 1, $R^1\pi_*$ is right exact on the category of coherent \mathcal{O}_B -modules. Since $R^1\pi_*(E^{\oplus N}) = \{0\}$, also $R^1\pi_*(\pi^*\mathcal{F} \otimes E) = \{0\}$, which proves the claim.

In particular, applying the long exact sequence of higher direct images to the short exact sequence,

$$0 \longrightarrow \pi^*\mathcal{I}_t \otimes E \longrightarrow E \longrightarrow E|_{B_t} \longrightarrow 0,$$

$\pi_*(E) \rightarrow H^0(B_t, E|_{B_t})$ is surjective. Since $E|_{B_t}$ is generated by global sections for every closed point $t \in T$, E is π -relatively generated by global sections. So E is π -relatively deformation ample. \square

Lemma 2.8. *Let $\pi : B \rightarrow T$ be a proper, flat family of connected, smooth curves of genus 0 and let E be a locally free sheaf of positive rank. Then E is π -relatively deformation ample iff E is π -relatively ample.*

Proof. Both properties are local on T and can be checked after étale, surjective base-change of T . So it suffices to consider the case when $\pi : B \rightarrow T$ is isomorphic to $\pi_T : T \times \mathbb{P}^1 \rightarrow T$.

(\Rightarrow): Denote $F = (\pi_T)_*(E \otimes \pi_{\mathbb{P}^1}^*\mathcal{O}_{\mathbb{P}^1}(-1))$. Tensoring the map $\mu_{E \otimes \pi_{\mathbb{P}^1}^*\mathcal{O}_{\mathbb{P}^1}(-1)}$ with the identity map on $\pi^*\mathcal{O}_{\mathbb{P}^1}(-1)$ gives a map $\nu : \pi_T^*F \otimes \pi_{\mathbb{P}^1}^*\mathcal{O}_{\mathbb{P}^1}(1) \rightarrow E$. Assume that E is deformation ample. The claim is that ν is surjective. To prove this, it suffices to prove,

- (i) For every geometric point t of T , $h^1(\mathbb{P}_{\kappa(t)}^1, E|_{B_t} \otimes \mathcal{O}_{\mathbb{P}^1}(-1)) = 0$,
- (ii) $F \otimes_{\mathcal{O}_T} \kappa(t) = H^0(\mathbb{P}_{\kappa(t)}^1, E|_{B_t} \otimes \mathcal{O}_{\mathbb{P}^1}(-1))$, and
- (iii) the map $H^0(\mathbb{P}_{\kappa(t)}^1, E|_{B_t} \otimes \mathcal{O}_{\mathbb{P}^1}(-1)) \otimes \mathcal{O}_{\mathbb{P}^1}(1) \rightarrow E|_{B_t}$ is surjective.

By Grothendieck's lemma [13, Exer. V.2.6], $E|_{B_t}$ splits as a direct sum $\mathcal{O}_{\mathbb{P}^1}(a_1) \oplus \cdots \oplus \mathcal{O}_{\mathbb{P}^1}(a_r)$ for some integers $a_1 \leq \cdots \leq a_r$. By Lemma 2.7, $E|_{B_t}$ is deformation ample, and in particular $h^1(\mathbb{P}^1, E|_{B_t}(-2)) = 0$. Hence $a_1 \geq 1$, and $h^1(\mathbb{P}^1, E|_{B_t} \otimes \mathcal{O}_{\mathbb{P}^1}(-1)) = 0$, i.e., (i) holds. By [13, Thm. III.12.11(b)], also (ii) holds. Finally, for $a_i \geq 1$, $H^0(\mathbb{P}_{\kappa}^1, \mathcal{O}_{\mathbb{P}^1}(a_i - 1)) \otimes \mathcal{O}_{\mathbb{P}^1}(1) \rightarrow \mathcal{O}_{\mathbb{P}^1}(a_i)$ is surjective. Thus (iii) holds and the claim is proved.

Now $\pi_T^*F \otimes \pi_{\mathbb{P}^1}^*\mathcal{O}_{\mathbb{P}^1}(1)$ is π_T -relatively ample. Since E is a quotient of $\pi_T^*F \otimes \pi_{\mathbb{P}^1}^*\mathcal{O}_{\mathbb{P}^1}(1)$, also E is π_T -relatively ample, cf. Lemma 2.10 (i).

(\Leftarrow): The converse direction follows in the same way. \square

Lemma 2.9. *There exists an open subscheme $U : U \rightarrow T$ with the following property: for every morphism $f : T' \rightarrow T$, $f(T')$ is contained in U iff E' is π' -relatively deformation ample.*

Proof. By [10, Sec. 8.5.2, Prop. 8.9.1], it suffices to consider the case that T and T' are Noetherian affine schemes.

Let $Z_1 \subset T$ be the closed subset,

$$Z_1 = f[\text{Supp}(\text{coker}(\pi^*\pi_*E \rightarrow E))].$$

Let $Z_2 \subset T$ be the closed subset,

$$Z_2 = \text{Supp}(R^1\pi_*(E(K_\pi))).$$

And let $Z_3 \subset T$ be the closed subset,

$$Z_3 = \text{Supp}(R^1\pi_*E).$$

Let $i : U \rightarrow T$ be the open complement of $Z_1 \cup Z_2 \cup Z_3$.

Let $f : T' \rightarrow T$ be a morphism of Noetherian affine schemes. By [13, Thm. III.12.11, Prop. III.9.3] and Nakayama's lemma, $R^1\pi'_*(E'(K_{\pi'})) = \{0\}$ iff for each closed point $t' \in T'$, $h^1(B'_{t'}, E'(K_{\pi'})|_{B'_{t'}}) = 0$. Denoting $t = f(t')$, $h^1(B'_{t'}, E'(K_{\pi'})|_{B'_{t'}}) = 0$ iff $h^1(B_t, E(K_\pi)|_{B_t}) = 0$, i.e., iff t is contained in the complement of Z_2 . Hence $R^1\pi'_*(E'(K_{\pi'})) = \{0\}$ iff $f(T')$ is contained in the complement of Z_2 .

By the same argument as in the converse direction of the proof of Lemma 2.7, E' is π' -relatively generated by global sections iff for every closed point $t' \in T'$, $E'|_{B'_{t'}}$ is generated by global sections. Denoting $t = f(t')$, $E'|_{B'_{t'}}$ is generated by global sections iff $E|_{B_t}$ is generated by global sections. If $E|_{B_t}$ is generated by global sections, then $h^1(B_t, E|_{B_t}) = 0$. By [13, Thm. III.12.11], t is not in Z_3 and t is not in Z_1 . Conversely, if t is not in Z_3 , then $E|_{B_t}$ is generated by global sections iff t is not in Z_1 . Thus E' is π' -relatively generated by global sections iff $f(T')$ is contained in the complement of $Z_1 \cup Z_3$. So E' is π' -relatively deformation ample iff $f(T')$ is contained in U . \square

Lemma 2.10.

- (i) *If $\chi : E \rightarrow E''$ is a morphism of locally free sheaves on B whose cokernel is torsion in every fiber (in particular, if χ is surjective), if E'' is nonzero, and if E is π -relatively deformation ample, then also E'' is π -relatively deformation ample.*
- (ii) *If E' and E'' are π -relatively deformation ample, then for every short exact sequence of coherent \mathcal{O}_B -modules,*

$$0 \longrightarrow E' \xrightarrow{v} E \xrightarrow{\chi} E'' \longrightarrow 0,$$

E is π -relatively deformation ample.

- (iii) *If E is π -relatively deformation ample, then for every integer $n \geq 1$, also $E^{\otimes n}$ is π -relatively deformation ample.*

Proof. (i): Let Q denote the cokernel of χ and let $I \subset E''$ denote the image of χ . There is a short exact sequence of coherent \mathcal{O}_B -modules:

$$0 \longrightarrow I \longrightarrow E'' \longrightarrow Q \longrightarrow 0.$$

When we “twist” this exact sequence by $\mathcal{O}_B(K_\pi)$, it remains exact. Because π has relative dimension 1, $R^1\pi_*$ is right exact on the category of coherent \mathcal{O}_B -modules. In particular, since $R^1\pi_*E(K_\pi) = \{0\}$, also $R^1\pi_*I(K_\pi) = \{0\}$. Since Q is torsion in every fiber, $R^1\pi_*Q(K_\pi) = \{0\}$. Thus, by the long exact sequence of higher direct images associated to the twisted exact sequence above, $R^1\pi_*E''(K_\pi) = \{0\}$.

The surjective composition map

$$\pi^*\pi_*E_1 \xrightarrow{\mu_E} E_1 \longrightarrow I$$

factors through the natural map $\mu_I : \pi^*\pi_*I \rightarrow I$. Hence μ_I is surjective, i.e., I is π -relatively generated by global sections. Since Q is torsion in every fiber, the support of Q is finite over T and it follows that Q is π -relatively generated by global sections. By Lemma 2.3 (iii), E'' is π -relatively generated by global sections. So E'' is π -relatively deformation ample.

(ii): By hypothesis, $R^1\pi_*E'(K_\pi) = R^1\pi_*E''(K_\pi) = \{0\}$. By the long exact sequence of higher direct images, also $R^1\pi_*E(K_\pi) = \{0\}$. By Lemma 2.3 (iii), E is π -relatively generated by global sections. So E is π -relatively deformation ample.

(iii): This is proved by induction on n , the case $n = 1$ being tautological. It suffices to consider the case when T is affine. Suppose $n > 1$ and suppose the result is known for $n - 1$. In particular, $E^{\otimes(n-1)}$ is generated by global sections. There is a natural surjection

$$\pi^*\pi_*\left(E^{\otimes(n-1)}\right) \otimes_{\mathcal{O}_B} E \rightarrow E^{\otimes n}.$$

And there is a surjective map $\mathcal{O}_T^{\oplus N} \rightarrow \pi_*(E^{\otimes(n-1)})$. Hence there is a surjection $E^{\oplus N} \rightarrow E^{\otimes n}$. By (ii) and induction, $E^{\oplus N}$ is π -relatively deformation ample. By (i), the quotient $E^{\otimes n}$ is π -relatively deformation ample. Thus (iii) is proved by induction. \square

Lemma 2.11. *Let B be a proper, connected, at-worst-nodal curve of arithmetic genus 0 over an algebraically closed field k . Let E be a locally free sheaf of positive rank such that,*

- (i) *for every irreducible component $B_i \subset B$, $E|_{B_i}$ is generated by global sections, and*
- (ii) *there exists a nonempty, connected, closed subcurve $B' \subset B$ such that $E|_{B'}$ is deformation ample.*

Then E is deformation ample.

Proof. Let δ be the number of irreducible components of B which are not contained in B' . The result is proved by induction on δ . The base case $\delta = 0$ is tautological, for then $B = B'$. Assume that $\delta > 0$ and that the result is true for all smaller values of δ .

Let $B_1 \subset B$ be an irreducible component of B . Let $B_2 \subset B$ denote the union of all irreducible components other than B_1 . There exists an irreducible component B_1 not contained in B' such that B_2 is connected: if the dual graph of B' contains every leaf (= vertex of valence 1) of the dual graph of B , then the two graphs are equal. The intersection $B_1 \cap B_2$ is a single node, denoted b . By the induction hypothesis, $E|_{B_2}$ is deformation ample.

The claim is that E is generated by global sections. Denote by $F \subset E$ the image of $H^0(B, E) \otimes_k \mathcal{O}_B \rightarrow E$. There is a short exact sequence of coherent sheaves:

$$0 \longrightarrow E \otimes_{\mathcal{O}_B} \mathcal{O}_{B_1}(-b) \longrightarrow E \longrightarrow E \otimes_{\mathcal{O}_B} \mathcal{O}_{B_2} \longrightarrow 0.$$

Since $E|_{B_1}$ is a locally free sheaf on \mathbb{P}^1 generated by global sections, Grothendieck's lemma and the cohomology of line bundles on \mathbb{P}^1 imply that $h^1(B, E \otimes_{\mathcal{O}_B} \mathcal{O}_{B_1}(-b)) = 0$ ($h^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(a)) = 0$ for $a \geq -1$). Hence all the global sections of $E|_{B_2}$ lift to global sections of E , i.e., $F \rightarrow E|_{B_2}$ is surjective. So E/F is supported on B_1 . Hence E/F is a quotient of $E|_{B_1}$. Since $E|_{B_1}$ is generated by global sections, also E/F is generated by global sections. There is a short exact sequence

$$0 \longrightarrow F \longrightarrow E \longrightarrow E/F \longrightarrow 0.$$

By Lemma 2.3 (iii), E is generated by global sections.

There is a short exact sequence of coherent sheaves,

$$0 \rightarrow E(K_B) \otimes_{\mathcal{O}_B} \mathcal{O}_{B_2}(-b) \rightarrow E(K_B) \rightarrow E(K_B) \otimes_{\mathcal{O}_B} \mathcal{O}_{B_1} \rightarrow 0.$$

This gives a long exact sequence in cohomology, part of which is,

$$H^1(B, E(K_B) \otimes_{\mathcal{O}_B} \mathcal{O}_{B_2}(-b)) \rightarrow H^1(B, E(K_B)) \rightarrow H^1(B, E(K_B) \otimes_{\mathcal{O}_B} \mathcal{O}_{B_1}) \rightarrow 0.$$

The inclusion map $B_2 \rightarrow B$ is finite, therefore there is a canonical isomorphism of \mathcal{O}_B -modules (cf. [12, Sec. III.6], [17, Cor. 5.68]),

$$\mathcal{O}_{B_2}(K_{B_2}) \cong \text{Hom}_{\mathcal{O}_B}(\mathcal{O}_{B_2}, \mathcal{O}_B(K_B)) \cong \mathcal{O}_B(K_B) \otimes_{\mathcal{O}_B} \mathcal{O}_{B_2}(-b).$$

Hence $h^1(B, E(K_B) \otimes_{\mathcal{O}_B} \mathcal{O}_{B_2}(-b)) = h^1(B_2, E \otimes_{\mathcal{O}_B} \mathcal{O}_{B_2}(K_{B_2}))$, which is zero by the induction assumption. Similarly, $E(K_B) \otimes_{\mathcal{O}_B} \mathcal{O}_{B_1}$ is isomorphic to $E \otimes_{\mathcal{O}_B} \mathcal{O}_{B_1}(-1)$ (identifying B_1 with \mathbb{P}^1). Since $E|_{B_1}$ is generated by global sections, it follows by Grothendieck's lemma and the cohomology of line bundles on \mathbb{P}^1 that $h^1(B_1, E|_{B_1}(-1)) = 0$. Hence $h^1(B, E(K_B)) = 0$, and E is deformation ample. Therefore the lemma is proved by induction on δ . \square

Remark 2.12. A particular case of Lemma 2.11 is when B' is one irreducible component of B , in which case the lemma says that a locally free sheaf on B which is *generically ample* in the sense of Lazarsfeld [8] is deformation ample.

3. DEFORMATION THEORY OF STABLE MAPS

The Kontsevich moduli space of genus 0 stable maps, $\overline{\mathcal{M}}_{0,r}(X, e)$, and the Behrend-Manin moduli spaces, $\overline{\mathcal{M}}(X, \tau)$, are described in Subsection 1.1. The deformation theory of stable maps has been worked out in [4] and [3]. Many specific deformation-theoretic results follow easily from these papers and are known to the experts, but have not been written down. Some of these specific results are proved in this section. Although these results will only be applied to *genus 0* stable maps in this paper, the same arguments work for stable maps of arbitrary genus; in this section only, stable maps and stable A -graphs are not necessarily assumed to be of genus 0.

Let X and T be schemes.

Definition 3.1. A family of r -pointed, genus g prestable maps to X over B is a triple

$$\zeta = ((p : \Sigma \rightarrow B, \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X)$$

consisting of a family $p : \Sigma \rightarrow B$ of prestable curves of genus g over B , a sequence of r disjoint sections $\sigma_i : B \rightarrow \Sigma$ with image contained in the smooth locus of p (if $r = 0$, the sections are omitted), and a morphism $g : \Sigma \rightarrow X$.

Let $((\Sigma, \sigma_1, \dots, \sigma_r), g)$ be a prestable map over an algebraically closed field. An irreducible component $\Sigma_i \subset \Sigma$ is *stable* if the restriction of the *log-dualizing sheaf*, $\mathcal{O}_\Sigma(K_\Sigma + \sigma_1 + \dots + \sigma_r) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_i}$, is g -relatively ample, i.e., one of the following hold,

- (i) $g : \Sigma_i \rightarrow X$ is nonconstant,
- (ii) $p_a(\Sigma_i) > 1$,
- (iii) $p_a(\Sigma_i) = 1$ and Σ_i contains at least one marked point or external node of Σ , or
- (iv) $p_a(\Sigma_i) = 0$ and Σ_i contains at least three marked points and nodes of B .

The curve Σ is *stable* if $\mathcal{O}_\Sigma(K_\Sigma + \sigma_1 + \dots + \sigma_r)$ is g -relatively ample, i.e., every irreducible component of Σ is stable. The family ζ is *stable* if for each geometric point $t \in T$, the curve Σ_t is stable.

Now assume that X is smooth.

Notation 3.2. Denote by L_ζ the complex of coherent sheaves on Σ

$$\begin{array}{ccc} -1 & & 0 \\ g^* \Omega_X & \xrightarrow{(dg)^\dagger} & \Omega_p(\sigma_1(T) + \dots + \sigma_r(T)) \end{array} \quad (1)$$

For a scheme T and a bounded above complex of coherent sheaves C on T , C^\vee denotes the object in the derived category of coherent sheaves on T ,

$$C^\vee := \mathbb{R} \text{Hom}_{\mathcal{O}_T}(C, \mathcal{O}_T).$$

In particular, L_ζ^\vee is the object

$$L_\zeta^\vee = \mathbb{R}Hom_{\mathcal{O}_\Sigma}(L_\zeta, \mathcal{O}_\Sigma)$$

in the derived category of coherent sheaves on Σ .

The relevance of the complex L_ζ^\vee is the following.

Lemma 3.3. *Let X be a smooth quasi-projective scheme. Let $\overline{\mathcal{M}}_{g,r}(X, \beta)$ denote the Deligne-Mumford stack of r -pointed stable maps to X of arithmetic genus g and degree β . Let $p : \Sigma \rightarrow \overline{\mathcal{M}}_{g,r}(X, \beta)$ denote the universal curve, let $\sigma_i : \overline{\mathcal{M}}_{g,r}(X, \beta) \rightarrow \Sigma$ denote the universal sections, and let $g : \Sigma \rightarrow X$ denote the universal map, i.e.*

$$\zeta = ((p : \Sigma \rightarrow \overline{\mathcal{M}}_{g,r}(X, \beta), \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X)$$

is the universal family of stable maps. There is an obstruction theory for $\overline{\mathcal{M}}_{g,r}(X, \beta)$ in the sense of [4, Def. 4.4] of the form

$$(\mathbb{R}p_*(L_\zeta^\vee)[1])^\vee \rightarrow L_{\overline{\mathcal{M}}_{g,r}(X, \beta)}.$$

A similar result holds for prestable maps, cf. Remark 3.4.

Proof. Essentially this follows from [4] and [3]. \square

Remark 3.4. Explicitly, if $\zeta = ((\Sigma, \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X)$ is a stable map or a prestable map, the space of first order deformations of ζ is $\mathbb{E}xt_{\mathcal{O}_\Sigma}^1(L_\zeta, \mathcal{O}_\Sigma)$ and the obstruction group is a subgroup of $\mathbb{E}xt_{\mathcal{O}_\Sigma}^2(L_\zeta, \mathcal{O}_\Sigma)$. In the case of a prestable map, the space of infinitesimal automorphisms of the map is $\mathbb{E}xt_{\mathcal{O}_\Sigma}^0(L_\zeta, \mathcal{O}_\Sigma)$ (for stable maps this group is zero). In particular, if $\mathbb{E}xt_{\mathcal{O}_\Sigma}^2(L_\zeta, \mathcal{O}_\Sigma)$ vanishes, then $\overline{\mathcal{M}}_{g,r}(X, \beta)$ is smooth at the point $[\zeta]$.

Lemma 3.5. *Let $((\Sigma, \sigma_1, \dots, \sigma_r), g)$ be a prestable map. Let Θ_Σ denote the tangent sheaf of Σ , i.e. the dual of Ω_Σ . The space of infinitesimal automorphisms of ζ , $\mathbb{E}xt_{\mathcal{O}_\Sigma}^0(L_\zeta, \mathcal{O}_\Sigma)$, is canonically isomorphic to a subspace of the space of infinitesimal automorphisms of $(\Sigma, \sigma_1, \dots, \sigma_r)$, $H^0(\Sigma, \Theta_\Sigma(-(\sigma_1 + \dots + \sigma_r)))$.*

An irreducible component $\Sigma_i \subset \Sigma$ is stable iff the restriction of every infinitesimal automorphism of ζ to $\Theta_\Sigma(-(\sigma_1 + \dots + \sigma_r)) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_i}$ is zero. Moreover, if Σ_i is unstable, then every infinitesimal automorphism of the nodal curve Σ_i that fixes all marked points and nodes of Σ is the image of an infinitesimal automorphism of ζ .

Proof. Analyzing the spectral sequence for hypercohomology, $\mathbb{E}xt_{\mathcal{O}_\Sigma}^0(L_\zeta, \mathcal{O}_\Sigma)$ is canonically isomorphic to the kernel of

$$\text{Ext}_{\mathcal{O}_\Sigma}^0(\Omega_\Sigma(\sigma_1 + \dots + \sigma_r), \mathcal{O}_\Sigma) \rightarrow \text{Ext}_{\mathcal{O}_\Sigma}^0(g^*\Omega_X, \mathcal{O}_\Sigma).$$

And $\text{Ext}_{\mathcal{O}_\Sigma}^0(\Omega_\Sigma(\sigma_1 + \dots + \sigma_r), \mathcal{O}_\Sigma)$ equals $H^0(\Sigma, \Theta_\Sigma(-(\sigma_1 + \dots + \sigma_r)))$.

Assume that Σ_i is stable, i.e., Σ_i satisfies one of the Cases (i)–(iv) in Definition 3.1. In Cases (ii)–(iv), the image of

$$H^0(\Sigma, \Theta_\Sigma(-(\sigma_1 + \dots + \sigma_r))) \rightarrow H^0(\Sigma_i, \Theta_\Sigma(-(\sigma_1 + \dots + \sigma_r)) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_i})$$

is zero, hence the restriction of every infinitesimal automorphism of ζ is zero. In Case (i), chasing through diagrams, the image of $H^0(\Sigma, \Theta_\Sigma(-(\sigma_1 + \dots + \sigma_r)))$ is contained in the subsheaf which is the kernel of $d(g|_{\Sigma_i}) : \Theta_{\Sigma_i} \rightarrow g^*T_X \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_i}$. Since g is non-constant, this map is nonzero. And Θ_{Σ_i} is a torsion-free sheaf, so the kernel of $d(g|_{\Sigma_i})$ is zero. Hence, also in Case (i), the restriction of every infinitesimal automorphism of ζ is zero.

Assume that Σ_i is not stable. Then g contracts Σ_i to a point, and either $\Sigma = \Sigma_i$ is a curve of arithmetic genus 1 and there are no marked points, or Σ_i is a smooth rational curve which contains at most two marked points and nodes of Σ . In both cases there is a positive dimensional group of automorphisms of Σ_i which fix all marked points and nodes of Σ : $Pic^\tau(\Sigma_i)$ acting by translation if Σ_i has arithmetic genus 1, and the group of automorphisms of \mathbb{P}^1 fixing (at most) two points if Σ_i is a smooth rational curve. The Lie algebra of this positive dimensional group is the Lie algebra of infinitesimal automorphisms of the marked curve Σ_i . This group of automorphisms of Σ_i extends to a group of automorphisms of the map ζ . Therefore every infinitesimal automorphism of the marked curve Σ_i extends to an infinitesimal automorphism of ζ . In particular there exists an infinitesimal automorphism of ζ whose restriction to Σ_i is nonzero. \square

Let $f : X \rightarrow Y$ be a smooth morphism of smooth quasi-projective varieties. Let $\zeta = ((\Sigma, \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X)$ be a prestable map. Denote by $h : \Sigma \rightarrow Y$ the composition $h = f \circ g$, and denote by ξ the prestable map $((\Sigma, \sigma_1, \dots, \sigma_r), h : \Sigma \rightarrow Y)$. There is a short exact sequence of complexes

$$0 \longrightarrow L_\xi \longrightarrow L_\zeta \longrightarrow g^*\Omega_f[1] \longrightarrow 0,$$

defined by the following commutative diagram,

$$\begin{array}{ccc} & -1 & 0 \\ L_\xi : & g^*f^*\Omega_Y & \xrightarrow{dh^\dagger} \Omega_\Sigma(\sigma_1 + \dots + \sigma_r) \\ & \downarrow g^*(df)^\dagger \downarrow & \downarrow = \\ L_\zeta : & g^*\Omega_X & \xrightarrow{df^\dagger} \Omega_\Sigma(\sigma_1 + \dots + \sigma_r) \\ & \downarrow \quad \downarrow & \downarrow \\ g^*\Omega_f[1] : & g^*\Omega_f & \longrightarrow 0 \end{array}$$

Lemma 3.6. (i) *If ξ is stable, then ζ is stable.*

(ii) *If the dimension of the obstruction group of ζ is 0, then the dimension of the obstruction group of ξ is 0.*

(iii) *If $h^1(\Sigma, g^*\Omega_f^\vee) = 0$ and if the dimension of the obstruction group of ξ is 0, then the dimension of the obstruction group of ζ is 0, and the map from the space of first order deformations of ζ to the space of first order deformations of ξ is surjective.*

Proof. (i): If the log dualizing sheaf of $(\Sigma, \sigma_1, \dots, \sigma_r)$ is h -ample, then it is g -ample.

(ii): Associated to the short exact sequence of complexes above, there is a long exact sequence of hyperExt, part of which is

$$\mathbb{E}xt_{\mathcal{O}_\Sigma}^2(L_\zeta, \mathcal{O}_\Sigma) \longrightarrow \mathbb{E}xt_{\mathcal{O}_\Sigma}^2(L_\xi, \mathcal{O}_\Sigma) \longrightarrow \mathbb{E}xt_{\mathcal{O}_\Sigma}^3(g^*\Omega_f[1], \mathcal{O}_\Sigma).$$

Of course $\mathbb{E}xt_{\mathcal{O}_\Sigma}^k(g^*\Omega_f[1], \mathcal{O}_\Sigma) = H^{k-1}(\Sigma, g^*\Omega_f^\vee)$. In particular $\dim \mathbb{E}xt_{\mathcal{O}_\Sigma}^3(g^*\Omega_f[1], \mathcal{O}_\Sigma) = h^2(\Sigma, g^*\Omega_f^\vee) = 0$. Therefore if $\dim \mathbb{E}xt_{\mathcal{O}_\Sigma}^2(L_\zeta, \mathcal{O}_\Sigma) = 0$, then $\dim \mathbb{E}xt_{\mathcal{O}_\Sigma}^2(L_\xi, \mathcal{O}_\Sigma) = 0$.

(iii): If $h^1(\Sigma, g^*\Omega_f^\vee) = 0$, then $\mathbb{E}xt_{\mathcal{O}_\Sigma}^2(L_\zeta, \mathcal{O}_\Sigma) \rightarrow \mathbb{E}xt_{\mathcal{O}_\Sigma}^2(L_\xi, \mathcal{O}_\Sigma)$ is an isomorphism. Hence if the dimension of the obstruction group of ζ is 0, then the dimension of the obstruction group of ξ is 0. Moreover the preceding two terms in the long exact sequence of hyperExts give a surjection $\mathbb{E}xt_{\mathcal{O}_\Sigma}^1(L_\zeta, \mathcal{O}_\Sigma) \rightarrow \mathbb{E}xt_{\mathcal{O}_\Sigma}^1(L_\xi, \mathcal{O}_\Sigma)$. \square

Let $q : S \rightarrow B$ be a smooth morphism and let $i : \Sigma \rightarrow S$ be an unramified morphism of B -schemes. Then $(g, i) : \Sigma \rightarrow X \times S$ is an unramified morphism of B -schemes, hence the map of

coherent sheaves

$$d(g, i)^\dagger : (g, i)^*\Omega_{(X \times S/B)} \rightarrow \Omega_p$$

is surjective. Because p is flat of relative dimension 1 and the geometric fibers are reduced, local complete intersection schemes, the kernel of $d(g, i)^\dagger$ is a locally free sheaf $N_{(g, i)}^\vee$. Similarly the kernel of $di^\dagger : i^*\Omega_q \rightarrow \Omega_p$ is a locally free sheaf N_i^\vee .

There is a short exact sequence of coherent sheaves

$$0 \longrightarrow N_i^\vee \longrightarrow N_{(g, i)}^\vee \longrightarrow g^*\Omega_X \longrightarrow 0.$$

Denote by $N_{(\zeta, i)}^\vee$ the subsheaf of $N_{(g, i)}^\vee(\sigma_1(B) + \cdots + \sigma_r(B))$ that contains $N_i^\vee(\sigma_1(B) + \cdots + \sigma_r(B))$ and such that $N_{(\zeta, i)}^\vee/N_i^\vee(\sigma_1(B) + \cdots + \sigma_r(B))$ is identified with $g^*\Omega_X \subset g^*\Omega_X(\sigma_1(B) + \cdots + \sigma_r(B))$. Observe that $N_{(\zeta, i)}^\vee$ is a locally free sheaf.

Notation 3.7. There is a canonical map $N_{(g, i)}^\vee(\sigma_1(B) + \cdots + \sigma_r(B)) \rightarrow (g, i)^*\Omega_{X \times S/B}$. The two projections give a canonical isomorphism of $\Omega_{X \times S/B}$ with $\pi_X^*\Omega_X \oplus \pi_S^*\Omega_q$. Denote by $\gamma_{(\zeta, i)} : N_{(\zeta, i)}^\vee \rightarrow g^*\Omega_X(\sigma_1(B) + \cdots + \sigma_r(B)) \oplus i^*\Omega_q(\sigma_1(B) + \cdots + \sigma_r(B))$ the induced morphism. Observe that the composition of $\gamma_{(\zeta, i)}$ with projection on the first summand factors through $g^*\Omega_X \subset g^*\Omega_X(\sigma_1(B) + \cdots + \sigma_r(B))$. Denote by $\alpha_{(\zeta, i)} : N_{(\zeta, i)}^\vee \rightarrow g^*\Omega_X$ and $\beta_{(\zeta, i)} : N_{(\zeta, i)}^\vee \rightarrow i^*\Omega_q(\sigma_1(B) + \cdots + \sigma_r(B))$ the composition of $\gamma_{(\zeta, i)}$ with the two projections. Denote by $L_{(\zeta, i)}$ the complex of locally free \mathcal{O}_Σ -modules concentrated in degrees $[-1, 0]$,

$$\begin{array}{ccc} -1 & & 0 \\ N_{(\zeta, i)}^\vee & \xrightarrow{\beta_{(\zeta, i)}} & i^*\Omega_q(\sigma_1(B) + \cdots + \sigma_r(B)). \end{array}$$

Denote by $\lambda_{(\zeta, i)} : L_{(\zeta, i)} \rightarrow L_\zeta$ the quasi-isomorphism of complexes of coherent \mathcal{O}_Σ -modules,

$$\begin{array}{ccc} N_{(\zeta, i)}^\vee & \xrightarrow{\beta_{(\zeta, i)}} & i^*\Omega_q(\sigma_1(B) + \cdots + \sigma_r(B)) \\ \alpha_{(\zeta, i)} \downarrow & & \downarrow (di)^\dagger \\ g^*\Omega_X & \xrightarrow{(dg)^\dagger} & \Omega_p(\sigma_1(B) + \cdots + \sigma_r(B)) \end{array}.$$

The relevance of $\lambda_{(\zeta, i)} : L_{(\zeta, i)} \rightarrow L_\zeta$ is that the complex $L_{(\zeta, i)}^\vee$ is easy to compute since $L_{(\zeta, i)}$ is a complex of locally free sheaves; it is simply

$$\begin{array}{ccc} 0 & & 1 \\ i^*T_q(-(\sigma_1(B) + \cdots + \sigma_r(B))) & \xrightarrow{\beta_{(\zeta, i)}^\dagger} & N_{(\zeta, i)}^\vee. \end{array}$$

In most applications, B will be the spectrum of a field and S will be a surface.

3.1. Contracting unstable components. In this subsection, the base B will always be the spectrum of an algebraically closed field. The changes necessary to get relative versions of the lemmas over a more general base are straightforward.

Let

$$\zeta = ((\Sigma, \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X)$$

be a prestable map, let

$$(\Sigma', \sigma'_1, \dots, \sigma'_r, \sigma''_1, \dots, \sigma''_s)$$

be a proper, connected, at-worst-nodal curve, and let

$$u : (\Sigma', \sigma'_1, \dots, \sigma'_r) \rightarrow (\Sigma, \sigma_1, \dots, \sigma_r)$$

be a map which contracts some of the unstable components of $(\Sigma', \sigma'_1, \dots, \sigma'_r)$ (i.e., u contracts some of the irreducible components of Σ' which have arithmetic genus 0 and contain fewer than three nodes and marked points). Denote $g' = g \circ u$ and denote by ζ' the prestable map

$$\zeta' = ((\Sigma', \sigma'_1, \dots, \sigma'_r, \sigma''_1, \dots, \sigma''_s), g' : \Sigma' \rightarrow X).$$

What is the relationship of $\mathbb{E}xt_{\mathcal{O}_\Sigma}(L_\zeta, \mathcal{O}_\Sigma)$ and $\mathbb{E}xt_{\mathcal{O}_{\Sigma'}}(L_{\zeta'}, \mathcal{O}_{\Sigma'})$?

Any morphism $u : \Sigma' \rightarrow \Sigma$ as above can be factored as a sequence of *elementary morphisms*, defined below.

- Definition 3.8.**
- (i) The morphism u is a *Type I elementary morphism* if $\zeta = (\Sigma, g : \Sigma \rightarrow X)$ is a prestable map without marked points, $u : \Sigma' \rightarrow \Sigma$ contracts a single unstable component to a smooth point of Σ , and $\zeta' = (\Sigma', g' = g \circ u : \Sigma' \rightarrow X)$.
 - (ii) The morphism u is a *Type II elementary morphism* if $\zeta = (\Sigma, g : \Sigma \rightarrow X)$ is a prestable map without marked points, $u : \Sigma' \rightarrow \Sigma$ contracts a single unstable component to a node of Σ , and $\zeta' = (\Sigma', g' = g \circ u : \Sigma' \rightarrow X)$.
 - (iii) The morphism u is a *Type III elementary morphism* if $\zeta = (\Sigma, (\sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X)$ is a marked prestable map, ζ' is the same prestable map but with one extra marked point, and $u : \Sigma' \rightarrow \Sigma$ is the identity map.

Let S be a smooth surface and let $i : \Sigma \rightarrow S$ be a closed immersion. Let $s \in \Sigma \subset S$ be a closed point.

Notation 3.9. Denote by $v : S' \rightarrow S$ the blowing up of S at s . Denote by $E \subset S$ the exceptional divisor. Denote by $i' : \Sigma' \rightarrow S'$ the *reduced total transform of B* ; i.e., the reduced scheme of $v^{-1}(\Sigma)$. Denote by v also the morphism of pairs $v : (S', \Sigma') \rightarrow (S, \Sigma)$. Denote by $u : \Sigma' \rightarrow \Sigma$ the restriction of v to Σ' . Denote by $g' : \Sigma' \rightarrow X$ the composition $g' = g \circ u$. Denote by ζ' the prestable map $(\Sigma', g' : \Sigma' \rightarrow X)$. Denote by $\Gamma \subset \Sigma'$ the closed (not necessarily connected) subcurve which is the union of all irreducible components other than E , and denote $D = E \cap \Gamma$.

- Definition 3.10.**
- (i) The morphism of pairs is *Type Ia* if $s \in \Sigma$ is a smooth point that lies on a stable component.
 - (ii) The morphism of pairs is *Type Ib* if $s \in \Sigma$ is a smooth point that lies on an unstable component.
 - (iii) The morphism of pairs is *Type I* if it is Type Ia or Type Ib.
 - (iv) The morphism of pairs is *Type IIa* if $s \in \Sigma$ is a node, and there exists a first order deformation of ζ that smoothes the node s (to first order).
 - (v) The morphism of pairs is *Type IIb* if $s \in \Sigma$ is a node, and there is no first order deformation of ζ that smoothes the node s (to first order).
 - (vi) The morphism of pairs is *Type II* if it is Type IIa or Type IIb.

If v is Type I, then $v^*\Sigma = \Sigma'$ as Cartier divisors. If v is Type II, then $v^*\Sigma = \Sigma' + E$ as Cartier divisors.

Lemma 3.11. *For every integer k the pullback morphism of sheaf hypercohomology groups, $\mathbb{H}^k(\Sigma, L_\zeta^\vee) \rightarrow \mathbb{H}^k(\Sigma', \mathbb{L}u^*L_\zeta^\vee)$, is an isomorphism.*

Proof. Of course $v_*\mathcal{O}_{S'} = \mathcal{O}_S$ and $R^k v_*\mathcal{O}_{S'} = \{0\}$ if $k > 0$. Also $v_*(\mathcal{O}_{S'}(E)) = \mathcal{O}_S$ and $R^k v_*(\mathcal{O}_{S'}(E)) = \{0\}$ for $k > 0$. If v is Type I then $\mathcal{O}_{S'}(-\Sigma') = v^*\mathcal{O}_S(-\Sigma)$, and if v is Type II then $\mathcal{O}_{S'}(-\Sigma') = v^*\mathcal{O}_S(-\Sigma) \otimes_{\mathcal{O}_{S'}} \mathcal{O}_{S'}(E)$. For both types, the projection formula implies that $v_*(\mathcal{O}_{S'}(-\Sigma')) = \mathcal{O}_S(-\Sigma)$ and $R^k v_*(\mathcal{O}_{S'}(-\Sigma')) = \{0\}$ for $k > 0$.

Associated to the short exact sequence of coherent $\mathcal{O}_{S'}$ -modules,

$$0 \longrightarrow \mathcal{O}_{S'}(-\Sigma') \longrightarrow \mathcal{O}_{S'} \longrightarrow \mathcal{O}_{\Sigma'} \longrightarrow 0,$$

there is a long exact sequence of higher direct images $R^k v_*$, and the higher direct images of $\mathcal{O}_{S'}$ and $\mathcal{O}_{S'}(-\Sigma')$ have just been computed. The conclusion is that $u_* \mathcal{O}_{\Sigma'} = \mathcal{O}_{\Sigma}$ and $R^k u_* \mathcal{O}_{\Sigma'} = \{0\}$ for $k > 0$. In other words, the canonical morphism of complexes of coherent \mathcal{O}_{Σ} -modules, $\mathcal{O}_{\Sigma}[0] \rightarrow \mathbb{R}u_* \mathcal{O}_{\Sigma'}$, is a quasi-isomorphism. From this and the projection formula, it follows that the canonical morphism

$$L_{(\zeta, i)}^{\vee} \rightarrow \mathbb{R}u_* \mathbb{L}u^*(L_{(\zeta, i)}^{\vee})$$

is a quasi-isomorphism. Therefore the pullback morphisms

$$\mathbb{H}^k(\Sigma, L_{\zeta}^{\vee}) \rightarrow \mathbb{H}^k(\Sigma', \mathbb{L}u^* L_{\zeta}^{\vee})$$

are isomorphisms. \square

There is a canonical map of coherent sheaves $\mathcal{O}_{S'}(\Sigma') \rightarrow v^* \mathcal{O}_S(\Sigma)$. It is an isomorphism if v is Type I, and is injective with cokernel $v^* \mathcal{O}_S(\Sigma) \otimes_{\mathcal{O}_{S'}} \mathcal{O}_E$ if v is Type II. If v is Type II, then $v^* \mathcal{O}_X(\Sigma) \otimes_{\mathcal{O}_{S'}} \mathcal{O}_E \cong M \otimes_{\mathbb{C}} \mathcal{O}_E$ where $M = \mathcal{O}_S(\Sigma)|_s$ is a one-dimensional vector space. If v is Type I, the canonical morphism $N_{(\zeta', i')} \rightarrow u^* N_{(\zeta, i)}$ is an isomorphism. If v is Type II, there is an exact sequence:

$$0 \rightarrow M \otimes_{\mathbb{C}} \text{Tor}_1^{\mathcal{O}_{S'}}(\mathcal{O}_{\Sigma'}, \mathcal{O}_E) \rightarrow N_{(\zeta', i')} \rightarrow u^* N_{(\zeta, i)} \rightarrow M \otimes_{\mathbb{C}} \mathcal{O}_E \rightarrow 0$$

Lemma 3.12. *Let $N_{E/S'}^{\vee}$ denote the conormal sheaf of $E \subset S'$. There is a canonical isomorphism $\text{Tor}_1^{\mathcal{O}_{S'}}(\mathcal{O}_{\Sigma'}, \mathcal{O}_E) \cong N_{E/S'}^{\vee} \otimes_{\mathcal{O}_E} \mathcal{O}_E(-D)$.*

Proof. There is an $\mathcal{O}_{S'}$ -flat resolution of \mathcal{O}_E ,

$$0 \longrightarrow \mathcal{O}_{S'}(-E) \longrightarrow \mathcal{O}_S \longrightarrow \mathcal{O}_E \longrightarrow 0.$$

Tensoring this resolution with $\mathcal{O}_{\Sigma'}$ over $\mathcal{O}_{S'}$ gives a canonical isomorphism,

$$\text{Tor}_1^{\mathcal{O}_{S'}}(\mathcal{O}_{\Sigma'}, \mathcal{O}_E) \cong \mathcal{I}_{\Gamma} \otimes_{\mathcal{O}_{S'}} \mathcal{O}_{S'}(-E) \cong N_{E/S'}^{\vee} \otimes_{\mathcal{O}_E} \mathcal{O}_E(-D).$$

where $\mathcal{I}_{\Gamma} \subset \mathcal{O}_{\Sigma'}$ is the ideal sheaf of $\Gamma \subset \Sigma'$. \square

In particular, if v is Type II there is an exact sequence:

$$0 \rightarrow M \otimes_{\mathbb{C}} N_{E/S'}^{\vee} \otimes_{\mathcal{O}_E} \mathcal{O}_E(-D) \rightarrow N_{(\zeta', i')} \rightarrow u^* N_{(\zeta, i)} \rightarrow M \otimes_{\mathbb{C}} \mathcal{O}_E \rightarrow 0.$$

Lemma 3.13. *There is a long exact sequence,*

$$0 \rightarrow T_E(-D) \rightarrow (i')^* T_{S'} \xrightarrow{dv} u^* i^* T_S \rightarrow T_E \otimes_{\mathcal{O}_E} N_{E/S'} \rightarrow 0.$$

(Both if v is Type I and if v is Type II.)

Proof. For both types, there is a short exact sequence of coherent $\mathcal{O}_{S'}$ -modules

$$0 \longrightarrow v^* \Omega_S \xrightarrow{(dv)^{\dagger}} \Omega_{S'} \longrightarrow \Omega_E \longrightarrow 0.$$

There is an associated long exact sequence of higher derived functors of $\text{Hom}_{\mathcal{O}_{S'}}(\cdot, \mathcal{O}_{S'})$, part of which is the short exact sequence,

$$0 \longrightarrow T_{S'} \longrightarrow v^* T_S \longrightarrow \text{Ext}_{\mathcal{O}_{S'}}^1(\Omega_E, \mathcal{O}_{S'}) \longrightarrow 0.$$

The resolution of \mathcal{O}_E from the proof of Lemma 3.12 gives a canonical isomorphism,

$$\text{Ext}_{\mathcal{O}_{S'}}^1(\mathcal{O}_E, \mathcal{O}_{S'}) \cong N_{E/S'},$$

where $N_{E/S'}$ is the dual of $N_{E/S'}^{\vee}$. So the previous exact sequence is,

$$0 \longrightarrow T_{S'} \longrightarrow v^* T_S \longrightarrow T_E \otimes_{\mathcal{O}_E} N_{E/S'} \longrightarrow 0$$

Of course $T_E \otimes_{\mathcal{O}_E} N_{E/S'}$ is a locally free \mathcal{O}_E -module; in particular, we have a canonical isomorphism,

$$\mathrm{Tor}_1^{\mathcal{O}_{\Sigma'}}(\mathcal{O}_{\Sigma'}, T_E \otimes_{\mathcal{O}_E} N_{E/S'}) \cong \mathrm{Tor}_1^{\mathcal{O}_{S'}}(\mathcal{O}_{\Sigma'}, \mathcal{O}_E) \otimes_{\mathcal{O}_E} T_E \otimes_{\mathcal{O}_E} N_{E/S'}.$$

Tensoring the short exact sequence above with $\mathcal{O}_{\Sigma'}$ over $\mathcal{O}_{S'}$ and using Lemma 3.12, produces the exact sequence,

$$0 \rightarrow T_E(-D) \rightarrow (i')^* T_{S'} \xrightarrow{dv} u^* i^* T_S \rightarrow T_E \otimes_{\mathcal{O}_E} N_{E/S'} \rightarrow 0.$$

□

The maps $N_{(\zeta', i')} \rightarrow u^* N_{(\zeta, i)}$ and $(i')^* T_{S'} \rightarrow u^* i^* T_S$ are compatible with $\alpha_{(\zeta', i')}^\dagger$ and $u^* \alpha_{(\zeta, i)}^\dagger$. So there is an induced map of complexes $L_{(\zeta', i')}^\vee \rightarrow u^* L_{(\zeta, i)}^\vee$.

- Notation 3.14.** (i) Denote by $du : L_{(\zeta', i')}^\vee \rightarrow u^* L_{(\zeta, i)}^\vee$ the induced map of complexes.
(ii) Denote by $\mathrm{Image}(du) \hookrightarrow u^* L_{(\zeta, i)}^\vee$ the image of du in the Abelian category of complexes of coherent $\mathcal{O}_{\Sigma'}$ -modules.
(iii) If v is Type I, denote $K_I = T_E(-D)[0]$ and $Q_I = (T_E \otimes_{\mathcal{O}_E} N_{E/S'})[0]$.
(iv) If v is Type II, denote

$$\begin{aligned} K_{II} &= T_E(-D)[0] \oplus \left(M \otimes_{\mathbb{C}} N_{E/S'}^\vee(-D) \right) [-1] \\ Q_{II} &= (T_E \otimes_{\mathcal{O}_E} N_{E/S'})[0] \oplus (M \otimes_{\mathbb{C}} \mathcal{O}_E) [-1] \end{aligned}$$

Lemma 3.15. *Both if v is Type I and if v is Type II, there are short exact sequences of complexes of coherent $\mathcal{O}_{\Sigma'}$ -modules,*

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K & \longrightarrow & L_{(\zeta', i')}^\vee & \xrightarrow{du} & \mathrm{Image}(du) & \longrightarrow & 0 \\ 0 & \longrightarrow & \mathrm{Image}(du) & \longrightarrow & u^* L_{(\zeta, i)}^\vee & \longrightarrow & Q & \longrightarrow & 0 \end{array}$$

Proof. If v is Type I, there is a commutative diagram with exact rows,

$$\begin{array}{ccccccccc} & & L_{(\zeta', i')}^\vee & \xrightarrow{du} & u^* L_{(\zeta, i)}^\vee & & & & \\ 0 & \rightarrow & T_E(-D) & \rightarrow & (i')^* T_{S'} & \xrightarrow{(i')^* dv} & u^* i^* T_S & \rightarrow & T_E \otimes_{\mathcal{O}_E} N_{E/S'} \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ & & N_{(\zeta', i')} & \cong & h^* N_{(\zeta, i)} & & & & \end{array}$$

The middle two columns of this diagram give $du : L_{(\zeta', i')}^\vee \rightarrow u^* L_{(\zeta, i)}^\vee$. By inspection the kernel of du is K_I and the cokernel of du is Q_I .

If v is Type II, there is a commutative diagram with exact rows,

$$\begin{array}{ccccccccc} & & L_{(\zeta', i')}^\vee & \xrightarrow{du} & u^* L_{(\zeta, i)}^\vee & & & & \\ 0 & \rightarrow & T_E(-D) & \rightarrow & (i')^* T_{S'} & \xrightarrow{(i')^* dv} & u^* i^* T_S & \rightarrow & T_E \otimes_{\mathcal{O}_E} N_{E/S'} \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & M \otimes_{\mathbb{C}} N_{E/S'}^\vee(-D) & \rightarrow & N_{(\zeta', i')} & \rightarrow & u^* N_{(\zeta, i)} & \rightarrow & M \otimes_{\mathbb{C}} \mathcal{O}_E \rightarrow 0 \end{array}$$

The middle two columns of this diagram give $du : L_{(\zeta', i')}^\vee \rightarrow u^* L_{(\zeta, i)}^\vee$.

The claim is that the induced map of kernels, $T_E(-D) \rightarrow M \otimes_{\mathbb{C}} N_{E/S'}^\vee(-D)$, is zero, and the induced map of cokernels, $T_E \otimes_{\mathcal{O}_E} N_{E/S'} \rightarrow M \otimes_{\mathbb{C}} \mathcal{O}_E$ is zero. To see this, observe that in each

map, the domain and target are locally free sheaves on E . So each map is really a section of a $\text{Hom}_{\mathcal{O}_E}(\cdot, \cdot)$ sheaf. Up to canonical isomorphisms, the sheaf in both cases is simply

$$\text{Hom}_{\mathcal{O}_E}(T_E, M \otimes_{\mathbb{C}} N_{E/S'}^\vee) \cong M \otimes_{\mathbb{C}} \Omega_{S'}^2 \otimes_{\mathcal{O}_S} \mathcal{O}_E.$$

Identifying E with \mathbb{P}^1 , there are isomorphisms $N_{E/S'} \cong \mathcal{O}_E(-1)$ and $\mathcal{O}_E(K_E) \cong \mathcal{O}_E(-2)$. By the adjunction formula for divisors on surfaces $\Omega_{S'}^2 \otimes_{\mathcal{O}_S} \mathcal{O}_E = \mathcal{O}_{S'}(K_{S'}) \otimes_{\mathcal{O}_S} \mathcal{O}_E$ is isomorphic to $\mathcal{O}_E(K_E) \otimes_{\mathcal{O}_E} N_{E/S'}^\vee$, i.e., $\Omega_{S'}^2 \otimes_{\mathcal{O}_S} \mathcal{O}_E \cong \mathcal{O}_E(-2+1) = \mathcal{O}_E(-1)$. Since this sheaf has no nonzero global sections, both the induced maps are zero. Therefore the kernel of du is K_{II} and cokernel of du is Q_{II} . \square

Assume first that v is Type I. Since $T_E(-D) \cong \mathcal{O}_E(1)$, $\mathbb{H}^0(\Sigma', K_I) = H^0(E, T_E(-D))$ is 2-dimensional, and $\dim \mathbb{H}^i(\Sigma', K_I) = 0$ for $i \neq 0$. Similarly $\mathbb{H}^0(\Sigma', Q_I) = H^0(E, T_E(E))$ is 2-dimensional and $\dim \mathbb{H}^i(\Sigma', Q_I) = 0$ for $i > 0$. Therefore there is a long exact sequence of hypercohomology groups:

$$\begin{aligned} 0 \rightarrow H^0(E, T_E(-D)) &\rightarrow \mathbb{H}^0(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^0(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow \dots \\ \dots \rightarrow H^0(E, T_E \otimes_{\mathcal{O}_E} N_{E/S'}) &\rightarrow \mathbb{H}^1(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^1(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0 \\ 0 \rightarrow \mathbb{H}^2(\Sigma', L_{(\zeta', i')}^\vee) &\rightarrow \mathbb{H}^2(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0 \end{aligned}$$

Lemma 3.16. *If v is Type Ia then there are exact sequences,*

$$\begin{aligned} 0 \rightarrow H^0(E, T_E(-D)) &\rightarrow \mathbb{H}^0(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^0(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0, \\ 0 \rightarrow H^0(E, T_E \otimes_{\mathcal{O}_E} N_{E/S'}) &\rightarrow \mathbb{H}^1(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^1(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0, \\ 0 \rightarrow \mathbb{H}^2(\Sigma', L_{(\zeta', i')}^\vee) &\rightarrow \mathbb{H}^2(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0. \end{aligned}$$

In other words, the canonical map from the space of infinitesimal automorphisms of ζ' to the space of infinitesimal automorphisms of ζ is surjective with 2-dimensional kernel, the canonical map from the space of first order deformations of ζ' to the space of first order deformations of ζ is surjective with 2-dimensional kernel, and the obstruction space of ζ' equals the obstruction space of ζ .

Proof. The only claim that doesn't follow from the long exact sequence of cohomology is that $\mathbb{H}^0(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^0(\Sigma, L_{(\zeta, i)}^\vee)$ is surjective. Since the irreducible component $\Sigma_j \subset \Sigma$ containing s is stable, Lemma 3.5 states that every infinitesimal automorphism of ζ vanishes on Σ_j . The infinitesimal automorphisms of ζ that vanish at s are the same as the infinitesimal automorphisms of ζ' that vanish on E . Therefore every infinitesimal automorphism of ζ is the image of an infinitesimal automorphism of ζ' that vanishes on E . \square

Let v be a morphism of Type Ib. Let $\Sigma_j \subset \Sigma$ denote the unstable component containing s . Let N denote the 1-dimensional vector space $\Theta_{\Sigma_j} \otimes_{\mathcal{O}_{\Sigma_j}} \mathcal{O}_s$.

Lemma 3.17. *If v is Type Ib then there are exact sequences,*

$$\begin{aligned} 0 \rightarrow H^0(E, T_E(-D)) &\rightarrow \mathbb{H}^0(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^0(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow N \rightarrow 0 \\ 0 \rightarrow H^0(E, T_E \otimes_{\mathcal{O}_E} N_{E/S'})/N &\rightarrow \mathbb{H}^1(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^1(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0 \\ 0 \rightarrow \mathbb{H}^2(\Sigma', L_{(\zeta', i')}^\vee) &\rightarrow \mathbb{H}^2(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0 \end{aligned}$$

In other words, the canonical map from the space of infinitesimal automorphisms of ζ' to the space of infinitesimal automorphisms of ζ has a 2-dimensional kernel and a 1-dimensional cokernel, the canonical map from the space of first order deformations of ζ' to the space of first order deformations of ζ is surjective with 1-dimensional kernel, and the obstruction space of ζ' equals the obstruction space of ζ .

Proof. As in Lemma 3.16, the only claim that doesn't follow from the long exact sequence of cohomology is that the cokernel of $du : \mathbb{H}^0(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^0(\Sigma, L_{(\zeta, i)}^\vee)$ is N . Let $\Sigma'_j \subset \Sigma'$ denote the strict transform of Σ_j ; Σ'_j is canonically isomorphic to Σ_j . Composing du with the restriction map $\mathbb{H}^0(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow H^0(\Sigma_j, \Theta_{\Sigma_j})$ gives a map which is canonically isomorphic to the restriction map $\mathbb{H}^0(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow H^0(\Sigma'_j, \Theta_{\Sigma'_j})$. By Lemma 3.5, every infinitesimal automorphism of ζ' restricts to an infinitesimal automorphism of Σ'_j that vanishes at s . On the other hand, the infinitesimal automorphisms of ζ that vanish at s are the same as the infinitesimal automorphisms of ζ' that vanish on E . Hence the image of du is precisely the subspace of infinitesimal automorphisms of ζ that vanish at s .

Since Σ_j is unstable, either $\Sigma = \Sigma_j$ is a curve of arithmetic genus 1 with no markings and g is constant, or Σ_j is a smooth curve of genus 0 containing at most 2 marked points and nodes of Σ and $g|_{\Sigma_j}$ is constant. In each case, it is easy to see that there is an infinitesimal automorphism of Σ_j that vanishes at all marked points and nodes of Σ and that does not vanish at s . By Lemma 3.5, this infinitesimal automorphism of Σ_j is the image of an infinitesimal automorphism of ζ , i.e., there exists an infinitesimal automorphism of ζ that does not vanish at s . Therefore the cokernel of du is N . \square

Assume next that v is Type II. Because the divisor $D \subset E$ has degree 2, the \mathcal{O}_E -module $N_{E/S'}^\vee(-D)$ is isomorphic to $\mathcal{O}_E(-1)$. Since $h^0(E, \mathcal{O}_E(-1)) = h^1(E, \mathcal{O}_E(-1)) = 0$, the term $M \otimes_{\mathbb{C}} N_{E/S'}^\vee(-D)[-1]$ in K_{II} does not contribute to the hypercohomology; i.e., the hypercohomology of K_{II} is the sheaf cohomology of $T_E(-D)$. The \mathcal{O}_E -module $T_E(-D)$ is isomorphic to \mathcal{O}_E ; hence $h^0(E, T_E(-D)) = 1$ and $h^1(E, T_E(-D)) = 0$. Therefore $\dim \mathbb{H}^0(\Sigma', K_{II}) = H^0(E, T_E(-D))$ is 1-dimensional, and $\dim \mathbb{H}^i(\Sigma', K_{II}) = 0$ for $k \neq 1$.

For Q_{II} both $T_E \otimes_{\mathcal{O}_E} N_{E/S'}[0]$ and $M \otimes_{\mathbb{C}} \mathcal{O}_E[-1]$ contribute to the hypercohomology. The \mathcal{O}_E -module $T_E \otimes_{\mathcal{O}_E} N_{E/S'}$ is isomorphic to $\mathcal{O}_E(1)$. Hence $\mathbb{H}^0(\Sigma', Q_{II}) = H^0(E, T_E(E))$ is 2-dimensional, $\mathbb{H}^1(\Sigma', Q_{II}) = M$ is 1-dimensional (recall $M = \mathcal{O}_S(\Sigma)|_s$), and $\dim \mathbb{H}^k(\Sigma', Q_{II}) = 0$ for $k \neq 0, 1$. Therefore there is a long exact sequence in hypercohomology:

$$\begin{aligned} 0 \rightarrow H^0(E, T_E(-D)) \rightarrow \mathbb{H}^0(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^0(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow \dots \\ \dots \rightarrow H^0(E, T_E \otimes_{\mathcal{O}_E} N_{E/S'}) \rightarrow \mathbb{H}^1(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^1(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow \dots \\ \dots \rightarrow M \rightarrow \mathbb{H}^2(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^2(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0 \end{aligned}$$

Every infinitesimal automorphism of ζ vanishes at s . So the infinitesimal automorphisms of ζ are the same as the infinitesimal automorphisms of ζ' that vanish on E . In particular, $\mathbb{H}^0(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^0(\Sigma, L_{(\zeta, i)}^\vee)$ is surjective. The map $\mathbb{H}^1(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow \mathcal{O}_S(\Sigma)|_s$ is nonzero iff there are deformations of ζ that smooth the node s to first order. This proves the following two lemmas, which are stated separately for notational convenience.

Lemma 3.18. *If v is Type IIa then there are exact sequences,*

$$0 \rightarrow H^0(E, T_E(-D)) \rightarrow \mathbb{H}^0(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^0(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0 \quad (2)$$

$$0 \rightarrow H^0(E, T_E \otimes_{\mathcal{O}_E} N_{E/S'}) \rightarrow \mathbb{H}^1(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^1(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow \mathcal{O}_S(\Sigma)|_s \rightarrow 0 \quad (3)$$

$$0 \rightarrow \mathbb{H}^2(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^2(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0 \quad (4)$$

In other words, the canonical map from the space of infinitesimal automorphisms of ζ' to the space of infinitesimal automorphisms of ζ is surjective with a 1-dimensional kernel, the canonical map

from the space of first order deformations of ζ' to the space of first order deformations of ζ has a 2-dimensional kernel and a 1-dimensional cokernel, and the obstruction space to ζ' equals the obstruction space to ζ .

Lemma 3.19. *If v is Type IIb then there are exact sequences,*

$$0 \rightarrow H^0(E, T_E(-D)) \rightarrow \mathbb{H}^0(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^0(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0 \quad (5)$$

$$0 \rightarrow H^0(E, T_E \otimes_{\mathcal{O}_E} N_{E/S'}) \rightarrow \mathbb{H}^1(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^1(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0 \quad (6)$$

$$0 \rightarrow \mathcal{O}_S(\Sigma)|_s \rightarrow \mathbb{H}^2(\Sigma', L_{(\zeta', i')}^\vee) \rightarrow \mathbb{H}^2(\Sigma, L_{(\zeta, i)}^\vee) \rightarrow 0. \quad (7)$$

In other words, the canonical map from the space of infinitesimal automorphisms of ζ' to the space of infinitesimal automorphisms of ζ is surjective with a 1-dimensional kernel, the canonical map from the space of first order deformations of ζ' to the space of first order deformations of ζ is surjective with 1-dimensional kernel, and the map from the obstruction space of ζ' to the obstruction space of ζ is surjective and has a 1-dimensional kernel.

Finally, assume that $u : \Sigma' \rightarrow \Sigma$ is a Type III elementary morphism, i.e., u is the identity map, but there is one marked point $\sigma' \in \Sigma'$ that is not in Σ .

Definition 3.20. Let $u : \Sigma' \rightarrow \Sigma$ be a Type III elementary morphism.

- (i) The morphism u is *Type IIIa* if $\sigma' \in \Sigma$ lies on an unstable component.
- (ii) The morphism u is *Type IIIb* if $\sigma' \in \Sigma$ lies on a stable component.

In both cases, there is a canonical short exact sequence of complexes:

$$0 \longrightarrow L_\zeta \longrightarrow L_{\zeta'} \longrightarrow \Omega_\Sigma(\sigma') \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\sigma'}[0] \longrightarrow 0$$

Of course $\mathbb{E}xt_{\mathcal{O}_\Sigma}^1(\Omega_\Sigma(\sigma') \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\sigma'}, \mathcal{O}_\Sigma)$ is canonically isomorphic to $T_\Sigma \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\sigma'}$, and $\dim \mathbb{E}xt_{\mathcal{O}_\Sigma}^k(\Omega_\Sigma(\sigma') \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\sigma'}, \mathcal{O}_\Sigma) = 0$ for $k \neq 1$. In particular, there is an induced map

$$\mathbb{E}xt_{\mathcal{O}_\Sigma}^0(L_\zeta, \mathcal{O}_\Sigma) \rightarrow T_\Sigma \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\sigma'}.$$

This map is zero iff σ' lies on a stable component of ζ . Combined with the long exact sequence of hypercohomology associated to the short exact sequences, this proves the following two lemmas, which are stated separately for notational convenience.

Lemma 3.21. *If $u : \Sigma' \rightarrow \Sigma$ is Type IIIa then there are exact sequences,*

$$0 \rightarrow \mathbb{E}xt_{\mathcal{O}_{\Sigma'}}^0(L_{\zeta'}, \mathcal{O}_{\Sigma'}) \rightarrow \mathbb{E}xt_{\mathcal{O}_\Sigma}^0(L_\zeta, \mathcal{O}_\Sigma) \rightarrow T_\Sigma \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\sigma'} \rightarrow 0$$

$$0 \rightarrow \mathbb{E}xt_{\mathcal{O}_{\Sigma'}}^1(L_{\zeta'}, \mathcal{O}_{\Sigma'}) \rightarrow \mathbb{E}xt_{\mathcal{O}_\Sigma}^1(L_\zeta, \mathcal{O}_\Sigma) \rightarrow 0$$

$$0 \rightarrow \mathbb{E}xt_{\mathcal{O}_{\Sigma'}}^2(L_{\zeta'}, \mathcal{O}_{\Sigma'}) \rightarrow \mathbb{E}xt_{\mathcal{O}_\Sigma}^2(L_\zeta, \mathcal{O}_\Sigma) \rightarrow 0$$

In other words, the space of infinitesimal automorphisms of ζ' maps isomorphically to a codimension 1 linear subspace of the space of infinitesimal automorphisms of ζ , the space of first order deformations of ζ' equals the space of first order deformations of ζ , and the obstruction space of ζ' equals the obstruction space of ζ .

Lemma 3.22. *If $u : \Sigma' \rightarrow \Sigma$ is Type IIIb then there are exact sequences,*

$$0 \rightarrow \mathbb{E}xt_{\mathcal{O}_{\Sigma'}}^0(L_{\zeta'}, \mathcal{O}_{\Sigma'}) \rightarrow \mathbb{E}xt_{\mathcal{O}_\Sigma}^0(L_\zeta, \mathcal{O}_\Sigma) \rightarrow 0$$

$$0 \rightarrow T_\Sigma \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\sigma'} \rightarrow \mathbb{E}xt_{\mathcal{O}_{\Sigma'}}^1(L_{\zeta'}, \mathcal{O}_{\Sigma'}) \rightarrow \mathbb{E}xt_{\mathcal{O}_\Sigma}^1(L_\zeta, \mathcal{O}_\Sigma) \rightarrow 0$$

$$0 \rightarrow \mathbb{E}xt_{\mathcal{O}_{\Sigma'}}^2(L_{\zeta'}, \mathcal{O}_{\Sigma'}) \rightarrow \mathbb{E}xt_{\mathcal{O}_\Sigma}^2(L_\zeta, \mathcal{O}_\Sigma) \rightarrow 0$$

In other words, the space of infinitesimal automorphisms of ζ' equals the space of infinitesimal automorphisms of ζ , the canonical map from the space of first order deformations of ζ' to the space of first order deformations of ζ is surjective with 1-dimensional kernel, and the obstruction space of ζ' equals the obstruction space of ζ .

Taken together, Lemma 3.16 through Lemma 3.22, describe the canonical maps $\mathbb{E}xt_{\mathcal{O}_{\Sigma'}}^k(L_{\zeta'}, \mathcal{O}_{\Sigma'}) \rightarrow \mathbb{E}xt_{\mathcal{O}_{\Sigma}}^k(L_{\zeta}, \mathcal{O}_{\Sigma})$ for any morphism $u : \Sigma' \rightarrow \Sigma$ that removes a subset of marked points from Σ' and then contracts a subset of the unstable components.

3.2. Gluing stable maps. Let τ be a stable A -graph, not necessarily of genus 0. To describe the analogue of Lemma 3.3, a more precise description of the stack $\overline{\mathcal{M}}(X, \tau)$ is needed. Let B be a scheme. A 1-morphism $\tilde{\zeta} : B \rightarrow \overline{\mathcal{M}}(X, \tau)$ is equivalent to a pair (ζ, ϕ) consisting of a family of r -pointed stable maps,

$$\zeta = ((p : \Sigma \rightarrow B, \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X),$$

along with a natural assignment to each geometric point $b \in B$ of a contraction of stable A -graphs $\phi_b : \tau(\zeta_b) \rightarrow \tau$. The graph $\tau(\zeta_b)$ is the stable A -graph of $((\Sigma_b, \sigma_1(b), \dots, \sigma_r(b)), g_b)$. A contraction of stable A -graphs, $\phi : \tau' \rightarrow \tau$ is a map that contracts subgraphs of τ' to vertices of τ , cf. [5, Def. 1.8]. Geometrically the vertices of τ give a decomposition of Σ_b into connected subcurves. The main example of a contraction of stable A -graphs comes from a family of stable maps over a discrete valuation ring, $\zeta : \text{Spec } R \rightarrow \overline{\mathcal{M}}_{g,r}(X, e)$. Let ζ_0 denote the fiber over the geometric closed point of R and let ζ_η denote the fiber over the geometric generic point of R . There is a canonical contraction $\phi_{(\zeta, R)} : \tau(\zeta_0) \rightarrow \tau(\zeta_\eta)$; a vertex of $\tau(\zeta_0)$ corresponding to an irreducible component $\Sigma_{0,j} \subset \Sigma_0$ maps to a vertex of $\tau(\zeta_\eta)$ corresponding to an irreducible component of $\Sigma_{\eta,k} \subset \Sigma_\eta$ iff $\Sigma_{0,j}$ is in the closure of $\Sigma_{\eta,k}$. The assignment ϕ from above is called *natural* if it is compatible with the action of the Galois group of $\kappa(b)$ and for each map from a discrete valuation ring to B , $\rho : \text{Spec } R \rightarrow B$, the contractions $\phi_{\rho(0)}$ and $\phi_{\rho(\eta)}$ commute with the contraction $\phi_{\rho^*\zeta, R}$.

Notation 3.23. Associated to each edge $\epsilon = \{f_1, f_2\}$ of τ , there is a section $\sigma_\epsilon : B \rightarrow \Sigma$ such that for each geometric point $b \in B$, $\sigma_\epsilon(b) \in \Sigma_b$ is a node. Denote by N_ϵ the pullback $\sigma_\epsilon^* \text{Ext}_{\mathcal{O}_\Sigma}^1(\Omega_p(\sigma_1(B) + \dots + \sigma_r(B)), \mathcal{O}_\Sigma)$. Denote by N_τ the direct sum $\bigoplus_\epsilon N_\epsilon$ where ϵ ranges over all edges of τ .

Definition 3.24. Let T be a scheme. Let C be a complex of coherent sheaves on T and let n be an integer. The *good* ($\leq r$)-*truncation*, $C^{\leq r}$, is the complex of coherent sheaves on T ,

$$(C^{\leq r})^k = \begin{cases} C^k & k < r, \\ \text{Ker}(d^r : C^r \rightarrow C^{r+1}) & k = r, \\ \{0\} & k > r \end{cases}$$

The differentials on $C^{\leq r}$ are the obvious ones. The association $C \mapsto C^{\leq r}$ defines a functor on the category of complexes of coherent sheaves on T . This functor takes quasi-isomorphisms to quasi-isomorphisms and sends null-homotopic maps to null-homotopic maps, thus it induces a well-defined functor on the derived category of coherent sheaves on T . There is a natural transformation to the identity functor, $C^{\leq r} \rightarrow C$.

The \mathcal{O}_B -module N_ϵ is invertible and the localization of $\text{Ext}_{\mathcal{O}_\Sigma}^1(\Omega_p(\sigma_1(B) + \dots + \sigma_r(B)), \mathcal{O}_\Sigma)$ along $\sigma_\epsilon(B)$ is canonically isomorphic to $(\sigma_\epsilon)_* N_\epsilon$. By construction there is a map of complexes of coherent \mathcal{O}_Σ -modules, $L_\zeta^\vee \rightarrow \mathbb{R} \text{Hom}_{\mathcal{O}_\Sigma}(\Omega_p(\sigma_1(B) + \dots + \sigma_r(B)), \mathcal{O}_\Sigma)$. For all $k > 1$, $\text{Ext}_{\mathcal{O}_B}^k(\Omega_B(\sigma_1(B) + \dots + \sigma_r(B)), \mathcal{O}_B) = \{0\}$. So the complex is quasi-isomorphic to its (≤ 1)-good truncation,

$$\mathbb{R}^{\leq 1} \text{Hom}_{\mathcal{O}_\Sigma}(\Omega_p(\sigma_1(B) + \dots + \sigma_r(B)), \mathcal{O}_\Sigma) \xrightarrow{\text{qisom}} \mathbb{R} \text{Hom}_{\mathcal{O}_\Sigma}(\Omega_p(\sigma_1(B) + \dots + \sigma_r(B)), \mathcal{O}_\Sigma).$$

Notation 3.25. After replacing L_ζ^\vee by a quasi-isomorphic complex, there is a morphism of complexes denoted $a_{(\zeta,\tau)} : L_\zeta^\vee \rightarrow \mathbb{R}^{\leq 1} \text{Hom}_{\mathcal{O}_\Sigma}(\Omega_p(\sigma_1(B) + \cdots + \sigma_r(B)), \mathcal{O}_\Sigma)$ factoring the original morphism. There is a map of complexes of \mathcal{O}_Σ -modules denoted

$$b_{(\zeta,\tau)} : \mathbb{R}^{\leq 1} \text{Hom}_{\mathcal{O}_\Sigma}(\Omega_p(\sigma_1(B) + \cdots + \sigma_r(B)), \mathcal{O}_\Sigma) \rightarrow \text{Ext}_{\mathcal{O}_\Sigma}^1(\Omega_p(\sigma_1(B) + \cdots + \sigma_r(B)), \mathcal{O}_\Sigma)[-1].$$

Denote by,

$$c_{(\zeta,\tau)} : \text{Ext}_{\mathcal{O}_\Sigma}^1(\Omega_p(\sigma_1(B) + \cdots + \sigma_r(B)), \mathcal{O}_\Sigma)_\tau \rightarrow \text{Ext}_{\mathcal{O}_\Sigma}^1(\Omega_p(\sigma_1(B) + \cdots + \sigma_r(B)), \mathcal{O}_\Sigma),$$

the kernel of,

$$\text{Ext}_{\mathcal{O}_\Sigma}^1(\Omega_p(\sigma_1(B) + \cdots + \sigma_r(B)), \mathcal{O}_\Sigma) \rightarrow \oplus_\epsilon(\sigma_\epsilon)_* N_\epsilon.$$

Denote by

$$d_{(\zeta,\tau)} : \mathbb{R}^{\leq 1} \text{Hom}_{\mathcal{O}_\Sigma}(\Omega_p(\sigma_1(B) + \cdots + \sigma_r(B)), \mathcal{O}_\Sigma)_\tau \rightarrow \mathbb{R}^{\leq 1} \text{Hom}_{\mathcal{O}_\Sigma}(\Omega_p(\sigma_1(B) + \cdots + \sigma_r(B)), \mathcal{O}_\Sigma),$$

the fiber product of $b_{(\zeta,\tau)}$ and $c_{(\zeta,\tau)}$. Denote by,

$$e_{(\zeta,\tau)} : L_{(\zeta,\tau)}^\vee \rightarrow L_\zeta^\vee,$$

the fiber product of $a_{(\zeta,\tau)}$ and $d_{(\zeta,\tau)}$.

Let $|\tau|$ denote the underlying *modular graph* of τ , i.e., the graph of τ along with the genus function on vertices, but without the degree function on vertices, cf. [5]. There is a (highly non-separated) Artin stack of $|\tau|$ -prestable curves, $\mathfrak{M}(|\tau|)$, and a 1-morphism $\overline{\mathcal{M}}(X, \tau) \rightarrow \mathfrak{M}(|\tau|)$. The *relative obstruction theory* for this 1-morphism is described in [3]. From this the absolute obstruction theory of $\overline{\mathcal{M}}(X, \tau)$ readily follows.

Lemma 3.26. *Let X be a smooth quasi-projective scheme. Let τ be a stable A -graph. Let*

$$\zeta = ((p : \Sigma \rightarrow \overline{\mathcal{M}}(X, \tau), \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X)$$

denote the universal family of stable maps over $\overline{\mathcal{M}}(X, \tau)$. There exists an obstruction theory for $\overline{\mathcal{M}}(X, \tau)$ in the sense of [4, Def. 4.4] of the form

$$\left(\mathbb{R}p_*(L_{(\zeta,\tau)}^\vee)[1] \right)^\vee \rightarrow L_{\overline{\mathcal{M}}(X,\tau)}.$$

This obstruction theory is perfect, and there is a distinguished triangle in the derived category of complexes of coherent $\mathcal{O}_{\overline{\mathcal{M}}(X,\tau)}$ -modules,

$$\mathbb{R}p_*(L_{(\zeta,\tau)}^\vee)[1] \longrightarrow \mathbb{R}p_*(L_\zeta^\vee)[1] \longrightarrow N_\tau \longrightarrow \mathbb{R}p_*(L_{(\zeta,\tau)}^\vee)[2].$$

Remark 3.27. The definition of $L_{(\zeta,\tau)}^\vee$ seems very complicated, but in fact it is quite simple. Let $((\Sigma, \sigma_1, \dots, \sigma_r), g)$ be a stable τ -map. Let $i : \Sigma \rightarrow S$ be a closed immersion from Σ to a smooth surface S . Let N_i denote the normal sheaf of i . The complex L_ζ^\vee is quasi-isomorphic to the complex $L_{(\zeta,i)}^\vee$. There is a surjective map $N_{(\zeta,i)} \rightarrow N_i(\sigma_1(B) + \cdots + \sigma_r(B))$. Define $N_{(\zeta,i,\tau)} \subset N_{(\zeta,i)}$ to be the subsheaf which is the inverse image of $\mathcal{I} \cdot N_i(\sigma_1(B) + \cdots + \sigma_r(B)) \subset N_i(\sigma_1(B) + \cdots + \sigma_r(B))$, where \mathcal{I} is the ideal sheaf of $\cup_\epsilon \sigma_\epsilon$. Then $e_{(\zeta,\tau)} : L_{(\zeta,\tau)}^\vee \rightarrow L_\zeta^\vee$ is quasi-isomorphic to the following map of complexes.

$$\begin{array}{ccc} & 0 & 1 \\ & & \\ L_{(\zeta,i,\tau)}^\vee : i^* T_S(\sigma_1(B) + \cdots + \sigma_r(B)) & \longrightarrow & N_{(\zeta,i,\tau)} \\ \downarrow e_{(\zeta,i,\tau)} & \quad \quad \quad \downarrow = & \downarrow \\ L_{(\zeta,i)}^\vee : i^* T_S(\sigma_1(B) + \cdots + \sigma_r(B)) & \longrightarrow & N_{(\zeta,i)} \end{array}$$

Let $\epsilon = \{f_1, f_2\}$ be a disconnecting edge of τ . Let $\tau_1 \subset \tau$ be the maximal connected subgraph which contains f_1 and not f_2 , and let $\tau_2 \subset \tau$ be the maximal connected subgraph which contains f_2 and not f_1 . There are “forgetful” 1-morphisms $F_i : \overline{\mathcal{M}}(X, \tau) \rightarrow \overline{\mathcal{M}}(X, \tau_i)$ for $i = 1, 2$. Let $\tilde{\zeta} = (\zeta, \phi)$ be a map in $\overline{\mathcal{M}}(X, \tau)$, where $\zeta = ((\Sigma, \sigma_1, \dots, \sigma_r), g)$. Let $\sigma_\epsilon \in \Sigma$ be the node corresponding to ϵ . Let $\tilde{\zeta}_i = (\zeta_i, \phi_i) \in \overline{\mathcal{M}}(X, \tau_i)$ be the image $F_i(\tilde{\zeta})$. Denote $\zeta_i = ((\Sigma_i, \sigma_{i,0}, \dots, \sigma_{i,r_i}), g_i)$ for $i = 1, 2$ where $\sigma_{i,0} \in \Sigma_i$ is the point corresponding to the flag f_i of τ_i , i.e., $\sigma_{i,0} = \sigma_\epsilon$.

Lemma 3.28. *If*

- (i) *the dimension of the obstruction group of $\overline{\mathcal{M}}(X, \tau_i)$ at $\tilde{\zeta}_i$ is 0, $i = 1, 2$, and*
- (ii) *the evaluation morphism $ev_{f_1} : \overline{\mathcal{M}}(X, \tau_1) \rightarrow X$ is smooth at $\tilde{\zeta}_1$,*

then the dimension of the obstruction group of $\overline{\mathcal{M}}(X, \tau)$ at $\tilde{\zeta}$ is 0, and there is a short exact sequence,

$$0 \longrightarrow \zeta_1^* T_{ev_{f_1}} \longrightarrow \zeta^* T_{\overline{\mathcal{M}}(X, \tau)} \longrightarrow \zeta_2^* T_{\overline{\mathcal{M}}(X, \tau_2)} \longrightarrow 0,$$

where $T_{ev_{f_1}}$ is the dual of the sheaf of relative differentials of ev_{f_1} .

Proof. The proof follows from the fact that $\overline{\mathcal{M}}(X, \tau)$ is an open substack of the 2-fibered product:

$$\overline{\mathcal{M}}(X, \tau_1) \times_{\text{ev}_{f_1}, X, \text{ev}_{f_2}} \overline{\mathcal{M}}(X, \tau_2).$$

□

Let $\phi : \tau \rightarrow \tau'$ be the minimal contraction of stable A -graphs that contracts the edge $\{f_1, f_2\}$. The induced 1-morphism,

$$\overline{\mathcal{M}}(X, \phi) : \overline{\mathcal{M}}(X, \tau) \rightarrow \overline{\mathcal{M}}(X, \tau'),$$

is unramified and the image has codimension at most 1. In some circumstances, this morphism is the normalization of a Cartier divisor.

Lemma 3.29. *If*

- (i) *τ has genus 0,*
- (ii) *the dimension of the obstruction group of $\overline{\mathcal{M}}(X, \tau_2)$ at $\tilde{\zeta}_2$ is 0, and*
- (iii) *the \mathcal{O}_{Σ_1} -module $g_1^* T_X$ is generated by global sections,*

then the dimension of the obstruction group of $\overline{\mathcal{M}}(X, \tau')$ at $\overline{\mathcal{M}}(X, \phi)(\tilde{\zeta})$ is 0, the irreducible component of $\overline{\mathcal{M}}(X, \tau)$ containing $\tilde{\zeta}$ maps to a Cartier divisor under $\overline{\mathcal{M}}(X, \phi)$, and there is a short exact sequence,

$$0 \longrightarrow T_{\overline{\mathcal{M}}(X, \tau)}|_{\tilde{\zeta}} \longrightarrow T_{\overline{\mathcal{M}}(X, \tau')}|_{\tilde{\zeta}} \longrightarrow \Theta_{\Sigma_1}|_{\sigma_{1,0}} \otimes \Theta_{\Sigma_2}|_{\sigma_{2,0}} \longrightarrow 0.$$

Proof. There is a short exact sequence of complexes,

$$0 \longrightarrow L_{\Sigma}(\sigma_1 + \dots + \sigma_r) \longrightarrow L_{\zeta} \longrightarrow g^* \Omega_X[1] \longrightarrow 0,$$

that gives rise to a long exact sequence in hypercohomology, part of which is,

$$\mathbb{H}^1(\Sigma, L_{\zeta}^{\vee}) \rightarrow \mathbb{H}^1(\Sigma, L_{\Sigma}^{\vee}(-(\sigma_1 + \dots + \sigma_r))) \rightarrow H^1(\Sigma, g^* T_X) \rightarrow \mathbb{H}^2(\Sigma, L_{\zeta}^{\vee}) \rightarrow 0.$$

The goal is to prove that $\dim \mathbb{H}^2(\Sigma, L_{\zeta}^{\vee}) = 0$, i.e., that $\mathbb{H}^1(\Sigma, L_{\Sigma}^{\vee}(-(\sigma_1 + \dots + \sigma_r))) \rightarrow H^1(\Sigma, g^* T_X)$ is surjective. After replacing L_{Σ} and L_{ζ} by quasi-isomorphic complexes of locally free \mathcal{O}_{Σ} -modules,

there is a diagram of distinguished triangles of coherent \mathcal{O}_B -modules,

$$\begin{array}{ccccc} L_\Sigma(\sigma_1 + \cdots + \sigma_r) & \rightarrow & L_\zeta & \rightarrow & \cdots \\ \downarrow & & \downarrow & & \\ L_\Sigma(\sigma_1 + \cdots + \sigma_r) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2} & \rightarrow & L_\zeta \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2} & \rightarrow & \cdots \\ \\ \cdots & \rightarrow & g^* \Omega_X[1] & \rightarrow & L_\Sigma(\sigma_1 + \cdots + \sigma_r)[-1] \\ & & \downarrow & & \downarrow \\ \cdots & \rightarrow & g^* \Omega_X \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2}[1] & \rightarrow & L_\Sigma(\sigma_1 + \cdots + \sigma_r) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2}[-1] \end{array},$$

that gives rise to a commutative diagram of exact sequences in hypercohomology,

$$\begin{array}{ccc} \mathbb{H}^1(\Sigma, L_\Sigma^\vee(-(\sigma_1 + \cdots + \sigma_r))) & \rightarrow & \mathbb{H}^1(\Sigma, g^* T_X) \\ \downarrow & & \downarrow \\ \mathbb{H}^1(\Sigma, L_\Sigma^\vee(-(\sigma_1 + \cdots + \sigma_r)) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2}) & \rightarrow & \mathbb{H}^1(\Sigma, g^* T_X \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2}) \end{array}.$$

To prove the top horizontal arrow is surjective, it suffices to prove

- (i) the left vertical arrow is surjective,
- (ii) the right vertical arrow is an isomorphism, and
- (iii) the bottom vertical arrow is surjective.

Let S be a smooth surface, and let $i : \Sigma \rightarrow S$ be an unramified morphism. Then $L_\Sigma^\vee(-(\sigma_1 + \cdots + \sigma_r))$ is represented in the derived category of coherent sheaves by the complex

$$0 \quad \quad \quad 1 \\ L_i^\vee(-(\sigma_1 + \cdots + \sigma_r)) : i^* T_S(-(\sigma_1 + \cdots + \sigma_r)) \xrightarrow{\beta_i^\dagger} N_i(-(\sigma_1 + \cdots + \sigma_r)).$$

Let K denote the kernel of β_i^\dagger and let Q denote the cokernel of β_i^\dagger . The sheaf K is torsion-free and is locally free of rank 1 on a dense open subset of Σ . The sheaf Q is torsion. There is a short exact sequence of complexes

$$0 \longrightarrow K[0] \longrightarrow L_i^\vee(-(\sigma_1 + \cdots + \sigma_r)) \longrightarrow Q[-1] \longrightarrow 0.$$

This gives rise to a long exact sequence in hypercohomology, part of which is

$$H^1(\Sigma, K) \longrightarrow \mathbb{H}^1(\Sigma, L_i^\vee(-(\sigma_1 + \cdots + \sigma_r))) \longrightarrow H^0(\Sigma, Q) \longrightarrow 0.$$

Let K' denote the kernel of $\beta_i^\dagger \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2}$ and let Q' denote the cokernel. There are induced maps of coherent sheaves on Σ , $K \rightarrow K'$ and $Q \rightarrow Q'$. These maps give rise to a commutative diagram with exact rows,

$$\begin{array}{ccccccc} H^1(\Sigma, K) & \longrightarrow & \mathbb{H}^1(\Sigma, L_i^\vee(-(\sigma_1 + \cdots + \sigma_r))) & \longrightarrow & H^0(\Sigma, Q) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ H^1(\Sigma, K') & \longrightarrow & \mathbb{H}^1(\Sigma, L_i^\vee(-(\sigma_1 + \cdots + \sigma_r)) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2}) & \longrightarrow & H^0(\Sigma, Q') & \longrightarrow & 0. \end{array}$$

Now $Q \rightarrow Q'$ is a surjective map of torsion sheaves, so the map on the right is surjective. And the cokernel of $K \rightarrow K'$ is torsion, and hence the map on the left is surjective. Therefore the map in the middle is surjective, i.e., (i) holds.

There is a short exact sequence of \mathcal{O}_Σ -modules,

$$0 \longrightarrow (g_1)^* T_X(-\sigma_{1,0}) \longrightarrow g^* T_X \longrightarrow g^* T_X \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2} \longrightarrow 0$$

By hypothesis $(g_1)^*T_X$ is generated by global sections, hence $h^1(\Sigma_1, (g_1)^*T_X(-\sigma_{1,0})) = 0$ by Lemma 2.3. Therefore $H^1(\Sigma, g^*T_X) \rightarrow H^1(\Sigma, g^*T_X \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2})$ is an isomorphism, i.e., (ii) is true.

Denote by $j : \Sigma_2 \rightarrow \Sigma$ the canonical closed immersion. There is a morphism of complexes of \mathcal{O}_Σ -modules,

$$\begin{array}{ccc} & -1 & 0 \\ L_\zeta : & g^*\Omega_X & \xrightarrow{(dg)^\dagger} \Omega_\Sigma(\sigma_1 + \cdots + \sigma_r) \\ & \downarrow & \downarrow (dj)^\dagger \\ j_*L_{\zeta_2} : & j_*(g_2)^*\Omega_X & \xrightarrow{j_*(dg_2)^\dagger} j_*\Omega_\Sigma(\sigma_{2,0} + \cdots + \sigma_{2,r_2}) \end{array}$$

By adjunction, there is a morphism of complexes of \mathcal{O}_{Σ_2} -modules, $L_\zeta \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2} \rightarrow L_{\zeta_2}$. So there is a commutative diagram in hypercohomology

$$\begin{array}{ccc} \mathbb{H}^1(\Sigma_2, L_{\Sigma_2}^\vee(-(\sigma_1 + \cdots + \sigma_r))) & \longrightarrow & H^1(\Sigma_2, (g_2)^*T_X) \\ \downarrow & & \downarrow \\ \mathbb{H}^1(\Sigma_2, L_{\Sigma_2}^\vee(-(\sigma_{2,0} + \cdots + \sigma_{2,r_2})) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2}) & \longrightarrow & H^1(\Sigma_2, j^*g^*T_X) \end{array}$$

Of course $(g_2)^*T_X = j^*g^*T_X$, so $H^1(\Sigma_2, (g_2)^*T_X) \rightarrow H^1(\Sigma_2, j^*g^*T_X)$ is an isomorphism. By hypothesis, the obstruction group of ζ_2 vanishes, hence $\mathbb{H}^1(\Sigma_2, L_{\Sigma_2}^\vee(-(\sigma_{2,0} + \cdots + \sigma_{2,r_2}))) \rightarrow H^1(\Sigma_2, (g_2)^*T_X)$ is surjective. Therefore also $\mathbb{H}^1(\Sigma_2, L_{\Sigma_2}^\vee(-(\sigma_1 + \cdots + \sigma_r)) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_{\Sigma_2}) \rightarrow H^1(\Sigma_2, j^*g^*T_X)$ is surjective. This proves (iii). Therefore the dimension of the obstruction group of $\overline{\mathcal{M}}(X, \tau)$ at $\overline{\mathcal{M}}(X, \phi)(\tilde{\zeta})$ is 0.

Because the obstruction groups of $\overline{\mathcal{M}}(X, \tau)$ and $\overline{\mathcal{M}}(X, \tau')$ both vanish, each stack is smooth of the expected dimension at $\tilde{\zeta}$. The expected dimension of $\overline{\mathcal{M}}(X, \tau)$ is 1 less than the expected dimension of $\overline{\mathcal{M}}(X, \tau')$, because τ has one extra edge (before stabilization). Therefore the image of the irreducible component of $\overline{\mathcal{M}}(X, \tau)$ containing ζ is a Cartier divisor in $\overline{\mathcal{M}}(X, \tau')$.

Finally, the short exact sequence follows from the 2-Cartesian diagram of Artin stacks,

$$\begin{array}{ccc} \overline{\mathcal{M}}(X, \tau) & \xrightarrow{\overline{\mathcal{M}}(X, \phi)} & \overline{\mathcal{M}}(X, \tau') \\ \downarrow & & \downarrow \\ \mathfrak{M}(|\tau|) & \xrightarrow{\mathfrak{M}(|\phi|)} & \mathfrak{M}(|\tau'|). \end{array}$$

The image of $\mathfrak{M}(|\phi|)$ is a Cartier divisor. Because the image of $\overline{\mathcal{M}}(X, \phi)$ is a Cartier divisor, the normal bundle of $\overline{\mathcal{M}}(X, \phi)$ is simply the pullback of the normal bundle of $\mathfrak{M}(|\phi|)$, which is the bundle whose fiber at $\tilde{\zeta}$ is the space of first-order deformations of the node $\sigma_\epsilon \in \Sigma$. It is well-known that this space is canonically isomorphic to $\Theta_{\Sigma_1}|_{\sigma_{1,0}} \otimes \Theta_{\Sigma_2}|_{\sigma_{2,0}}$. \square

Remark 3.30. There is a relative version of Lemma 3.29. Let B be a scheme and let $\tilde{\zeta} = (\zeta, \phi)$ be a family of τ -maps over B . If for every geometric point $b \in B$, the map $\tilde{\zeta}_b$ satisfies the hypotheses of Lemma 3.29, then the obstruction group of $\overline{\mathcal{M}}(X, \sigma)$ at $\tilde{\zeta}_b$ vanishes, the image of $\overline{\mathcal{M}}(X, \tau)$ in $\overline{\mathcal{M}}(X, \tau')$ is a Cartier divisor, and there exists a short exact sequence of locally free \mathcal{O}_B -modules,

$$0 \longrightarrow \tilde{\zeta}^*T_{\overline{\mathcal{M}}(X, \tau)} \longrightarrow \tilde{\zeta}^*T_{\overline{\mathcal{M}}(X, \tau')} \longrightarrow \sigma_{1,0}^*\Theta_{\Sigma_1/B} \otimes_{\mathcal{O}_B} \sigma_{2,0}^*\Theta_{\Sigma_2/B} \longrightarrow 0.$$

4. CONDITIONS ON FAMILIES OF STABLE MAPS

This section defines and proves basic results about the many conditions discussed in Subsection 1.2. Because of the process of *modification* given in the next section, it is necessary to work with families of stable maps that are parametrized by reducible curves of genus 0. This is the setting in which all definitions are made.

Let $\zeta = ((\Sigma, \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X)$ be an r -pointed genus 0 stable map to X .

Definition 4.1. The stable map ζ is *very stable* if the prestable map $(\Sigma, g : B \rightarrow X)$ is stable.

Let X be a quasi-projective scheme. Let B be a scheme. Let $\zeta : B \rightarrow \overline{\mathcal{M}}_{g,r}(X, e)$ be a 1-morphism.

Notation 4.2. The 1-morphism $\zeta : B \rightarrow \overline{\mathcal{M}}_{g,r}(X, e)$ consists of a datum,

$$\zeta = ((p_\zeta : \Sigma_\zeta \rightarrow B, \sigma_{(\zeta,1)}, \dots, \sigma_{(\zeta,r)}), g_\zeta : \Sigma \rightarrow X),$$

where

- (i) $p_\zeta : \Sigma_\zeta \rightarrow B$ is a proper, flat family of connected prestable curves of arithmetic genus g ,
- (ii) $\sigma_{(\zeta,i)} : B \rightarrow \Sigma_\zeta$, $i = 1, \dots, r$ is a collection of everywhere disjoint sections with image in the smooth locus of p_ζ , and
- (iii) $g_\zeta : \Sigma_\zeta \rightarrow X$ is a morphism of schemes,

that satisfies the stability condition in Definition 3.1. Denote by $h_{\zeta,i} : B \rightarrow X$ the composition $g_\zeta \circ \sigma_{(\zeta,i)}$. When there is no risk of confusion, the subscript ζ will be omitted.

Let X be a smooth, quasi-projective variety. Let $\pi : B \rightarrow T$ be a proper, flat family of connected prestable curves of arithmetic genus 0. Let $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ be a 1-morphism $((p : \Sigma \rightarrow B, \sigma), f : \Sigma \rightarrow X)$.

Definition 4.3. The 1-morphism $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ is *twisting* (resp. *very twisting*) if,

- (i) $(\pi : B \rightarrow T, h : B \rightarrow X)$ is a family of stable maps to X , i.e., a 1-morphism $\xi : T \rightarrow \overline{\mathcal{M}}_{0,0}(X, e)$ for some $e \geq 0$,
- (ii) the dimension of the obstruction group of $\overline{\mathcal{M}}_{0,0}(X, e)$ at each point of $\xi(T)$ is 0,
- (iii) the dimension of the relative obstruction group of the evaluation morphism $\text{ev} : \overline{\mathcal{M}}_{0,1}(X, 1) \rightarrow X$ at each point of $\zeta(T)$ is 0,
- (iv) denoting by T_{ev} the dual of the sheaf of relative differentials Ω_{ev} , the pullback ζ^*T_{ev} is π -relatively generated by global sections (resp. π -relatively deformation ample), and
- (v) denoting by $\text{pr} : \overline{\mathcal{M}}_{0,1}(X, 1) \rightarrow \overline{\mathcal{M}}_{0,0}(X, 1)$ the projection map, and denoting by T_{pr} the dual of the sheaf of relative differentials Ω_{pr} , the pullback ζ^*T_{pr} is π -relatively generated by global sections, i.e., the line bundle $\sigma^*\mathcal{O}_\Sigma(\sigma(B))$ is π -relatively generated by global sections.

Remark 4.4.

- (i) The sheaf ζ^*T_{pr} is canonically isomorphic to $\sigma^*\mathcal{O}_\Sigma(\sigma(B))$.
- (ii) The product morphism $(p, g) : \Sigma \rightarrow B \times X$ is a closed immersion whose ideal sheaf is everywhere locally defined by a regular sequence, i.e., it is a regular embedding. Denote by \mathcal{N} the normal bundle of this regular embedding. Then (iii) of Definition 4.3 is equivalent to the condition that $R^1p_*(\mathcal{N}(-\sigma(B))) = \{0\}$. Under this hypothesis, ζ^*T_{ev} is canonically isomorphic to $p_*(\mathcal{N}(-\sigma(B)))$.
- (iii) Condition (ii) is superfluous. Since the prestable family of maps $(\pi : B \rightarrow T, \xi : B \rightarrow X)$ is stable, also $(\pi : B \rightarrow T, \zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1))$ is stable by Lemma 3.6 (i).
- (iv) The conditions in Definition 4.3 impose some restrictions on the degrees of the locally free sheaves involved. By [19, Lem. 2.2.2], the total degree of $\sigma^*\mathcal{O}_\Sigma(\sigma(B))$ is simply $\delta = 2e - e'$

where e is the degree of $h : B \rightarrow \mathbb{P}^N$ and e' is the degree of $f : \Sigma \rightarrow \mathbb{P}^N$. If ζ is twisting then $2e \geq e'$.

Let B be a prestable, connected curve of arithmetic genus 0. Let $B_1, B_2 \subset B$ be connected subcurves of B such that $B_1 \cap B_2 = \{q\}$ is a single node, and $B = B_1 \cup B_2$. Let $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ be a 1-morphism.

Lemma 4.5. *If $\zeta|_{B_i} : B_i \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ is twisting for $i = 1, 2$, then ζ is twisting. If both ζ_1, ζ_2 are twisting, and if at least one of them is very twisting, then ζ is very twisting.*

Proof. This follows from Lemma 2.11. \square

Let $\pi : B \rightarrow T$ be a proper, flat family of connected, prestable curves of arithmetic genus 0. Let $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ be a 1-morphism. For every morphism of schemes $g : T' \rightarrow T$, denote by $\pi_{T'} : B_{T'} \rightarrow T'$ the base-change of π by g , and denote by $\zeta_{T'} : B_{T'} \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ the composition of $B_{T'} \rightarrow B$ with ζ .

Lemma 4.6. *There is an open subscheme $U_{twist} \subset T$ (resp. $U_{vtwist} \subset T$) with the following property: for every morphism of schemes $g : T' \rightarrow T$, the pullback family $\pi_{T'} : B_{T'} \rightarrow T'$ and $\zeta_{T'} : B_{T'} \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ is twisting (resp. very twisting) iff $g(T') \subset U_{twist}$ (resp. $g(T') \subset U_{vtwist}$).*

Proof. By [3, Lemma 1] there is an open subscheme $U_1 \subset T$ with the property that for every morphism g , $(\pi_{T'} : B_{T'} \rightarrow T', h_{T'} : B_{T'} \rightarrow X)$ is a family of stable maps iff $g(T') \subset U_1$. For every morphism g such that $\zeta_{T'}$ is twisting, $g(T') \subset U_1$. Hence the lemma for ζ_{U_1} implies the lemma for ζ . After replacing T by U_1 , the morphism $h : B \rightarrow X$ is a family of stable maps over T , i.e., a 1-morphism $\xi : T \rightarrow \overline{\mathcal{M}}_{0,0}(X, e)$ for some integer e .

Because π has relative dimension 1, the cohomology sheaf $\mathcal{H}^k(\mathbb{R}\pi_*(L_h^\vee)) = \{0\}$ for $k \geq 2$. By cohomology and base change, for every geometric point t of the support of the sheaf $\mathcal{H}^1(\mathbb{R}\pi_*(L_h^\vee))$, $\dim \mathbb{H}^1(B_t, L_{h_t}^\vee) > 0$, i.e. the obstruction group of ξ_t does not vanish. Therefore the dimension of the obstruction group of $\overline{\mathcal{M}}_{0,0}(X, e)$ at each point of $\xi_{T'}(T')$ is 0 iff $g(T')$ is contained in the complement of the support of $\mathcal{H}^1(\mathbb{R}\pi_*(L_h^\vee))$. The complement of the support of this sheaf is an open subset of T . After replacing T by this open set, the dimension of the obstruction group of $\overline{\mathcal{M}}_{0,0}(X, e)$ at each point of $\xi(T)$ is 0.

For similar reasons, T can be replaced by the complement of

$$\pi[\text{Supp}(R^1 p_*(\mathcal{N}(-\sigma(B))))],$$

where \mathcal{N} is as in (ii) of Remark 4.4. After replacing T by this open subset, for every g , $\zeta_{T'}$ satisfies (i), (ii), and (iii) of Definition 4.3. In order that (v) is satisfied, it is necessary and sufficient that $g(T')$ is contained in the complement of the support of the sheaf

$$\text{Coker}(\pi^* \pi_* \sigma^* \mathcal{O}_\Sigma(\sigma) \rightarrow \sigma^* \mathcal{O}_\Sigma(\sigma)).$$

After replacing T by the complement of the support of this sheaf, for every g , $\zeta_{T'}$ satisfies (i), (ii), (iii) and (v) of Definition 4.3.

Define U_{twist} to be the complement in T of the image under π of the support of

$$\text{Coker}(\pi^* \pi_* \zeta^* T_{ev} \rightarrow \zeta^* T_{ev}).$$

For every morphism g , $(\zeta^* T_{ev})_{T'}$ is $\pi_{T'}$ -relatively generated by global sections iff $g(T') \subset U_{twist}$. Therefore, for every morphism g , $\zeta_{T'}$ is twisting iff $g(T') \subset U_{twist}$. By Lemma 2.9, there exists an open subset $U_{vtwist} \subset T$ such that for every g , $(\zeta^* T_{ev})_{T'}$ is $\pi_{T'}$ -relatively deformation ample iff $g(T') \subset U_{vtwist}$. Therefore $\zeta_{T'}$ is very twisting iff $g(T') \subset U_{vtwist}$. \square

Let $(\pi : B \rightarrow T, h : B \rightarrow X)$ be a family of genus 0 stable maps, i.e., a 1-morphism $\xi : T \rightarrow \overline{\mathcal{M}}_{0,0}(X, e)$ for some $e \geq 0$. For every morphism of schemes $g : T' \rightarrow T$, denote by $\pi_{T'} : B_{T'} \rightarrow T'$ the base-change of π by g , and denote by $\xi_{T'} : B_{T'} \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ the composition of $B_{T'} \rightarrow B$ with ξ .

Definition 4.7. The family of stable maps $\xi : T \rightarrow \overline{\mathcal{M}}_{0,0}(X, e)$ is *twistable* (resp. *very twistable*) if there exists a surjective étale morphism $u : T' \rightarrow T$ and a morphism $\zeta : B_{T'} \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ such that $\xi_\zeta = \xi_{T'}$ and such that ζ is twisting (resp. very twisting).

Proposition 4.8. *There is an open subscheme $U_{t\text{-able}} \subset T$ (resp. $U_{vt\text{-able}} \subset T$) such that for every morphism of schemes $g : T' \rightarrow T$, $\xi_{T'}$ is twistable (resp. very twistable) iff $g(T') \subset U_{t\text{-able}}$ (resp. $g(T') \subset U_{vt\text{-able}}$).*

Proof. It suffices to check that if $t \in T$ is a geometric point such that $h_t : B_t \rightarrow X$ is twistable (resp. very twistable), then there is an étale neighborhood of $t \in T$ over which ξ is twistable (resp. very twistable). Denote by $\zeta_t : B_t \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ the twisting morphism. Consider $\overline{\mathcal{M}}_{0,1}(X, 1)$ as a quasi-projective scheme via the Plücker and Segre embeddings of $\mathbb{G}(1, n) \times \mathbb{P}^n \hookrightarrow \mathbb{P}^{\frac{n(n+1)^2}{2}-1}$. Let β denote the degree of the stable map ζ_t .

Define $\mathcal{M} = T \times \overline{\mathcal{M}}_{0,0}(\overline{\mathcal{M}}_{0,1}(X, 1), \beta)$, i.e., \mathcal{M} parametrizes pairs (s, ζ) consisting of a point $s \in T$ together with a genus 0 stable map $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ of degree β . Denote the universal stable map over $\overline{\mathcal{M}}_{0,0}(\overline{\mathcal{M}}_{0,1}(X, 1), \beta)$ by

$$\begin{aligned} \rho : \mathcal{B} &\rightarrow \overline{\mathcal{M}}_{0,0}(\overline{\mathcal{M}}_{0,1}(X, 1), \beta), \\ \zeta : \mathcal{B} &\rightarrow \overline{\mathcal{M}}_{0,1}(X, 1). \end{aligned}$$

As in Notation 4.2, let $p : \Sigma \rightarrow \mathcal{B}$ be the pullback by ζ of the universal curve over $\overline{\mathcal{M}}_{0,1}(X, 1)$, let $\sigma : \mathcal{B} \rightarrow \Sigma$ be the pullback of the universal section, let $g : \Sigma \rightarrow X$ be the pullback of the universal map, and let $h = g \circ \sigma$. This gives a family of prestable maps,

$$\tilde{\xi} = (\rho : \mathcal{B} \rightarrow \overline{\mathcal{M}}_{0,0}(\overline{\mathcal{M}}_{0,1}(X, 1), \beta), h : \mathcal{B} \rightarrow X). \quad (8)$$

By [3, Lemma 1] there is a maximal open substack $\mathcal{U}_e \subset \overline{\mathcal{M}}_{0,0}(\overline{\mathcal{M}}_{0,1}(X, 1), \beta)$ over which $\tilde{\xi}$ is stable of degree e . By hypothesis, (t, ζ_t) is in $T \times \mathcal{U}_e$.

Because ζ_t is twisting, the hypotheses of Lemma 3.6 (iii) are satisfied where $f : X \rightarrow Y$ corresponds to $\text{ev} : \overline{\mathcal{M}}_{0,1}(X, 1) \rightarrow X$ and where $(\Sigma, g : \Sigma \rightarrow X)$ corresponds to $(B_t, \zeta_t : B_t \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1))$. Therefore at the point $(B_t, \zeta_t : B_t \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1))$, the stack \mathcal{U}_e is smooth and the morphism $\tilde{\xi} : \mathcal{U}_e \rightarrow \overline{\mathcal{M}}_{0,0}(X, e)$ is smooth by the Jacobian criterion.

Consider the 1-morphism $(1_T, \xi) : T \rightarrow T \times \overline{\mathcal{M}}_{0,0}(X, e)$. Denote by \mathcal{M} the 2-fibered product of the 1-morphism $(1_T, \xi)$ and the 1-morphism $(1_T, \tilde{\xi}) : T \times \mathcal{U}_e \rightarrow T \times \overline{\mathcal{M}}_{0,0}(X, e)$. The 2-fibered product \mathcal{M} is the stack whose objects are triples (t, ζ, θ) consisting of a point $t \in T$, an object $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ of \mathcal{U}_e , and an equivalence $\theta : \xi_t \rightarrow \zeta$ of objects in the groupoid $\overline{\mathcal{M}}_{0,0}(X, e)(\text{Spec } \kappa(t))$. Because $(1_T, \tilde{\xi})$ is smooth at $(t, (B_t, \zeta_t))$, the projection $\text{pr}_1 : \mathcal{M} \rightarrow T$ is smooth at $(t, (B_t, \zeta_t))$. Hence there exists an étale 1-morphism $f : M \rightarrow \mathcal{M}$ from a scheme M to \mathcal{M} such that $(t, (B_t, \zeta_t))$ is in the image of f and such that the composite morphism $\text{pr}_1 \circ f : M \rightarrow T$ is smooth. By [6, Prop. 2.2.14], there exists an étale morphism $u : (T', t') \rightarrow (T, t)$ and a section $z : T' \rightarrow M$ such that $f(z(t'))$ is $(t, (B_t, \zeta_t))$. Denote by $\zeta_{\text{pre}} : T' \rightarrow \mathcal{U}_e$ the composition $\text{pr}_2 \circ f \circ z$.

Denote by $\zeta : B' \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ the pullback by $\zeta_{\text{pre}} : T' \rightarrow \mathcal{U}_e$ of the universal stable map. By construction, $\tilde{\xi}(\zeta) : B' \rightarrow \overline{\mathcal{M}}_{0,0}(X, e)$ is equivalent to $u^*\xi : u^*B \rightarrow \overline{\mathcal{M}}_{0,0}(X, e)$. Hence, after replacing T' by a surjective, étale cover, $B' = u^*B$ as T' -schemes, and $\tilde{\xi}(\zeta) = u^*\xi$. The fiber of $\zeta : u^*B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ over every preimage of $(t, (B_t, \zeta_t))$ is twisting. So by Lemma 4.6, after replacing T' by a Zariski open subscheme whose image contains t , $\zeta : u^*B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ is twisting. Similarly, if $(t, (B_t, \zeta_t))$ is very twisting, after replacing T' by a Zariski open subscheme whose image contains t , ζ is very twisting. On the Zariski open subscheme of T that is the image of $u : T' \rightarrow T$, the family $\xi : B \rightarrow \overline{\mathcal{M}}_{0,0}(X, e)$ is twistable (resp. very twistable). \square

Let

$$\xi_i = ((\pi_i : B_i \rightarrow T, \sigma_i : T \rightarrow B_i), h_i : B_i \rightarrow X), i = 1, 2,$$

be two families of 1-pointed stable maps such that $h_1 \circ \sigma_1 = h_2 \circ \sigma_2$. Denote by

$$\xi = (\pi : B \rightarrow T, h : B \rightarrow X)$$

the family obtained by taking B to be the union of B_1 and B_2 where the section σ_1 is identified with the section σ_2 .

Lemma 4.9. *If for $i = 1, 2$ the family of unpointed stable maps $(\pi_i : B_i \rightarrow T, h_i : B_i \rightarrow X)$ is twistable, and if for every geometric point $t \in T$, the variety parametrizing free lines that contain $h(\sigma(t))$ is irreducible, then ξ is twistable. If also one of ξ_1, ξ_2 is very twistable, then ξ is very twistable.*

Proof. This follows from Lemma 4.5. First of all, using Proposition 4.8, it suffices to prove the result when $T = \text{Spec } k$ for some algebraically closed field k .

For each of $i = 1, 2$, let \mathcal{M}_i denote the 2-fibered product constructed in the proof of Proposition 4.8, i.e., the objects of \mathcal{M}_i are pairs (ζ_i, θ_i) consisting of a twisting family (resp. very twisting family), $\zeta_i : B_i \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$, such that the induced map

$$\tilde{\zeta}_i = ((B_i, \sigma_i), g_i \circ \rho_i : B_i \rightarrow X)$$

is stable, and an equivalence of objects, $\theta_i : \xi_i \rightarrow \tilde{\zeta}_i$. Since each of ξ_i is twistable, each of \mathcal{M}_i is nonempty.

By the proof of Proposition 4.8, each of \mathcal{M}_i is smooth. By the definition of twisting families, for each $i = 1, 2$ the morphism

$$e_i : \mathcal{M}_i \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1), \zeta_i \mapsto \zeta_i(\sigma_i)$$

has image contained in the unobstructed locus of $\text{ev} : \overline{\mathcal{M}}_{0,1}(X, 1) \rightarrow X$. Let $P \subset \overline{\mathcal{M}}_{0,1}(X, 1)$ denote the preimage under ev of the point $p = h_1(\sigma_1) = h_2(\sigma_2)$. The image of e_i is contained in the smooth locus of P . The claim is that $e_i : \mathcal{M}_i \rightarrow P$ is smooth. The obstruction space at a point ζ_i is a quotient of the cohomology group $H^1(B_i, \zeta_i^* T_{\text{ev}}(-\sigma_i))$. By Definition 4.3 (iv), $\zeta_i^* T_{\text{ev}}$ is generated by global sections. Hence, by Lemma 2.3 (ii), this cohomology group is zero, and e_i is smooth.

Since both $e_1 : \mathcal{M}_1 \rightarrow P$ and $e_2 : \mathcal{M}_2 \rightarrow P$ are smooth, both have nonempty, open image contained in the locus of free lines. By hypothesis, the open subset of P parametrizing free lines is irreducible. Therefore the image of e_1 and the image of e_2 intersect. Choose a family $\zeta_1 \in \mathcal{M}_1$ and $\zeta_2 \in \mathcal{M}_2$ such that $e_1(\zeta_1) = e_2(\zeta_2)$. Then ζ_1 and ζ_2 can be glued to obtain a morphism $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ such that $\zeta|_{B_1} = \zeta_1$ and $\zeta|_{B_2} = \zeta_2$. By Lemma 4.5, ζ is twisting. Moreover, if one of ζ_1, ζ_2 is very twisting, then ζ is very twisting. And $\tilde{\zeta} = \xi$. Therefore ξ is twistable, and it is very twistable if one of ξ_1, ξ_2 is very twistable. \square

Hypothesis 4.10. Let $U \subset \overline{\mathcal{M}}_{0,1}(X, 1)$ denote the preimage of $U_{\text{t-able}} \subset \overline{\mathcal{M}}_{0,0}(X, 1)$ under pr . The evaluation morphism $\text{ev} : U \rightarrow X$ has Zariski dense image, i.e., a general point of X is contained in a twistable line.

Let $\pi : B \rightarrow T$ be a family of prestable curves of arithmetic genus 0.

Definition 4.11. A 1-morphism $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ is *positive* (resp. *very positive*) if

- (i) $(\pi : B \rightarrow T, h : B \rightarrow X)$ is a family of stable maps, i.e., a 1-morphism $\xi : T \rightarrow \overline{\mathcal{M}}_{0,0}(X, \epsilon)$ for some $\epsilon \geq 0$,
- (ii) the dimension of the obstruction group of $\overline{\mathcal{M}}_{0,0}(X, \epsilon)$ is 0 at each point of $\xi(T)$,
- (iii) the dimension of the obstruction group of $\overline{\mathcal{M}}_{0,0}(X, e)$ is 0 at each point of $\text{pr}(\zeta(T))$,
- (iv) the pullback $(\text{pr} \circ \zeta)^* T_{\overline{\mathcal{M}}_{0,0}(X, e)}$ is π -relatively deformation ample, and
- (v) the pullback $\sigma^* \mathcal{O}_\Sigma(\sigma)$ is π -relatively generated by global sections (resp. π -relatively ample).

Remark 4.12. This definition is similar to Definition 4.3. It differs in that e need not equal 1 and that $\text{pr} \circ \zeta$ has image in the unobstructed locus, instead of requiring ζ to have image in the unobstructed locus of the morphism ev .

Lemma 4.13. *Let $\pi : B \rightarrow T$ be a family of prestable curves of arithmetic genus 0 and let $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ be a 1-morphism. There is an open subscheme $U_{\text{pos}} \subset T$ (resp. $U_{v\text{-pos}} \subset T$) with the following property: for every morphism of schemes $f : T' \rightarrow T$, the pullback family $f^* \pi : f^* B \rightarrow T'$ and $f^* \zeta : f^* B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ is positive (resp. very positive) iff $f(T') \subset U_{\text{pos}}$ (resp. $f(T') \subset U_{v\text{-pos}}$).*

Proof. The proof is almost identical to the proof of lemma 4.6. \square

Lemma 4.14. *Let B be a prestable curve of arithmetic genus 0, and let $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ be a positive 1-morphism whose image is contained in the locus of very stable maps.*

- (i) *If B is smooth, then $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ is free, i.e., $\zeta^* T_{\overline{\mathcal{M}}_{0,1}(X, e)}$ is generated by global sections. If ζ is very positive, then ζ is very free, i.e., $\zeta^* T_{\overline{\mathcal{M}}_{0,1}(X, e)}$ is ample.*
- (ii) *The dimension of the obstruction group of the 1-morphism $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ is 0. In particular there is a discrete valuation ring, R , a family of prestable curves, $\pi : \mathcal{B} \rightarrow \text{Spec } R$, and a positive 1-morphism $\zeta_R : \mathcal{B} \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ such that the geometric closed fiber is ζ , such that the geometric generic fiber \mathcal{B}_η is smooth, and such that $\zeta_R(\mathcal{B})$ is contained in the locus of very stable maps.*

Proof. (i): By hypothesis, ζ is positive (resp. very positive). By the relative version of Lemma 3.21, the image of ζ is in the smooth locus of $\overline{\mathcal{M}}_{0,1}(X, e)$ and there is a short exact sequence,

$$0 \longrightarrow \sigma^* \mathcal{O}_\Sigma(\sigma) \longrightarrow \zeta^* T_{\overline{\mathcal{M}}_{0,1}(X, e)} \longrightarrow (\text{pr} \circ \zeta)^* T_{\overline{\mathcal{M}}_{0,0}(X, e)} \longrightarrow 0.$$

By Definition 4.11 (iv) and (v), $(\text{pr} \circ \zeta)^* T_{\overline{\mathcal{M}}_{0,0}(X, e)}$ is deformation ample and $\sigma^* \mathcal{O}_\Sigma(\sigma)$ is generated by global sections (resp. deformation ample). Hence $\zeta^* T_{\overline{\mathcal{M}}_{0,1}(X, e)}$ is generated by global sections (resp. deformation ample by Lemma 2.10 (ii)). Since B is smooth ζ is free (resp. very free).

(ii): Let R' be a discrete valuation ring and let $\pi : \mathcal{B}' \rightarrow \text{Spec } R'$ be a smoothing of B , i.e., there is an isomorphism $i : B \rightarrow \mathcal{B}'_0$ and the generic fiber \mathcal{B}'_η is smooth. For technical reasons it is necessary to compactify $\overline{\mathcal{M}}_{0,1}(X, e)$. Let $j : X \hookrightarrow \overline{X}$ be an open immersion of X into a projective scheme, and let $j_* : \overline{\mathcal{M}}_{0,1}(X, e) \hookrightarrow \overline{\mathcal{M}}_{0,1}(\overline{X}, e)$ be the corresponding open immersion of moduli stacks. By [9], $\overline{\mathcal{M}}_{0,1}(\overline{X}, e)$ is a proper Deligne-Mumford stack with projective coarse moduli space. Form the R' -stack, $\mathcal{B}' \times \overline{\mathcal{M}}_{0,1}(\overline{X}, e)$. With respect to an ample invertible sheaf on the coarse moduli

space, let ϵ denote the degree of the map $(i, \zeta) : B \rightarrow \mathcal{B}' \times \overline{\mathcal{M}}_{0,1}(\overline{X}, e)$. Then (i, ζ) is an object of the *Abramovich-Vistoli stack*,

$$\mathcal{K} = \mathcal{K}_{0,0}(\mathcal{B}' \times \overline{\mathcal{M}}_{0,1}(\overline{X}, e), \epsilon) \rightarrow \mathrm{Spec} R',$$

that parametrizes *twisted stable maps* to $\mathcal{B}' \times \overline{\mathcal{M}}_{0,1}(\overline{X}, e)$ over $\mathrm{Spec} R'$, cf. [1]. The stack $\mathcal{K} \rightarrow \mathrm{Spec} R'$ is a Deligne-Mumford stack that is proper over $\mathrm{Spec} R'$, and the coarse moduli space is projective over $\mathrm{Spec} R'$.

The deformation theory of twisted stable maps is developed in [1, Section 5], and is exactly analogous to the deformation theory of stable maps. In particular, the obstruction space to the morphism $o : \mathcal{K} \rightarrow \mathrm{Spec} R'$ at $[(i, \zeta)]$ is a quotient of $H^1(B, \zeta^* T_{\overline{\mathcal{M}}_{0,1}(X,e)})$. Since $\zeta^* T_{\overline{\mathcal{M}}_{0,1}(X,e)}$ is generated by global sections, by Lemma 2.3 (i), $h^1(B, \zeta^* T_{\overline{\mathcal{M}}_{0,1}(X,e)}) = 0$. Therefore o is smooth at $[(i, \zeta)]$. By [6, Prop. 2.2.14], there exists an étale morphism of discrete valuation ring's, $\mathrm{Spec} R \rightarrow \mathrm{Spec} R'$ and a section $s : \mathrm{Spec} R \rightarrow \mathcal{K}$ of o . In other words, there exists a family of twisted stable maps over $\mathrm{Spec} R$ whose closed fiber is isomorphic to the pullback of (i, ζ) , i.e.,

$$(\pi : \mathcal{B} \rightarrow \mathrm{Spec} R', \lambda : \mathcal{B} \rightarrow \mathrm{Spec} R \times_{\mathrm{Spec} R'} (\mathcal{B}' \times \overline{\mathcal{M}}_{0,1}(\overline{X}, e))).$$

Because B is a scheme, the family of curves over $\mathrm{Spec} R$ is a scheme, i.e., $\pi : \mathcal{B} \rightarrow \mathrm{Spec} R$ is a morphism of schemes. Because the composition of (i, ζ) with projection onto the closed fiber of \mathcal{B}' is an isomorphism, the composition

$$\mathcal{B} \xrightarrow{\lambda} \mathrm{Spec} R \times_{\mathrm{Spec} R'} (\mathcal{B}' \times \overline{\mathcal{M}}_{0,1}(\overline{X}, e)) \xrightarrow{\mathrm{pr}_{\mathcal{B}'}} \mathrm{Spec} R \times_{\mathrm{Spec} R'} \mathcal{B}',$$

is an isomorphism. Since the image of (i, ζ) is contained in the open substack $\mathcal{B}' \times \overline{\mathcal{M}}_{0,1}(X, e)$, the image of λ is contained in the the open substack $\mathrm{Spec} R \times_{\mathrm{Spec} R'} (\mathcal{B}' \times \overline{\mathcal{M}}_{0,1}(X, e))$. Putting the pieces together, $\pi : \mathcal{B} \rightarrow \mathrm{Spec} R$ is a smoothing of B , and λ induces a 1-morphism of R -stacks, $\zeta_R : \mathcal{B} \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ such that the closed fiber is ζ .

By Lemma 4.13, ζ_R is positive. Since the locus of very stable maps in $\overline{\mathcal{M}}_{0,1}(X, e)$ is open, the image of ζ_R is contained in this locus. \square

Remark 4.15. The use of the Abramovich-Vistoli stack above is a bit contrived. The most natural stack to use is the Hom stack parametrizing 1-morphisms between two flat, proper, tame Deligne-Mumford stacks. Martin Olsson has proved existence and some foundational properties of the Hom stack in the generality needed above, but the details have not yet been published.

Following is the main definition of this section.

Definition 4.16. Let $\pi : B \rightarrow T$ be a family of prestable curves of arithmetic genus 0. An *inducting pair of degree e* is a pair of 1-morphisms

$$(\zeta_1 : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1), \overline{\zeta}_e : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)),$$

such that

- (i) ζ_1 is very twisting,
- (ii) $\overline{\zeta}_e$ is very positive and the image of $\overline{\zeta}_e$ is contained in the locus of very stable maps, and
- (iii) the morphisms $h_{\zeta_1} : B \rightarrow X$ and $h_{\overline{\zeta}_e} : B \rightarrow X$ are equal.

Lemma 4.17. *Let $\pi : B \rightarrow T$ be a family of prestable curves of arithmetic genus 0, and let*

$$(\zeta_1 : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1), \overline{\zeta}_e : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)),$$

be a pair of 1-morphisms such that $h_{\zeta_1} = h_{\bar{\zeta}_e}$. Then there is an open subscheme $U_{\text{induct}} \subset T$ with the following property: for every morphism of schemes $f : T' \rightarrow T$, the pullback of $(\zeta_1, \bar{\zeta}_e)$ is inducting iff $f(T') \subset U$.

Proof. Define U_{induct} to be the intersection of the open subset $U_{\text{vtwist}} \subset T$ from Lemma 4.6 for ζ_1 and the open subset $U_{\text{v-pos}} \subset T$ from Lemma 4.13 for $\bar{\zeta}_e$. \square

The final definition of this section is the following.

Definition 4.18. Let $(\pi : B \rightarrow T, \bar{\zeta}_e : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e))$ be a very positive family whose image is contained in the locus of very stable maps. The 1-morphism $\bar{\zeta}_e$ is *inductable* if there exists a surjective étale morphism $u : T' \rightarrow T$ and a morphism $\zeta_1 : u^*B \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ such that

- (i) $h_{\zeta_1} = u^*h_{\bar{\zeta}_e}$, and
- (ii) $(\zeta_1, \bar{\zeta}_e)$ is an inducting pair.

Lemma 4.19. Let $(\pi : B \rightarrow T, \bar{\zeta}_e : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e))$ be a very free family. There is an open subscheme $U_{i\text{-able}} \subset T$ with the following property: for every morphism of schemes $f : T' \rightarrow T$, the pullback $(f^*\pi : f^*B \rightarrow T', f^*\bar{\zeta}_e : f^*B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e))$ is inductable iff $f(T') \subset U_{i\text{-able}}$.

Proof. Apply Proposition 4.8 to

$$\xi := (\pi : B \rightarrow T, h_{\bar{\zeta}_e} : B \rightarrow X).$$

\square

5. THE INDUCTION ARGUMENT

In this section it is proved that if X satisfies Hypotheses 1.5, 1.6, 1.7, and 4.10, and if there exists an inductable 1-morphism $\bar{\zeta}_e : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$, then there exists an inductable 1-morphism $\bar{\zeta}_{e+1} : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e+1)$. The basic idea is, given an inducting pair $(\zeta_1, \bar{\zeta}_{e+1})$, to form the family of “connected sums”, i.e., the family of reducible curves obtained by gluing the families ζ_1 and $\bar{\zeta}_{e+1}$ along the two sections. However, this is a family of unpointed curves, and an inducting family is a family of pointed curves. By hypothesis, the section of $\bar{\zeta}_e$, $\bar{\sigma} : B \rightarrow \overline{\Sigma}$, is such that the Cartier divisor $\bar{\sigma}(B) \subset \overline{\Sigma}$ moves. A general member of the linear system $|\bar{\sigma}(B)|$ is the image of a section, $\sigma' : B \rightarrow \overline{\Sigma}$. Except at finitely many points of B , the sections $\bar{\sigma}$ and σ' are disjoint. Away from these points, the section of the family of reducible curves is taken to be σ' . The stable limit over the finitely many points of B is obtained by blowing up $\overline{\Sigma}$ along the zero-dimensional scheme $\bar{\sigma}(B) \cap \sigma'(B)$. Unfortunately, the resulting family of pointed, reducible curves is no longer very positive.

To make the resulting family very positive, the family is altered by a *modification*.

5.1. Modification.

Definition 5.1. An *input triple* is a triple $I = (\zeta, L, \sigma_i)$ where

- (i) $\zeta = ((\Sigma, \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X)$ is an r -pointed stable map to X of arithmetic genus g ,
- (ii) $L \subset \Sigma$ is an irreducible component that is smooth, and
- (iii) $\sigma_i \in L$ is a marked point.

Let $I = (\zeta, L, \sigma_i)$ be an input triple. Denote by $M \subset \Sigma$ the union of all irreducible components other than L . Denote by $R = (\rho_1, \dots, \rho_c)$ the intersection $L \cap M$, denote by $S = (\sigma_{j_1}, \dots, \sigma_{j_d})$ the marked points that are contained in L other than σ_i , and denote by $S' = (\sigma_{k_1}, \dots, \sigma_{k_e})$ the marked points that are contained in M .

Denote by $\Delta : L \rightarrow L \times L$ the diagonal, and denote by $u : \Lambda \rightarrow L \times L$ the blowing up of $L \times L$ along $\Delta(R \cup S)$. For each closed point $\lambda \in R \cup S$, denote by $F_\lambda \subset \Lambda$ the proper transform of $L \times \{\lambda\} \subset L \times L$. Denote by $F_\Delta \subset \Lambda$ the proper transform of $\Delta(L) \subset L \times L$. For each $\lambda \in R \cup S$, there exists a unique section of $\text{pr}_1 \circ u : \Lambda \rightarrow L$, $\sigma_{I,\lambda} : L \rightarrow \Lambda$, such that $F_\lambda = \sigma_{I,\lambda}(L)$. Also there is a section of $\text{pr}_1 \circ u$, $\sigma_{I,\Delta}$, such that $\sigma_{I,\Delta}(L) = F_\Delta$.

Consider the projection $\text{pr}_1 : L \times M \rightarrow L$. For each closed point $\lambda \in R \cup S'$ there is a unique section of pr_1 , $\sigma'_{I,\lambda}$, such that $\sigma'_{I,\lambda}(L) = L \times \{\lambda\}$. Let Σ_I be the simple normal crossing surface containing Λ and $L \times M$ that is obtained by identifying the divisor $\sigma_{I,\lambda}(L) \subset \Lambda$ with $\sigma'_{I,\lambda}(L) \subset L \times M$ for each $\lambda \in R$. The divisors are identified by $\sigma_{I,\lambda}(t) \leftrightarrow \sigma'_{I,\lambda}(t)$ for each $t \in L$. There is a unique morphism $p_I : \Sigma_I \rightarrow L$ whose restriction to Λ is $\text{pr}_1 \circ u$ and whose restriction to $L \times M$ is pr_1 . There is a unique morphism $g_I : \Sigma_I \rightarrow X$ whose restriction to Λ is $g|_L \circ \text{pr}_2 \circ u$ and whose restriction to $L \times M$ is $g|_M \circ \text{pr}_2$. For each integer $j = 1, \dots, r$ such that $j \neq i$, there exists a section of p_I , $\sigma_{I,j} : L \rightarrow \Sigma_I$ given by

$$\sigma_{I,j} = \begin{cases} \sigma_{I,\lambda}, & \lambda = \sigma_j \in S, \\ \sigma'_{I,\lambda}, & \lambda = \sigma_j \in S' \end{cases}$$

And there exists a section $\sigma_{I,i} : L \rightarrow \Sigma_I$ given by $\sigma_{I,i} = \sigma_{I,\Delta}$. By construction, the sections $\sigma_{I,j}$ are pairwise disjoint. Hence the datum

$$\zeta_I = ((p_I : \Sigma_I \rightarrow L, \sigma_{I,1}, \dots, \sigma_{I,r}), g_I : \Sigma_I \rightarrow X)$$

is a family of r -pointed stable maps to X of arithmetic genus g . And the fiber over $\sigma_i \in L$ is canonically isomorphic to ζ .

Definition 5.2. The family of stable maps, ζ_I , is the *modification associated to I* .

The relevance of this construction is the following. Let B be a prestable curve and let

$$\zeta = ((p : \Sigma \rightarrow B, \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X)$$

be a family of r -pointed stable maps to X of arithmetic genus g . Let $b \in B$ be a smooth point, let $L \subset \Sigma_b$ be an irreducible component that is smooth, and let $\sigma_i(b) \in L$ be a marked point. Then $I = (\zeta_b, L, \sigma_i(b))$ is an input triple. Let ζ_I be the modification associated to I . The fiber $\zeta_{I, \sigma_i(b)}$ is canonically isomorphic to the fiber ζ_b . Let \tilde{B} be the prestable curve containing B and L obtained by identifying $b \in B$ with $\sigma_i(b) \in L$. There is a unique family of r -pointed stable maps to X of arithmetic genus g ,

$$\tilde{\zeta} = ((\tilde{p} : \tilde{\Sigma} \rightarrow \tilde{B}, \tilde{\sigma}_1, \dots, \tilde{\sigma}_r), \tilde{g} : \tilde{\Sigma} \rightarrow X),$$

whose restriction to B is ζ and whose restriction to L is ζ_I .

Definition 5.3. The family of stable maps, $\tilde{\zeta}$, is the *modification of ζ associated to $(b, L, \sigma_i(b))$* .

Let $I = (\zeta, L, \sigma_i)$ be an input triple. Denote by $\zeta_{\hat{\sigma}_i}$ the $(r-1)$ -pointed stable map obtained by forgetting the marked point σ_i .

Lemma 5.4. *Let $I = (\zeta, L, \sigma_i)$ be an input triple.*

- (i) *The dimension of the obstruction space of $\zeta_{\hat{\sigma}_i}$ is 0 and for each closed point $\lambda \in R$, there exists a first-order deformation of $\zeta_{\hat{\sigma}_i}$ that smoothes the node λ iff the same is true for ζ .*
- (ii) *For every closed point $\lambda \in L$, the dimension of the obstruction space of $\zeta_{I,\lambda}$ is 0 iff both the dimension of the obstruction space of $\zeta_{\hat{\sigma}_i}$ is 0 and for each closed point $\lambda \in R$ there exists a first-order deformation of $\zeta_{\hat{\sigma}_i}$ that smoothes the node λ .*

(iii) If the equivalent conditions of (ii) are satisfied then there is an exact sequence,

$$\begin{array}{ccccccc} 0 & \rightarrow & T_L & \rightarrow & \zeta_I^* T_{\overline{\mathcal{M}}_{g,r}(X,e)} & \rightarrow & \dots \\ \dots & \rightarrow & T_{\overline{\mathcal{M}}_{g,r-1}(X,e), \zeta_{\sigma_i}} \otimes_{\mathbb{C}} \mathcal{O}_L & \rightarrow & \bigoplus_{\lambda \in R} T_{L,\lambda} \otimes_{\mathcal{O}_\lambda} T_{M,\lambda} & \rightarrow & 0. \end{array}$$

(iv) If the equivalent conditions of (iii) are satisfied, if L has genus 0, and if the map,

$$T_{\overline{\mathcal{M}}_{g,r-1}(X,e), \zeta_{\sigma_i}} \rightarrow \bigoplus_{\lambda \in R} T_{L,\lambda} \otimes T_{M,\lambda},$$

is surjective, then $h^1(L, \zeta_I^* T_{\overline{\mathcal{M}}_{g,r}(X,e)}) = 0$.

Proof. The item (i) follows by Lemma 3.22. The items (ii) and (iii) follow by Lemmas 3.18, 3.19, 3.21 and 3.22. If L has genus 0, then T_L is generated by global sections. So (iv) follows from the long exact sequences in cohomology associated to the exact sequence in (ii) and by Lemma 2.3 (i). \square

Let ζ be a stable map of arithmetic genus 0 and let $I = (\zeta, L, \sigma_i)$ be an input pair.

Lemma 5.5. *If the dimension of the obstruction group of $g|_L : L \rightarrow X$ is 0, and if the pullback of T_X by $g|_M : M \rightarrow X$ is generated by global sections, then for every closed point $\lambda \in L$, the dimension of the obstruction space of $\zeta_{I,\lambda}$ is 0, and $h^1(L, \zeta_I^* T_{\overline{\mathcal{M}}_{0,r}(X,e)}) = 0$.*

Proof. By Lemma 3.17, Lemma 3.21 and Lemma 3.22, the family of pointed stable curves,

$$((\text{pr}_1 \circ u : \Lambda \rightarrow L, \sigma_\lambda, \sigma_\Delta), g|_L \circ \text{pr}_2 \circ u : \Lambda \rightarrow X),$$

is such that for each point of L the dimension of the obstruction space is 0. Consider the families obtained by successively gluing on the connected components of $L \times M$. Applying Lemma 3.29 to each of these families, the hypotheses of Lemma 5.4 (iv) are satisfied for ζ_I . \square

Let B be a prestable curve of arithmetic genus 0, and let

$$\zeta = ((p : \Sigma \rightarrow B, \sigma_1, \dots, \sigma_r), g : \Sigma \rightarrow X),$$

be a family of r -pointed stable maps of arithmetic genus 0. Let $b \in B$ be a smooth point, let $L \subset \Sigma_b$ be an irreducible component that is not contracted by g , and let $\sigma_i(b) \in L$ be a marked point. Let M be the union of all irreducible components of Σ_b other than L .

Lemma 5.6. *If for every point $b' \in B$ the dimension of the obstruction group of $\zeta_{b'}$ is 0, if $\zeta^* T_{\overline{\mathcal{M}}_{0,r}(X,e)}$ is generated by global sections, if the dimension of the obstruction group of $g|_L : L \rightarrow X$ is 0, and if the pullback of T_X by $g|_M : M \rightarrow X$ is generated by global sections, then for every point $b' \in \tilde{B}$ the dimension of the obstruction group of $\tilde{\zeta}_{b'}$ is 0 and $h^1(\tilde{B}, \tilde{\zeta}^* T_{\overline{\mathcal{M}}_{0,r}(X,e)}) = 0$.*

Therefore the dimension of the obstruction group of $\tilde{\zeta} : \tilde{B} \rightarrow \overline{\mathcal{M}}_{0,r}(X,e)$ is 0. In particular there is a discrete valuation ring, R , a family of prestable curves, $\pi : \mathcal{B} \rightarrow \text{Spec } R$, and a 1-morphism $\zeta_R : \mathcal{B} \rightarrow \overline{\mathcal{M}}_{0,r}(X,e)$ such that the geometric closed fiber is $\tilde{\zeta}$ and such that the geometric generic fiber \mathcal{B}_η is smooth.

Proof. The hypotheses of Lemma 5.5 are satisfied for the modification associated to $I = (\zeta_b, L, \sigma_i(b))$, $\tilde{\zeta}_L : L \rightarrow \overline{\mathcal{M}}_{0,r}(X,e)$. Hence the dimension of the obstruction group of $\tilde{\zeta}$ at each closed point of L is 0 and $h^1(L, \tilde{\zeta}_L^* T_{\overline{\mathcal{M}}_{0,r}(X,e)}) = 0$. There is a short exact sequence of coherent sheaves on \tilde{B} ,

$$0 \longrightarrow \zeta^* T_{\overline{\mathcal{M}}_{0,r}(X,e)}(-b) \longrightarrow \tilde{\zeta}^* T_{\overline{\mathcal{M}}_{0,r}(X,e)} \longrightarrow \tilde{\zeta}_L^* T_{\overline{\mathcal{M}}_{0,r}(X,e)} \longrightarrow 0.$$

By Lemma 2.3 (ii), $h^1(B, \zeta^* T_{\overline{\mathcal{M}}_{0,r}(X,e)}(-b)) = 0$. Hence by the long exact sequence of cohomology associated to the short exact sequence above, $h^1(\tilde{B}, \tilde{\zeta}^* T_{\overline{\mathcal{M}}_{0,r}(X,e)}) = 0$.

The second part of the lemma follows by the same proof as in Lemma 4.14 (ii). \square

5.2. The induction argument.

Notation 5.7. The notation for the generators of the group of \mathbb{Q} -Cartier divisor classes on $\overline{\mathcal{M}}_{0,1}(\mathbb{P}^n, e)$ is taken from [19].

- (i) For each integer $0 \leq i \leq [e]$ denote by Δ_i , or sometimes $\Delta_{i,e-i}$, the class of the \mathbb{Q} -Cartier divisor that is the closure of the locus parametrizing embedded curves that have one irreducible component of degree i , one irreducible component of degree $e - i$ and where the marked point is on the irreducible component of degree i .
- (ii) Denote by \mathcal{L} the first Chern class of $\text{ev}^* \mathcal{O}_{\mathbb{P}^n}(1)$.
- (iii) Denote by \mathcal{H} the class of the \mathbb{Q} -Cartier divisor parametrizing stable maps $g : \Sigma \rightarrow \mathbb{P}^n$ such that $g(\Sigma) \cap \Lambda \neq \emptyset$, where $\Lambda \subset \mathbb{P}^n$ is a fixed linear \mathbb{P}^{n-2} .

Let B be a prestable curve of genus 0. Let $\zeta = ((p : \Sigma \rightarrow B, \sigma), g : \Sigma \rightarrow X)$ be a family of stable maps of genus 0 and degree $e > 1$. Denote by $h : B \rightarrow X$ the composition $g \circ \sigma$.

Definition 5.8. The 1-morphism $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ is *typical* if,

- (i) the curve B is smooth,
- (ii) for every closed point $b \in B$, the obstruction group of ζ_b has dimension 0,
- (iii) for every point $b \in B$, the stable map ζ_b has only the trivial automorphism,
- (iv) for every integer $j = 1, \dots, e - 1$, the preimage $\zeta^{-1}(\Delta_j)$ consists of finitely many points, and for each $b \in \zeta^{-1}(\Delta_j)$, the curve Σ_b has only two irreducible components,
- (v) for every point $b \in \zeta^{-1}(\Delta_1)$, Σ_b is a union of 2 irreducible components $L \cup M$ with $\sigma(b) \in L$ such that $g|_L : L \rightarrow X$ is an isomorphism to a twistable line.

Lemma 5.9. *Let $\pi : B \rightarrow T$ be a family of prestable curves of arithmetic genus 0 and let $\zeta : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ be a 1-morphism. There is an open subscheme $U_{\text{typ}} \subset T$ with the following property: for every morphism of schemes $f : T' \rightarrow T$, the pullback family $f^* \pi : f^* B \rightarrow T'$ and $f^* \zeta : f^* B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ is typical iff $f(T') \subset U_{\text{typ}}$.*

Proof. Each of the conditions in Definition 5.8 is clearly an open condition. \square

Lemma 5.10. *If X satisfies Hypotheses 1.5, 1.6, 1.7, and 4.10, if for every closed point $b \in B$ the obstruction group of ζ_b has dimension 0, and if $\zeta^* T_{\overline{\mathcal{M}}_{0,1}(X, e)}$ is generated by global sections, then there exists a discrete valuation ring, R , a family of prestable curves, $\pi : \mathcal{B} \rightarrow \text{Spec } R$, and a 1-morphism $\zeta_R : \mathcal{B} \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ such that the geometric closed fiber is ζ , such that the geometric generic fiber \mathcal{B}_η is smooth, and such that $(\zeta_R)_\eta$ is typical.*

Proof. By the same argument as in the proof of Lemma 4.13, there exists a family $\zeta_R : \mathcal{B} \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ such that \mathcal{B}_η is smooth. The condition that for every point the obstruction group has dimension 0 is stable under generization, so it also holds for $(\zeta_R)_\eta$. Similarly, $(\zeta_R)_\eta^* T_{\overline{\mathcal{M}}_{0,1}(X, e)}$ is generated by global sections. By [16, Prop. 3.7], the family ζ_R may be chosen so that $(\zeta_R)(\mathcal{B}_\eta)$ is disjoint from any given finite collection of closed substacks of codimension at least 2. By Hypothesis 1.5, the locus of stable maps such that the domain curve has 3 or more irreducible components has codimension at least 2. Let $Z \subset \Delta_i$ be a closed substack whose image in the coarse moduli space of Δ_i is an ample divisor. Again by Hypothesis 1.5, Z has codimension 2 in $\overline{\mathcal{M}}_{0,1}(X, e)$. So $(\zeta_R)(\mathcal{B}_\eta)$ does not intersect Z , and therefore it is not contained in Δ_i . By Hypothesis 1.7, the locus of stable maps that have a nontrivial automorphism has codimension at least 2. Finally, by Hypothesis 1.6 and Hypothesis 4.10, the locus in Δ_1 parametrizing stable maps $g : L \cup M \rightarrow X$ such that $g|_L : L \rightarrow X$ is an isomorphism to a twistable line is a dense open in Δ_1 . In particular the complement in Δ_1 is a closed substack that has codimension 2 in $\overline{\mathcal{M}}_{0,1}(X, e)$. Therefore $(\zeta_R)_\eta$ is typical. \square

By Hypothesis 4.10, a general line $L \subset X$ is twistable. Denote by,

$$\zeta_L = ((p_L : \Sigma_L \rightarrow L, \sigma_L), g_L : \Sigma_L \rightarrow X),$$

a twisting family such that $h_L : L \rightarrow X$ is the inclusion. By Remark 4.4 (iv), the degree of g_L is either 1 or 2, i.e., either $g_L : \Sigma_L \rightarrow X$ is a birational map to a linear \mathbb{P}^2 in X (obtained by blowing up a point on the \mathbb{P}^2), or g_L is an isomorphism of Σ_L to a nonsingular quadric surface in X .

Definition 5.11. Let X be a variety that satisfies Hypotheses 1.6 and 4.10. The variety X is *planar type* if for a general line L there exists a twisting family such that the degree of g_L is 1. The variety is *quadric type* if it is not planar type.

Remark 5.12. It is not hard to prove that if X is planar type, then there exists an integer $r \geq 2$ such that for a general point $x \in X$, the set of lines in X that contain x sweep out a linear \mathbb{P}^r . Let Q denote the irreducible component of the Hilbert scheme of \mathbb{P}^r 's in X that contains the \mathbb{P}^r 's constructed in this way. Then X is birationally a \mathbb{P}^r -bundle over Q and, with its natural Plücker embedding, Q is not uniruled by lines. If X satisfies Hypothesis 1.5, then Q is not uniruled by rational curves of any degree. If also X contains a very twistable curve, then Q is a point and X is isomorphic to \mathbb{P}^r (not merely birational to \mathbb{P}^r). Thus, in what follows, X will usually be of quadric type.

Let $(\zeta_1, \bar{\zeta}_e)$ be an inducting pair denoted by,

$$\begin{aligned} \zeta_1 &= ((p : \Sigma \rightarrow B, \sigma), g : \Sigma \rightarrow X), \\ \bar{\zeta}_e &= ((\bar{p} : \bar{\Sigma} \rightarrow B, \bar{\sigma}), \bar{g} : \bar{\Sigma} \rightarrow X). \end{aligned}$$

Denote

$$\begin{aligned} \delta &= \deg(\zeta_1^*(2\mathcal{L} - \mathcal{H})), \\ \bar{\delta} &= \deg(\bar{\zeta}_e^*(\frac{2}{e}\mathcal{L} - \frac{1}{e^2}\mathcal{H} - \sum_{i=1}^{e-1} \frac{(e-i)^2}{e^2}\Delta_i)). \end{aligned}$$

By [19, Lem. 2.2.2], $\deg(\sigma^*\mathcal{O}_\Sigma(\sigma(B))) = \delta$ and $\deg(\bar{\sigma}^*\mathcal{O}_{\bar{\Sigma}}(\bar{\sigma}(B))) = \bar{\delta}$. In particular, both δ and $\bar{\delta}$ are nonnegative integers, and $\bar{\delta}$ is positive by Definition 4.11.

Theorem 5.13. *For each integer $d = 1, \dots, \bar{\delta}$, there exists an inducting pair $(\xi_{d,1}, \bar{\xi}_{d,e+1})$ such that $\bar{\xi}_{d,e+1}$ is typical and such that the following conditions are satisfied,*

$$\left\{ \begin{array}{ll} \deg(\bar{\xi}_{d,e+1}^*\mathcal{H}) &= \deg(\bar{\zeta}_e^*\mathcal{H}) + \deg(\zeta_1^*\mathcal{H}), \\ \deg(\xi_{d,1}^*\mathcal{H}) &= \deg(\zeta_1^*\mathcal{H}) + d, & \text{if } X \text{ is planar type,} \\ \deg(\xi_{d,1}^*\mathcal{H}) &= \deg(\zeta_1^*\mathcal{H}) + 2d, & \text{if } X \text{ is quadric type,} \\ \deg(\xi_{d,1}^*\mathcal{L}) &= \deg(\zeta_1^*\mathcal{L}) + d, \\ \deg(\bar{\xi}_{d,e+1}^*\mathcal{L}) &= \deg(\bar{\zeta}_e^*\mathcal{L}) + d, \\ \deg(\bar{\xi}_{d,e+1}^*\Delta_i) &= \deg(\bar{\zeta}_e^*\Delta_{i-1}), & i = 2, \dots, e-1, \\ \deg(\bar{\xi}_{d,e+1}^*\Delta_e) &= \deg(\bar{\zeta}_e^*\Delta_{e-1}) + \delta + d, & \text{if } i > 1, \\ \deg(\bar{\xi}_{d,e+1}^*\Delta_1) &= \bar{\delta} - d, & \text{if } i > 1, \\ \deg(\bar{\xi}_{d,2}^*\Delta_1) &= \delta + \bar{\delta}, & \text{if } i = 1. \end{array} \right. \quad (9)$$

The families are denoted,

$$\begin{aligned} \bar{\xi}_{d,e+1} &= ((\bar{p}_{d,e+1} : \bar{\Sigma}_{d,e+1} \rightarrow B, \bar{\sigma}_{d,e+1}), \bar{g}_{d,e+1} : \bar{\Sigma}_{d,e+1} \rightarrow X), \\ \xi_{d,1} &= ((p_{d,1} : \Sigma_{d,1} \rightarrow B, \sigma_{d,1}), g_{d,1} : \Sigma_{d,1} \rightarrow X). \end{aligned}$$

Also $\bar{h}_{d,e+1}$ denotes $\bar{g}_{d,e+1} \circ \bar{\sigma}_{d,e+1}$ and $h_{d,1}$ denotes $g_{d,1} \circ \sigma_{d,1}$.

Proof. By Lemma 4.14 (ii) and Lemma 4.19, it suffices to consider the case that B is smooth. By Lemma 4.14(i) $\bar{\zeta}_e : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$ is very free. Hence, the same argument as in the proof of Lemma 4.14 (ii) and [16, Prop. 3.7], it suffices to consider the case that $\bar{\zeta}_e(B_e)$ is in *general position*, i.e., for any finite collection of codimension 2 subvarieties $(Z_\alpha | \alpha = 1, \dots, M)$ and any finite collection of divisors $(D_\beta | \beta = 1, \dots, N)$, $\bar{\zeta}_e(B_e)$ is disjoint from each Z_α and has 0-dimensional intersection with each D_β .

The family $\zeta^{(1)}$: Denote by $\Sigma^{(1)}$ the surface containing Σ and $\bar{\Sigma}$ obtained by identifying the divisor $\sigma(B) \subset \Sigma$ and $\bar{\sigma}(B) \subset \bar{\Sigma}$ via $\sigma(b) \leftrightarrow \bar{\sigma}(b)$. Denote by $p^{(1)} : \Sigma^{(1)} \rightarrow B$ the unique morphism whose restriction to Σ is p and whose restriction to $\bar{\Sigma}$ is \bar{p} . Denote by $g^{(1)} : \Sigma^{(1)} \rightarrow X$ the unique morphism whose restriction to Σ is g and whose restriction to $\bar{\Sigma}$ is \bar{g} . Then $\zeta^{(1)} = (p^{(1)} : \Sigma^{(1)} \rightarrow B, g^{(1)} : \Sigma^{(1)} \rightarrow X)$ is a family of stable maps in the boundary divisor $\Delta_1 \subset \overline{\mathcal{M}}_{0,0}(X, e+1)$. Denote by τ the stable A -graph with two vertices v and \bar{v} of degree 1 and e respectively, with no tails, and with one edge connecting v and \bar{v} . Then $\zeta^{(1)}$ factors through the canonical 1-morphism of Behrend-Manin stacks $\overline{\mathcal{M}}(X, \tau) \rightarrow \overline{\mathcal{M}}_{0,0}(X, e+1)$.

Claim 5.14. *The pullback $(\zeta^{(1)})^*T_{\overline{\mathcal{M}}_{0,0}(X,e)}$ is ample.*

By Lemma 3.28, there is a short exact sequence,

$$0 \longrightarrow \zeta_1^*T_{\text{ev}} \longrightarrow (\zeta^{(1)})^*T_{\overline{\mathcal{M}}(X,\tau)} \longrightarrow \bar{\zeta}_e^*T_{\overline{\mathcal{M}}_{0,1}(X,e)} \longrightarrow 0. \quad (10)$$

Since ζ_1 is very twisting, by Definition 4.3 $\zeta_1^*T_{\text{ev}}$ is ample. Since $\bar{\zeta}_e$ is very positive, by Lemma 4.14, $\bar{\zeta}_e^*T_{\overline{\mathcal{M}}_{0,1}(X,e)}$ is ample. Hence $(\zeta^{(1)})^*T_{\overline{\mathcal{M}}(X,\tau)}$ is ample. By Lemma 3.29, there is a short exact sequence,

$$0 \rightarrow (\zeta^{(1)})^*T_{\overline{\mathcal{M}}(X,\tau)} \rightarrow (\zeta^{(1)})^*T_{\overline{\mathcal{M}}_{0,0}(X,e+1)} \rightarrow \sigma^*\mathcal{O}_\Sigma(\sigma) \otimes \bar{\sigma}^*\mathcal{O}_{\bar{\Sigma}}(\bar{\sigma}) \rightarrow 0. \quad (11)$$

By Definition 4.3 and Definition 4.11, both $\sigma^*\mathcal{O}_\Sigma(\sigma)$ and $\bar{\sigma}^*\mathcal{O}_{\bar{\Sigma}}(\bar{\sigma})$ are ample. Therefore the tensor product is ample, and $(\zeta^{(1)})^*T_{\overline{\mathcal{M}}_{0,0}(X,e+1)}$ is ample.

The family $\zeta^{(2)}$: The family $\zeta^{(1)}$ cannot be an inductable family, because it is a family of unpointed curves rather than 1-pointed curves. The next approximation to $\xi_{d,e+1}$ “adds” a marked section to $\zeta^{(1)}$. The self-intersection of $\bar{\sigma}(B) \subset \bar{\Sigma}$ is the degree of the invertible sheaf $\bar{\sigma}^*\mathcal{O}_{\bar{\Sigma}}(\bar{\sigma}(B))$, which is also

$$\bar{\delta} = \deg \left(2\bar{\zeta}_e^*\mathcal{L} - \bar{\zeta}_e^*\mathcal{H} \right).$$

Let $\varsigma : B \rightarrow \bar{\Sigma}$ be a section such that $\varsigma(B) \subset \bar{\Sigma}$ is a general member of the linear system $|\bar{\sigma}(B)|$. Because $\bar{\sigma}^*\mathcal{O}_{\bar{\Sigma}}(\bar{\sigma}(B))$ is generated by global sections, there exists ς such that $\varsigma(B)$ has only transverse intersections with $\bar{\sigma}(B)$. Denote the points of intersection by $q_1, \dots, q_{\bar{\delta}} \in \bar{\Sigma}$. Denote by $w : \bar{\Sigma}^{(2)} \rightarrow \bar{\Sigma}$ the blowing up of $\bar{\Sigma}$ at the points $q_1, \dots, q_{\bar{\delta}}$. Let $\bar{p}^{(2)} : \bar{\Sigma}^{(2)} \rightarrow B$ denote the projection $\bar{p} \circ w$. Let $\bar{g}^{(2)}$ denote $\bar{g} \circ w$. Let $\bar{\sigma}^{(2)} : B \rightarrow \bar{\Sigma}^{(2)}$ and $\varsigma^{(2)} : B \rightarrow \bar{\Sigma}^{(2)}$ denote the proper transforms of $\bar{\sigma}$ and ς respectively. Notice that $\bar{\sigma}^{(2)}(B)$ and $\varsigma^{(2)}(B)$ are disjoint by construction. So the data

$$\bar{\zeta}_e^{(2)} = ((\bar{p}^{(2)} : \bar{\Sigma}^{(2)} \rightarrow B, \bar{\sigma}^{(2)}, \varsigma^{(2)}), \bar{g}^{(2)} : \bar{\Sigma}^{(2)} \rightarrow X)$$

is a family of stable 2-pointed maps, i.e., a 1-morphism $\bar{\zeta}_e^{(2)} : B \rightarrow \overline{\mathcal{M}}_{0,2}(X, e)$.

Claim 5.15. *The pullback $(\bar{\zeta}_e^{(2)})^*T_{\overline{\mathcal{M}}_{0,2}(X,e)}$ is generated by global sections.*

By Lemmas 3.16, 3.21 and 3.22, there is a short exact sequence:

$$0 \longrightarrow (\varsigma^{(2)})^*\mathcal{O}_{\bar{\Sigma}^{(2)}}(\varsigma^{(1)}(B)) \longrightarrow (\bar{\zeta}_e^{(2)})^*T_{\overline{\mathcal{M}}_{0,2}(X,e)} \longrightarrow \bar{\zeta}_e^*T_{\overline{\mathcal{M}}_{0,1}(X,e)} \longrightarrow 0.$$

By construction, the self-intersection of $\zeta^{(2)}(B) \subset \overline{\Sigma}^{(2)}$ is 0, i.e., $(\zeta^{(2)})^* \mathcal{O}_{\overline{\Sigma}^{(2)}}(\zeta^{(2)}(B))$ is the trivial invertible sheaf \mathcal{O}_B . By hypothesis $\overline{\zeta}_e^* T_{\overline{\mathcal{M}}_{0,1}(X,e)}$ is generated by global sections. Therefore by Lemma 2.3 (iii), $(\overline{\zeta}_e^{(2)})^* T_{\overline{\mathcal{M}}_{0,2}(X,e)}$ is generated by global sections.

Denote by $\Sigma^{(2)}$ the surface containing $\overline{\Sigma}^{(2)}$ and Σ obtained by identifying the divisors $\overline{\sigma}^{(2)}(B) \subset \overline{\Sigma}^{(2)}$ and $\sigma(B) \subset \Sigma$ via $\overline{\sigma}^{(2)}(b) \leftrightarrow \sigma(b)$. Denote by $p^{(2)} : \Sigma^{(2)} \rightarrow B$ the unique morphism whose restriction to $\overline{\Sigma}^{(2)}$ is $\overline{p}^{(2)}$ and whose restriction to Σ is p . Denote by $g^{(2)} : \Sigma^{(2)} \rightarrow X$ the unique morphism whose restriction to $\overline{\Sigma}^{(2)}$ is $\overline{g}^{(2)}$ and whose restriction to Σ is g . Denote by $\sigma^{(2)} : B \rightarrow \Sigma^{(2)}$ the morphism obtained by composing $\zeta^{(2)}$ with the inclusion $\overline{\Sigma}^{(2)} \subset \Sigma^{(2)}$. Then

$$\zeta^{(2)} = ((p^{(2)} : \Sigma^{(2)} \rightarrow B, \sigma^{(2)}), g^{(2)} : \Sigma^{(2)} \rightarrow X)$$

is a family of stable maps in the boundary $\Delta_e \subset \overline{\mathcal{M}}_{0,1}(X, e+1)$. Denote by $\tau^{(2)}$ the stable A -graph with two vertices v and \overline{v} of degree 1 and e respectively, a single tail attached to \overline{v} , and a single edge connecting v and \overline{v} . Then $\zeta^{(2)}$ factors through the canonical 1-morphism of Behrend-Manin stacks $\overline{\mathcal{M}}(X, \tau^{(2)}) \rightarrow \overline{\mathcal{M}}_{0,1}(X, e+1)$.

Claim 5.16. *The pullback $(\zeta^{(2)})^* T_{\overline{\mathcal{M}}_{0,1}(X,e)}$ is generated by global sections.*

By Lemma 3.28, there is a short exact sequence,

$$0 \longrightarrow \zeta_1^* T_{\text{ev}} \longrightarrow (\zeta^{(2)})^* T_{\overline{\mathcal{M}}(X, \tau^{(2)})} \longrightarrow (\overline{\zeta}_e^{(2)})^* T_{\overline{\mathcal{M}}_{0,2}(X,e)} \longrightarrow 0.$$

By Claim 5.15, the third term is generated by global sections. Since ζ_1 is very twisting, by Definition 4.3, $\zeta_1^* T_{\text{ev}}$ is ample. In particular it is generated by global sections. By Lemma 2.3 (iii), $(\zeta^{(2)})^* T_{\overline{\mathcal{M}}(X, \tau^{(2)})}$ is generated by global sections. By Lemma 3.29 there is a short exact sequence,

$$0 \rightarrow (\zeta^{(2)})^* T_{\overline{\mathcal{M}}(X, \tau^{(2)})} \rightarrow (\zeta^{(2)})^* T_{\overline{\mathcal{M}}_{0,1}(X, e+1)} \rightarrow \sigma^* \mathcal{O}_{\Sigma}(\sigma(B)) \otimes_{\mathcal{O}_B} (\overline{\sigma}^{(2)})^* \mathcal{O}_{\overline{\Sigma}^{(2)}}(\overline{\sigma}^{(2)}(B)) \rightarrow 0.$$

By construction, $\mathcal{O}_{\overline{\Sigma}^{(2)}}(\overline{\sigma}^{(2)}(B))$ is isomorphic to \mathcal{O}_B . Since ζ_1 is very twisting, by Definition 4.3, $\sigma^* \mathcal{O}_{\Sigma}(\sigma(B))$ is generated by global sections. Hence the third term in the short exact sequence is generated by global sections. Since also the first term is generated by global sections, by Lemma 2.3 (iii), $(\zeta^{(2)})^* T_{\overline{\mathcal{M}}_{0,1}(X, e+1)}$ is generated by global sections.

If $e > 1$, the image of $\zeta^{(2)}$ intersects the divisor Δ_1 transversely at the images of the points $q_1, \dots, q_{\overline{\delta}}$. In particular, the degree of the \mathbb{Q} -Cartier divisor class $(\zeta^{(2)})^* \mathcal{O}_{\overline{\mathcal{M}}_{0,1}(X, e+1)}(\Delta_1)$ is $\overline{\delta}$, which is positive. If $e = 1$, then $\Delta_1 = \Delta_e$. In this case $\overline{\mathcal{M}}(X, \tau^{(2)})$ is the normalization of $\Delta_{1,1}$ in a neighborhood of $\zeta^{(2)}(B)$. So the degree of $(\zeta^{(2)})^* \mathcal{O}_{\overline{\mathcal{M}}_{0,1}(X, 2)}(\Delta_1)$ is the sum of the degree of the pullback of the normal sheaf of $\overline{\mathcal{M}}(X, \tau^{(2)}) \rightarrow \overline{\mathcal{M}}_{0,1}(X, 2)$ and the degree of the divisor $q_1 + \dots + q_{\overline{\delta}}$, i.e. $\delta + \overline{\delta}$. So also in this case the degree of $(\zeta^{(2)})^* \mathcal{O}_{\overline{\mathcal{M}}_{0,1}(X, e+1)}(\Delta_1)$ is positive.

Similar computations give that $\deg((\zeta^{(2)})^* \Delta_i) = \overline{\zeta}_e^* \Delta_{i-1}$ for $i = 2, \dots, e-1$. The curve $\zeta^{(2)}(B)$ is contained in the divisor Δ_e , which is the image of $\overline{\mathcal{M}}(X, \tau^{(2)})$. As noted above, by Lemma 3.29, the morphism $\overline{\mathcal{M}}(X, \tau^{(2)}) \rightarrow \overline{\mathcal{M}}_{0,1}(X, e+1)$ is unramified and the pullback by $\zeta^{(2)}$ of the normal sheaf has degree $\deg(\sigma^* \mathcal{O}_{\Sigma}(\sigma(B))) = \delta$. If $e > 1$, then the image of $\zeta^{(2)}(B)$ is contained in the smooth locus of Δ_e so that the degree of $(\zeta^{(2)})^* \mathcal{O}_{\overline{\mathcal{M}}_{0,1}(X, e+1)}(\Delta_e)$ is precisely δ . If $e = 1$, then each point $q_1, \dots, q_{\overline{\delta}}$ maps to a point of Δ_1 where Δ_1 intersects itself transversely. So if $e = 1$, the total degree of $(\zeta^{(2)})^* \mathcal{O}_{\overline{\mathcal{M}}_{0,1}(X, 2)}(\Delta_1)$ is $\delta + \overline{\delta}$ (this is the same result from the last paragraph – it is included for the sake of consistency).

It is evident that,

$$\deg((\zeta^{(2)})^*\mathcal{H}) = \deg(\bar{\zeta}_e^*\mathcal{H}) + \deg(\zeta_1^*\mathcal{H}),$$

and $\deg((\zeta^{(2)})^*\mathcal{L}) = \deg(\bar{\zeta}_e^*\mathcal{L})$. As a ‘‘consistency check’’ for the computations of these degrees, observe that by [19, Lem. 2.2.2],

$$\deg((\sigma^{(2)})^*\mathcal{O}_{\Sigma^{(2)}}(\sigma^{(2)}(B))) = \deg\left((\zeta^{(2)})^*\left(\frac{2}{e+1}\mathcal{L} - \frac{1}{(e+1)^2}\mathcal{H} - \sum_{i=1}^e \frac{(e+1-i)^2}{(e+1)^2}\Delta_i\right)\right).$$

Substituting the values above, using the formulas for δ and $\bar{\delta}$, and using that $\zeta_1^*\mathcal{L} = \bar{\zeta}_{e+1}^*\mathcal{L}$, the degree of the right-hand-side is 0. Of course this is correct; $\sigma^{(2)}(B) \subset \Sigma^{(2)}$ has self-intersection 0 by construction.

The family $\bar{\xi}_{0,e+1}$: Before proceeding to the construction of the families $\bar{\xi}_{d,e+1}$, the family $\zeta^{(2)}$ is deformed to a typical family $\bar{\xi}_{0,e+1}$, cf. Definition 5.8. The families $\bar{\xi}_{d,e+1}$ will be constructed as suitable modifications of $\bar{\xi}_{0,e+1}$.

Since $\zeta^{(2)} : B \rightarrow \bar{\mathcal{M}}_{0,1}(X, e+1)$ is free, i.e., $(\zeta^{(2)})^*T_{\bar{\mathcal{M}}_{0,1}(X, e+1)}$ is generated by global sections, it follows by the same argument as in the proof of Lemma 4.14 (ii) and [16, Prop. 3.7] that there exists a deformation $\bar{x}i_{0,e+1} : B \rightarrow \bar{\mathcal{M}}_{0,1}(X, e+1)$ of $\zeta^{(2)}$ that is in general position. Here *deformation* means that both $\zeta^{(2)}$ and $\bar{\xi}_{0,e+1}$ have a common generalization. Moreover, for any finite collection of open conditions that are satisfied by $\zeta^{(2)}$, the deformation $\bar{\xi}_{0,e+1}$ may be chosen to also satisfy these conditions. Denote the family of stable maps by,

$$\bar{\xi}_{0,e+1} = ((\bar{p}_{0,e+1} : \bar{\Sigma}_{0,e+1} \rightarrow B, \bar{\sigma}_{0,e+1}), \bar{g}_{0,e+1} : \bar{\Sigma}_{0,e+1} \rightarrow X).$$

Denote by $\bar{h}_{0,e+1} : B \rightarrow X$ the composition $\bar{g}_{0,e+1} \circ \bar{\sigma}_{0,e+1}$.

Claim 5.17. *There exists a deformation $\bar{\xi}_{0,e+1}$ of $\zeta^{(2)}$ such that*

- (i) $\bar{\xi}_{0,e+1}$ is typical,
- (ii) $(\text{pr} \circ \bar{\xi}_{0,e+1})^*T_{\bar{\mathcal{M}}_{0,0}(X, e+1)}$ is ample,
- (iii) $\bar{h}_{0,e+1} : B \rightarrow X$ is very twistable,
- (iv) and $(\bar{\sigma}_{0,e+1})^*\mathcal{O}_{\bar{\Sigma}_{0,e+1}}(\bar{\sigma}_{0,e+1}(B))$ is isomorphic to \mathcal{O}_B .

It suffices to prove that for a general deformation $\bar{\xi}_{0,e+1}$ of $\zeta^{(2)}$, $\bar{\xi}_{0,e+1}$ satisfies the properties in Claim 5.17, i.e., for every irreducible component of the Hom stack $\text{Hom}(B, \bar{\mathcal{M}}_{0,1}(X, e))$ that contains $\zeta^{(2)}$, each of the conditions (i)–(iv) holds on a dense open substack.

(i): By Lemmas 5.9 and 5.10, a general deformation of $\zeta^{(2)}$ is typical.

(ii): The pullback by $\text{pr} \circ \zeta^{(2)} : B \rightarrow \bar{\mathcal{M}}_{0,0}(X, e+1)$ of $T_{\bar{\mathcal{M}}_{0,0}(X, e+1)}$ equals $(\zeta^{(1)})^*T_{\bar{\mathcal{M}}_{0,0}(X, e)}$. By Claim 5.14, this is ample. The condition on deformations of $\zeta^{(2)}$ that the pullback of $T_{\bar{\mathcal{M}}_{0,0}(X, e)}$ is ample is an open condition by Lemma 2.8 and Lemma 2.9 (or more direct arguments). Therefore the pullback $(\text{pr} \circ \bar{\xi}_{0,e+1})^*T_{\bar{\mathcal{M}}_{0,0}(X, e)}$ is ample.

(iii): The morphism $g^{(2)} \circ \sigma^{(2)} : B \rightarrow X$ equals $\bar{g} \circ \zeta$, and this is a deformation of $\bar{g} \circ \bar{\sigma} = g \circ \sigma$. Because ζ_1 is very twisting, $g \circ \sigma$ is very twistable. By Proposition 4.8, ζ can be chosen so that $g^{(2)} \circ \sigma^{(2)}$ is very twistable. Since $\bar{h}_{0,e+1}$ is a deformation of $g^{(2)} \circ \sigma^{(2)}$, by Proposition 4.8, $\bar{\xi}_{0,e+1}$ can be chosen so that $\bar{h}_{0,e+1}$ is very twistable.

(iv): Since $(\sigma^{(2)})^* \mathcal{O}_{\Sigma^{(2)}}(\sigma^{(2)}(B))$ is isomorphic to \mathcal{O}_B , also $(\bar{\sigma}_{0,e+1})^* \mathcal{O}_{\bar{\Sigma}_{0,e+1}}(\bar{\sigma}_{0,e+1}(B))$ is isomorphic to \mathcal{O}_B . This proves Claim 5.17. In particular, $\bar{\xi}_{0,e+1}$ is positive and $\bar{\xi}_{0,e+1}(B)$ is contained in the locus of very stable maps. Unfortunately it is *not* very positive!

To be a bit more precise in the proof of (iii), there exists a deformation of $\zeta_1, \xi_{0,1} : B \rightarrow \bar{\mathcal{M}}_{0,1}(X, 1)$ that is very twisting and such that $h_{0,1} := g_{0,1} \circ \sigma_{0,1} : B \rightarrow X$ equals $\bar{h}_{0,e+1}$.

Because $\bar{\xi}_{0,e+1}$ is a deformation of $\zeta^{(2)}$, the intersection number of $\bar{\xi}_{0,e+1}(B)$ with any divisor of $\bar{\mathcal{M}}_{0,1}(X, e+1)$ is equal to the intersection number of $\zeta^{(2)}(B)$ with that divisor. These intersection numbers were computed above. Similarly, the intersection number of $\xi_{0,1}(B)$ with any divisor in $\bar{\mathcal{M}}_{0,1}(X, 1)$ is equal to the intersection number of $\zeta_1(B)$ with that divisor.

The families $\bar{\xi}_{d,e+1}$: The families $\bar{\xi}_{d,e+1}$ and $\xi_{d,1}$ are constructed by induction on d . The base case $d = 0$ is the pair $(\xi_{0,1}, \bar{\xi}_{0,e+1})$ constructed above, and the induction step consists of performing a single modification and then deforming.

Let d be an integer $1 \leq d \leq \bar{\delta}$. By way of induction, suppose that a pair $(\xi_{d-1,1}, \bar{\xi}_{d-1,e+1})$ has been constructed such that $\xi_{d-1,1} : B \rightarrow \bar{\mathcal{M}}_{0,1}(X, 1)$ is very twisting, $\bar{\xi}_{d-1,e+1} : B \rightarrow \bar{\mathcal{M}}_{0,1}(X, e+1)$ is very positive and typical (in particular $\bar{\xi}_{d-1,e+1}(B)$ is contained in the locus of very stable maps), the map $h_{d-1,1} : B \rightarrow X$ equals the map $\bar{h}_{d-1,e+1} : B \rightarrow X$, and the degree conditions of Equation (9) hold for $d-1$. In particular, because $d \leq \bar{\delta}$, $\deg(\bar{\xi}_{d-1,e+1}^* \Delta_1) \geq 1$.

Let $b \in B$ be a closed point such that the stable map $(\bar{\xi}_{d-1,e+1})_b$ is in Δ_1 . By Definition 5.8, this stable map is of the form,

$$(\bar{\xi}_{d-1,e+1})_b = ((L \cup M, \bar{\sigma}_{d-1,e+1}(b)), (\bar{g}_{d-1,e+1})_b : L \cup M \rightarrow X),$$

where $\bar{\sigma}_{d-1,e+1}(b) \in L$ and $g|_L : L \rightarrow X$ is an isomorphism to a twistable line.

Denote by \tilde{B} the prestable curve of genus 0 containing B and L obtained by identifying the divisor $b \in B$ with the divisor $\bar{\sigma}_{d-1,e+1}(b) \in L$. Denote by,

$$\tilde{\xi}_{d-1,e+1} = ((\tilde{p}_{d-1,e+1} : \tilde{\Sigma}_{d-1,e+1} \rightarrow \tilde{B}, \tilde{\sigma}_{d-1,e+1}), \tilde{g}_{d-1,e+1} : \tilde{B} \rightarrow X),$$

the modification of $\bar{\xi}_{d-1,e+1}$ associated to $(b, L, \bar{\sigma}_{d-1,e+1}(b))$, cf. Definition 5.3.

By construction, for each divisor in $\bar{\mathcal{M}}_{0,1}(X, e+1)$, the difference of the intersection number with $\tilde{\xi}_{d-1,e+1}(\tilde{B})$ and the intersection number with $\bar{\xi}_{d-1,e+1}(B)$ equals the intersection number with $\tilde{\xi}_{d-1,e+1}(L)$, i.e.,

$$\begin{cases} \tilde{\xi}_{d-1,e+1}^* \mathcal{H} &= \bar{\xi}_{d-1,e+1}^* \mathcal{H}, \\ \tilde{\xi}_{d-1,e+1}^* \mathcal{L} &= \bar{\xi}_{d-1,e+1}^* \mathcal{L} + 1, \\ \tilde{\xi}_{d-1,e+1}^* \Delta_i &= \bar{\xi}_{d-1,e+1}^* \Delta_i, & i = 2, \dots, e-1, \\ \tilde{\xi}_{d-1,e+1}^* \Delta_1 &= \bar{\xi}_{d-1,e+1}^* \Delta_1 - 1, & i > 1, \\ \tilde{\xi}_{d-1,e+1}^* \Delta_e &= \bar{\xi}_{d-1,e+1}^* \Delta_e + 1, & i > 1, \\ \tilde{\xi}_{d-1,2}^* \Delta_1 &= \bar{\xi}_{d-1,e+1}^* \Delta_1, & i = 1. \end{cases}$$

Just a few remarks about this list: The second line arises because $\tilde{h}_{d-1,e+1} : L \rightarrow X$ equals $g|_L : L \rightarrow X$, which is an isomorphism to a line. The fourth line arises by applying Lemma 3.29

and using the fact that the pullback of $\mathcal{O}_{\overline{\mathcal{M}}_{0,1}(X,e+1)}(\Delta_1)$ equals

$$\sigma_{I,\lambda}^* \mathcal{O}_\Lambda(\sigma_{I,\lambda}(L)) \otimes (\sigma'_{I,\lambda})^* \mathcal{O}_{L \times M}(\sigma'_{I,\lambda}(L)),$$

in the notation of Subsection 5.1, where λ is the point in $L \cap M$. Of course the second factor in the tensor product is isomorphic to \mathcal{O}_L and the first factor is $\mathcal{O}_L(-\lambda)$ (because of the blowing up of $L \times L$ at (λ, λ)); hence the net degree change is -1 . Similarly, the fifth line arises because the intersection of $\tilde{\xi}_{d-1,e+1}(L)$ and Δ_e is exactly $\tilde{\xi}_{d-1,e+1}(\lambda)$; hence the degree increases by 1. In case $e = 1$, the last two contributions exactly cancel each other, which gives the last line.

As a ‘‘consistency check’’, substituting the computations above into the formula [19, Lem 2.2.2] for the self-intersection of $\tilde{\sigma}_{d-1,e+1}(\tilde{B}) \subset \tilde{\Sigma}_{d-1,e+1}$ yields a net change of $+1$. Of course this is correct because the change equals the self-intersection of $\sigma_{I,\Delta}(L) \subset \Lambda$, which is $T_L(-\lambda)$ (because of the blowing up of $L \times L$ at (λ, λ)), and this has degree $2 - 1 = 1$.

Claim 5.18. *The 1-morphism $\tilde{\xi}_{d-1,e+1} : \tilde{B} \rightarrow \overline{\mathcal{M}}_{0,1}(X, e+1)$ is very positive, the image is contained in the locus of stable maps with only the trivial automorphism, and the hypotheses of Lemma 5.6 are satisfied.*

First of all, the restriction of $\tilde{h}_{d-1,e+1} : \tilde{B} \rightarrow X$ to B is $\bar{h}_{d-1,e+1}$ and the restriction to L is $g|_L : L \rightarrow X$. By hypothesis, $\bar{h}_{d-1,e+1}$ is stable, and $g|_L$ is stable because it is a closed immersion. Therefore $\tilde{h}_{d-1,e+1}$ is stable. By hypothesis, the dimension of the obstruction group of $\bar{h}_{d-1,e+1}$ is 0. Moreover, possibly after deforming $\bar{\xi}_{d-1,e+1}$, it may be assumed that the node of $L \cap M$ maps to a very general point of X . Therefore $g|_L : L \rightarrow X$ is free by [7, Prop. 4.14]. So the dimension of the obstruction group of $g|_L$ is 0. Also, by Hypothesis 4.10 it may be assumed that $\text{ev} : \overline{\mathcal{M}}_{0,1}(X, 1) \rightarrow X$ is smooth at $((L, \bar{\sigma}_{d-1,e+1}(b)), g|_L : L \rightarrow X)$. So by Lemma 3.28, the dimension of the obstruction group of $\tilde{h}_{d-1,e+1}$ is 0. Hence (i) and (ii) of Definition 4.11 are satisfied.

By hypothesis, the dimension of the obstruction group of $\overline{\mathcal{M}}_{0,0}(X, e+1)$ at each point of $\text{pr}(\tilde{\xi}_{d-1,e+1}(B)) = \text{pr}(\bar{\xi}_{d-1,e+1}(B))$ is 0. And $\text{pr}(\tilde{\xi}_{d-1,e+1}(L))$ is the point $\text{pr}((\tilde{\xi}_{d-1,e+1})_b)$, which is one of the points above. Thus (iii) of Definition 4.11 is satisfied. Similarly, the restriction to B of the pullback,

$$(\text{pr} \circ \tilde{\xi}_{d-1,e+1})^* T_{\overline{\mathcal{M}}_{0,0}(X,e+1)},$$

equals the pullback associated to $\text{pr} \circ \bar{\xi}_{d-1,e+1}$, which is deformation ample by hypothesis. And the restriction to L is the pullback of a vector bundle by a constant map, hence it is isomorphic to a direct sum of copies of \mathcal{O}_L . In particular the restriction to L is generated by global sections. By Lemma 2.11, the pullback is deformation ample on all of \tilde{B} , hence (iv) of Definition 4.11 is satisfied.

Finally, the restriction to B of the pullback,

$$\tilde{\sigma}_{d-1,e+1}^* \mathcal{O}_{\tilde{\Sigma}_{d-1,e+1}}(\tilde{\sigma}_{d-1,e+1}(\tilde{B})),$$

is equal to the analogous sheaf for $\bar{\xi}_{d-1,e+1}$, and this is generated by global sections by construction. The restriction to L is the pullback by the diagonal of T_L , twisted down by $\bar{\sigma}_{d-1,e+1}(b)$, i.e., $\mathcal{O}_L(2-1) = \mathcal{O}_L(1)$. Thus by Lemma 2.11, the restriction to all of \tilde{B} is deformation ample. Hence (v) of Definition 4.11 is satisfied, therefore $\tilde{\xi}_{d-1,e+1}$ is very positive.

By hypothesis $\tilde{\xi}_{d-1,e+1}(B) = \bar{\xi}_{d-1,e+1}(B)$ is contained in the locus of stable maps that have only the trivial automorphism. In particular, $(\tilde{\xi}_{d-1,e+1})_b$ has only the trivial automorphism, from which it easily follows that $\tilde{\xi}_{d-1,e+1}(L)$ is contained in the locus of stable maps with only the trivial

automorphism. Therefore the image of all of \tilde{B} is contained in the locus of stable maps with only the trivial automorphism.

Because $\bar{\xi}_{d-1,e+1} : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e+1)$ is positive and the image is contained in the locus of very stable curves, the hypotheses of Lemma 3.22 are satisfied and there is a short exact sequence of locally free \mathcal{O}_B -modules,

$$0 \rightarrow \bar{\sigma}_{d-1,e+1}^* \mathcal{O}_{\bar{\Sigma}_{d-1,e+1}}(\bar{\sigma}_{d-1,e+1}(B)) \rightarrow \dots \\ \bar{\xi}_{d-1,e+1}^* T_{\overline{\mathcal{M}}_{0,1}(X,e+1)} \rightarrow (\text{pr} \circ \bar{\xi}_{d-1,e+1})^* T_{\overline{\mathcal{M}}_{0,0}(X,e+1)} \rightarrow 0.$$

By hypothesis, the third term is deformation ample, and the first term is generated by global sections. By Lemma 2.3 (iii), the pullback of $T_{\overline{\mathcal{M}}_{0,1}(X,e+1)}$ is generated by global sections. Because the node of $L \cap M$ is mapped to a very general point of X , by [7, Prop. 4.14], the pullback of T_X to L and the pullback to M are each generated by global sections; in particular the dimension of the obstruction group of $g|_L : L \rightarrow X$ is 0. Hence the hypotheses of Lemma 5.6 are satisfied; therefore Claim 5.18 is true.

Of course the next step will be to apply Lemma 5.6 to construct $\bar{\xi}_{d,e+1}$. But first the family $\tilde{\xi}_{d-1,1}$ is constructed. By hypothesis, the line L is twistable. Denote by

$$\xi_{L,1} = ((p_{L,1} : \Sigma_{L,1} \rightarrow L, \sigma_{L,1}), g_{L,1} : \Sigma_{L,1} \rightarrow X),$$

a twisting family such that $h_{L,1} = g|_L$. If X is of planar type, assume that the degree of $g_{L,1}$ is 1; otherwise the degree of $g_{L,1}$ is 2. By Hypotheses 1.6 and 4.10, Lemma 4.9 applies to $\xi_{d-1,1}$ and $\xi_{L,1}$, i.e., possibly after deforming the two families (without deforming $\tilde{h}_{d-1,e+1}$),

$$(\xi_{d-1,1})_b = (\xi_{L,1})_{\bar{\sigma}_{d-1,e+1}(b)}.$$

Define $\tilde{\xi}_{d-1,1} : \tilde{B} \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ to be the unique 1-morphism whose restriction to B is $\xi_{d-1,1}$ and whose restriction to L is $\xi_{L,1}$. By Lemma 4.5, $\tilde{\xi}_{d-1,1}$ is very twisting. By construction, $\tilde{h}_{d-1,1} = \tilde{h}_{d-1,e+1}$. Also,

$$\deg(\tilde{\xi}_{d-1,1}^* \mathcal{H}) = \deg(\xi_{d-1,1}^* \mathcal{H}) + \deg(g_{L,1}).$$

By definition, $\deg(g_{L,1})$ is 1 if X is planar type, and 2 if X is quadric type. Finally,

$$\deg(\tilde{\xi}_{d-1,1}^* \mathcal{L}) = \deg(\xi_{d-1,1}^* \mathcal{L}) + 1,$$

because $h_{L,1} : L \rightarrow X$ is an isomorphism to a line.

By Lemma 5.6, there exists a 1-morphism $\bar{\xi}_{d,e+1} : B \rightarrow \overline{\mathcal{M}}_{0,1}(X, e+1)$, i.e., both $\tilde{\xi}_{d-1,e+1}$ and $\bar{\xi}_{d,e+1}$ have a common generization. Since the image of $\tilde{\xi}_{d-1,e+1}$ is contained in the locus of stable maps with only the trivial automorphism, $\bar{\xi}_{d,e+1}$ can be chosen with the same property. Because the dimension of the obstruction group at every point of the image of $\tilde{\xi}_{d-1,e+1}$ equals 0, the same is true for $\bar{\xi}_{d,e+1}$. Hence the image of $\bar{\xi}_{d,e+1}$ is contained in the locus of $\overline{\mathcal{M}}_{0,1}(X, e+1)$ that is a smooth scheme. Because $\tilde{\xi}_{d-1,e+1}$ can be chosen to contain a very general point of $\overline{\mathcal{M}}_{0,1}(X, e+1)$ (because $\tilde{\xi}_{d-1,e+1}$ is typical), the same is true of $\bar{\xi}_{d,e+1}$. By [7, Prop. 4.14], the pullback $\bar{\xi}_{d,e+1}^* T_{\overline{\mathcal{M}}_{0,1}(X,e+1)}$ is generated by global sections. Therefore, after deforming further, $\bar{\xi}_{d,e+1}$ can be chosen to be typical. Because $\tilde{\xi}_{d-1,e+1}$ is very positive, by Lemma 4.13, $\bar{\xi}_{d,e+1}$ can be chosen to be very positive. Also, by the proof of Proposition 4.8, $\bar{\xi}_{d,e+1}$ can be chosen so that there exists a deformation $\xi_{d,1}$ of $\tilde{\xi}_{d-1,1}$ that is very twisting and such that $h_{d,1} = \bar{h}_{d,e+1}$.

Of course for any divisor in $\overline{\mathcal{M}}_{0,1}(X, e+1)$ (resp. $\overline{\mathcal{M}}_{0,1}(X, 1)$), the intersection number with $\overline{\xi}_{d,e+1}$ (resp. $\xi_{d,1}$) equals the intersection number with $\tilde{\xi}_{d-1,e+1}$ (resp. $\tilde{\xi}_{d-1,1}$), and these are computed above. This finishes the proof of the induction step. Hence, by induction on d , for each $d = 1, \dots, \bar{d}$, there exists an inducting pair $(\xi_{d,1}, \overline{\xi}_{d,e+1})$ as claimed. \square

6. TWISTABLE LINES ON HYPERSURFACES

In this section it is proved that if $n+1 \geq d^2$, if $X \subset \mathbb{P}^n$ is a general hypersurface of degree d , and if $L \subset X$ is a general line on X , then L is twistable.

Remark 6.1. There is one exceptional case, namely $d = 1$. For $n = 1, 2$, there is no twistable line on a hyperplane in \mathbb{P}^n , i.e., there is no twistable line on \mathbb{P}^{n-1} . For $n \geq 3$ there is a twistable line on \mathbb{P}^{n-1} . In this section it is proved there is a twistable line on \mathbb{P}^{n-1} if $n \geq 4$. Let $((\pi : \Sigma \rightarrow B, \sigma), g)$ be a twisting family in \mathbb{P}^n such that $h : B \rightarrow \mathbb{P}^n$ is a line, and let $p \in \mathbb{P}^n - g(\Sigma)$. Denote by $\text{proj}_p : \mathbb{P}^n \dashrightarrow \mathbb{P}^{n-1}$ projection from p . Then $((\pi : \Sigma \rightarrow B, \sigma), \text{proj}_p \circ g)$ is a twisting family in \mathbb{P}^{n-1} and $\text{proj}_p \circ h : B \rightarrow \mathbb{P}^{n-1}$ is a line. Therefore there is also a twistable line on \mathbb{P}^2 .

Notation 6.2. Denote $N_d = \binom{n+d}{n} - 1$ and denote by \mathbb{P}^{N_d} the projective space parametrizing hypersurfaces $X \subset \mathbb{P}^n$ of degree d . Denote by $\mathcal{X} \subset \mathbb{P}^{N_d} \times \mathbb{P}^n$ the universal family of degree d hypersurfaces in \mathbb{P}^n . Denote by $\mathbb{G}(1, n)$ the Grassmannian variety of lines in \mathbb{P}^n . Denote by $F(\mathcal{X}) \subset \mathbb{P}^{N_d} \times \mathbb{G}(1, n)$ the parameter space of pairs $([X], [L])$ consisting of a hypersurface of degree d , $X \subset \mathbb{P}^n$, and a line $L \subset X$. Denote by $P(t) = (t+1)^2$ the Hilbert polynomial of a quadric surface in \mathbb{P}^3 . Denote by $U \subset \text{Hilb}_{\mathbb{P}^n}^{P(t)}$ the open subscheme of the Hilbert scheme that parametrizes subschemes of \mathbb{P}^n projectively equivalent to a smooth quadric surface in $\mathbb{P}^3 \subset \mathbb{P}^n$. Denote by $V \subset U \times \mathbb{G}(1, n)$ the parameter space of pairs $([\Sigma], [L])$ consisting of a smooth quadric surface, Σ , and a line $L \subset \Sigma$. Denote by $V \rightarrow \tilde{U} \rightarrow U$ the Stein factorization of the projection $\text{pr}_U : V \rightarrow U$. Denote by $W \subset \mathbb{P}^{N_d} \times U \times \mathbb{G}(1, n)$ the parameter space of triples $([X], [\Sigma], [L])$ consisting of a hypersurface of degree d , $X \subset \mathbb{P}^n$, a smooth quadric surface $\Sigma \subset X$, and a line $L \subset \Sigma$.

Observe that the projection $F(\mathcal{X}) \rightarrow \mathbb{G}(1, d)$ is a projective bundle of relative dimension $N_d - (d+1)$. Observe that $\tilde{U} \rightarrow U$ is a finite, étale morphism of degree 2. Observe that $V \rightarrow \tilde{U}$ is a \mathbb{P}^1 -bundle. Observe that $W \rightarrow V$ is a projective space bundle of relative dimension $N_d - (d+1)^2$.

Let $([X], [\Sigma], [L])$ be a triple in W . There is a map (well-defined up to nonzero scalar) $\partial_X : \mathbb{C}^{n+1} \rightarrow H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d-1))$ that evaluates the partial derivatives of a defining equation of X . Compose this map with the restriction map $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d-1)) \rightarrow H^0(\Sigma, \mathcal{O}_{\Sigma}(d-1))$, and denote the composition by,

$$\partial_{X,\Sigma} : \mathbb{C}^{n+1} \rightarrow H^0(\Sigma, \mathcal{O}_{\Sigma}(d-1)).$$

Denote $E = \mathcal{O}_W^{\oplus(n+1)}$. Denote by G the unique quotient of $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d-1)) \otimes_{\mathbb{C}} \mathcal{O}_U$ that is locally free and whose fiber at each point Σ is the quotient,

$$H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d-1)) \rightarrow H^0(\Sigma, \mathcal{O}_{\Sigma}(d-1)).$$

Denote by F the locally free \mathcal{O}_W -module,

$$F = \text{pr}_{\mathbb{P}^{N_d}}^* \mathcal{O}_{\mathbb{P}^{N_d}}(1) \otimes \text{pr}_U^* G.$$

There is a map of \mathcal{O}_W -modules, $\partial : E \rightarrow F$ whose fiber at each point $([X], [\Sigma], [L])$ is the map $\partial_{X,\Sigma}$. Denote by $W^o \subset W$ the open subscheme that is the complement of the support of $\text{Coker}(\partial)$, i.e., W^o is the maximal open subscheme on which ∂ is surjective.

Lemma 6.3. *Let $([X], [\Sigma], [L])$ be a point in W^o . Then,*

- (i) X is smooth along Σ ,

- (ii) $h^i(\Sigma, N_{\Sigma/\mathbb{P}^n}) = h^i(\Sigma, N_{\Sigma/\mathbb{P}^n}(-L)) = h^i(\Sigma, N_{\Sigma/\mathbb{P}^n}(-1)) = 0$, for $i > 0$,
- (iii) $h^i(\Sigma, N_{\Sigma/X}) = h^i(\Sigma, N_{\Sigma/X}(-L)) = h^i(\Sigma, N_{\Sigma/X} \otimes \mathcal{O}_{\Sigma}(-1)) = 0$, for $i > 0$,
- (iv) $h^1(L, N_{L/X}(-1)) = h^1(L, N_{L/X}) = 0$,
- (v) the projection morphism $pr_{\mathbb{P}^{N_d}} : W \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [\Sigma], [L])$,
- (vi) the projection morphism $pr_{\mathbb{P}^{N_d}} : F(\mathcal{X}) \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [L])$, and
- (vii) the projection morphism $pr_{F(\mathcal{X})} : W \rightarrow F(\mathcal{X})$ is smooth at $([X], [\Sigma], [L])$.

Proof. (i): Since the partial derivatives of a defining equation of X generate $H^0(\Sigma, \mathcal{O}_{\Sigma}(d-1))$, the subscheme of X where the partial derivatives all vanish is disjoint from Σ . By the Jacobian criterion, X is smooth at each point of Σ .

(ii): Denote by $\mathbb{P}^3 \subset \mathbb{P}^n$ the span of Σ . There is a short exact sequence,

$$0 \longrightarrow N_{\Sigma/\mathbb{P}^3} \longrightarrow N_{\Sigma/\mathbb{P}^n} \longrightarrow N_{\mathbb{P}^3/\mathbb{P}^n}|_{\Sigma} \longrightarrow 0.$$

Since $N_{\Sigma/\mathbb{P}^3} \cong \mathcal{O}_{\Sigma}(2)$ and since $N_{\mathbb{P}^3/\mathbb{P}^n}|_{\Sigma} \cong \mathcal{O}_{\Sigma}(1)^{\oplus(n-3)}$, the short exact sequence above is,

$$0 \longrightarrow \mathcal{O}_{\Sigma}(2) \longrightarrow N_{\Sigma/\mathbb{P}^n} \longrightarrow \mathcal{O}_{\Sigma}(1)^{\oplus(n-3)} \longrightarrow 0.$$

From this it is easy to compute that,

$$h^i(\Sigma, N_{\Sigma/\mathbb{P}^n}) = h^i(\Sigma, N_{\Sigma/\mathbb{P}^n} \otimes \mathcal{O}_{\Sigma}(-1)) = h^i(\Sigma, N_{\Sigma/\mathbb{P}^n} \otimes \mathcal{O}_{\Sigma}(-L)) = 0,$$

for $i > 0$.

(iii): There is a short exact sequence,

$$0 \longrightarrow N_{\Sigma/X} \longrightarrow N_{\Sigma/\mathbb{P}^n} \longrightarrow N_{X/\mathbb{P}^n}|_{\Sigma} \longrightarrow 0.$$

Of course $N_{X/\mathbb{P}^n}|_{\Sigma} \cong \mathcal{O}_{\Sigma}(d)$. For the three cases $\mathcal{L} = \mathcal{O}_{\Sigma}$, $\mathcal{L} = \mathcal{O}_{\Sigma}(-L)$, and $\mathcal{L} = \mathcal{O}_{\Sigma}(-1)$, $h^i(\Sigma, \mathcal{O}_{\Sigma}(d) \otimes \mathcal{L}) = 0$ for $i > 0$. By (ii), $h^i(\Sigma, N_{\Sigma/\mathbb{P}^n} \otimes \mathcal{L}) = 0$ for $i > 0$. By the long exact sequence in cohomology associated to the twist by \mathcal{L} of the short exact sequence above, $h^i(\Sigma, N_{\Sigma/X} \otimes \mathcal{L}) = 0$ for $i > 2$. Also $h^1(\Sigma, N_{\Sigma/X} \otimes \mathcal{L}) = 0$ iff the map $e_{\mathcal{L}} : H^0(\Sigma, N_{\Sigma/\mathbb{P}^n} \otimes \mathcal{L}) \rightarrow H^0(\Sigma, N_{X/\mathbb{P}^n}|_{\Sigma})$ is surjective.

The map $e_{\mathcal{O}_{\Sigma}}$ factors the map $\partial_{X,\Sigma}$. Since $\partial_{X,\Sigma}$ is surjective, also $e_{\mathcal{O}_{\Sigma}}$ is surjective and $h^1(\Sigma, N_{\Sigma/X}) = 0$.

For the case $\mathcal{L} = \mathcal{O}_{\Sigma}(-L)$, observe that $\mathcal{O}_{\Sigma}(1) \cong \mathcal{O}_{\Sigma}(L+L')$ where $L' \subset \Sigma$ is a line of the ruling opposite to L . There is a commutative diagram,

$$\begin{array}{ccc} H^0(\Sigma, N_{\Sigma/\mathbb{P}^n}(-1)) \otimes_{\mathbb{C}} H^0(\Sigma, \mathcal{O}_{\Sigma}(L')) & \longrightarrow & H^0(\Sigma, N_{\Sigma/\mathbb{P}^n}(-L)) \\ \downarrow & & \downarrow \\ H^0(\Sigma, N_{X/\mathbb{P}^n}|_{\Sigma}(-1)) \otimes_{\mathbb{C}} H^0(\Sigma, \mathcal{O}_{\Sigma}(L')) & \longrightarrow & H^0(\Sigma, N_{X/\mathbb{P}^n}|_{\Sigma}(-L)) \end{array}$$

The left vertical arrow is surjective by (ii). The bottom arrow is

$$H^0(\Sigma, \mathcal{O}_{\Sigma}((d-1)L + (d-1)L')) \otimes H^0(\Sigma, \mathcal{O}_{\Sigma}(L')) \rightarrow H^0(\Sigma, \mathcal{O}_{\Sigma}((d-1)L + dL')),$$

which is surjective. Therefore also the right vertical arrow is also surjective, i.e., $h^1(\Sigma, N_{\Sigma/X}(-L)) = 0$. The proof that $h^1(\Sigma, N_{\Sigma/X}) = 0$ is almost identical to the proof that $h^1(\Sigma, N_{\Sigma/X}(-L)) = 0$.

(iv) There is a short exact sequence,

$$0 \longrightarrow N_{\Sigma/X}(-1) \longrightarrow N_{\Sigma/X}(-L') \longrightarrow N_{\Sigma/X}|_L(-1) \longrightarrow 0.$$

By the associated long exact sequence in cohomology and the computations above, $h^1(L, N_{\Sigma/X}|_L(-1)) = 0$. There is a short exact sequence,

$$0 \longrightarrow N_{L/\Sigma}(-1) \longrightarrow N_{L/X}(-1) \longrightarrow N_{\Sigma/X}|_L(-1) \longrightarrow 0.$$

Of course $N_{L/\Sigma} \cong \mathcal{O}_L(1)$, so $h^1(L, N_{L/\Sigma}(-1)) = 0$. As proved above, $h^1(L, N_{\Sigma/X}|_L(-1)) = 0$. Therefore by the long exact sequence in cohomology, $h^1(L, N_{L/X}(-1)) = 0$. Since $N_{L/X}$ is a locally free \mathcal{O}_L -module and $h^1(L, N_{L/X}(-1)) = 0$, by Grothendieck's lemma $N_{L/X}$ is generated by global sections. In particular, also $h^1(L, N_{L/X}) = 0$.

(v): By [16, Prop. I.2.14.2], the obstruction space for the relative Hilbert scheme, $\text{Hilb}_{\mathcal{X}/\mathbb{P}^{N_d}}^{P(t)}$, at the point $([X], [\Sigma])$ is contained in $H^1(\Sigma, N_{\Sigma/X})$, which has dimension 0 by (iv). Since the obstruction space vanishes, it follows by [16, Thm. 2.10] that $\text{Hilb}_{\mathcal{X}/\mathbb{P}^{N_d}}^{P(t)} \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [\Sigma])$. Also $W \rightarrow \text{Hilb}_{\mathcal{X}/\mathbb{P}^{N_d}}^{P(t)}$ is smooth. Therefore the composition $\text{pr}_{\mathbb{P}^{N_d}} : W \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [\Sigma], [L])$.

(vi): The proof is almost identical to the proof of (v).

(vii): By (v), $W \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [\Sigma], [L])$. By (vi), $F(\mathcal{X}) \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [L])$. So the Jacobian criterion for the smoothness of $\text{pr}_{F(\mathcal{X})}$ at $([X], [\Sigma], [L])$ is that the map of vertical tangent bundles, $d\text{pr}_{F(\mathcal{X})} : T_{W/\mathbb{P}^{N_d}} \rightarrow \pi^* T_{F(\mathcal{X})/\mathbb{P}^{N_d}}$ is surjective at $([X], [\Sigma], [L])$. This reduces to the surjectivity of $H^0(\Sigma, N_{\Sigma/X}) \rightarrow H^0(L, N_{\Sigma/X}|_L)$. The cokernel is contained in $H^1(\Sigma, N_{\Sigma/X}(-L))$, which is zero by (iii). Therefore $\text{pr}_{F(\mathcal{X})}$ is smooth at $([X], [\Sigma], [L])$. \square

We associate to each $([X], [\Sigma], [L]) \in W^\circ$ a morphism $\zeta : L \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ as follows. Let $\sigma : L \rightarrow \Sigma$ be the inclusion and let $\text{pr}_L : \Sigma \rightarrow L$ be the unique projection such that σ is a section of pr_L . Let $g : \Sigma \rightarrow X$ be the inclusion. Then,

$$\zeta = ((\text{pr}_L : \Sigma \rightarrow L, \sigma), g : \Sigma \rightarrow X),$$

is a family of stable maps, i.e., a morphism $\zeta : L \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$.

Lemma 6.4. *For every point $([X], [\Sigma], [L]) \in W^\circ$, the morphism $\zeta : L \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ is twisting.*

Proof. Since $g \circ \sigma : L \rightarrow X$ is an embedding, Axiom (i) of Definition 4.3 is satisfied. By Lemma 6.3 (vi), the dimension of the obstruction group of $\overline{\mathcal{M}}_{0,0}(X, 1)$ at $[g \circ \sigma : L \rightarrow X]$ is 0, i.e., Axiom (ii) of Definition 4.3 is satisfied.

Denote by \mathcal{N} the normal bundle of the regular embedding $(\text{pr}_L, g) : \Sigma \rightarrow L \times X$. There is a short exact sequence,

$$0 \longrightarrow \text{pr}_L^* T_L \longrightarrow \mathcal{N} \longrightarrow N_{\Sigma/X} \longrightarrow 0.$$

By Remark 4.4 (ii), Axiom (iii) holds if $R^1(\text{pr}_L)_* \mathcal{N}(-\sigma) = \{0\}$. For each fiber L' of $\text{pr}_L : \Sigma \rightarrow L$, $\mathcal{N}(-\sigma)|_{L'} \cong N_{L'/X}(-1)$. Since $([X], [\Sigma], [L'])$ is also in W° , by Lemma 6.3 (iv), $h^1(L', N_{L'/X}(-1)) = 0$. Therefore $R^1(\text{pr}_L)_* \mathcal{N}(-\sigma) = \{0\}$, i.e., Axiom (iii) holds.

By Remark 4.4 (ii), $\zeta^* T_{\text{ev}} \cong (\text{pr}_L)_* \mathcal{N}(-\sigma)$. Part of the long exact sequence of higher direct images associated to the twist by $\mathcal{O}_\Sigma(-L)$ of the short exact sequence above is,

$$(\text{pr}_L)_* \text{pr}_L^* T_L(-\sigma(L)) \rightarrow (\text{pr}_L)_* \mathcal{N}(-\sigma(L)) \rightarrow (\text{pr}_L)_* N_{\Sigma/X}(-\sigma(L)) \rightarrow R^1(\text{pr}_L)_* \text{pr}_L^* T_L(-\sigma(L)) \rightarrow 0.$$

For each fiber L' of pr_L , $T_L(-\sigma(L))|_{L'} \cong \mathcal{O}_{L'}(-1)$. Therefore $(\text{pr}_L)_* \text{pr}_L^* T_L(-\sigma(L)) = \{0\}$ and $R^1(\text{pr}_L)_* \text{pr}_L^* T_L(-\sigma(L)) = \{0\}$, i.e., $(\text{pr}_L)_* \mathcal{N}(-\sigma(L)) \cong (\text{pr}_L)_* N_{\Sigma/X}(-\sigma(L))$.

Let $p' \in L$ be a closed point and denote $L' = \text{pr}_L^{-1}\{p'\}$. Since $R^1(\text{pr}_L)_* \text{pr}_L^* T_L(-L - L') = \{0\}$ and $R^1(\text{pr}_L)_* \mathcal{N}(-L - L') = \{0\}$, by the long exact sequence of higher direct images, also $R^1(\text{pr}_L)_* N_{\Sigma/X}(-L - L') = \{0\}$. By the Leray spectral sequence, $h^1(\Sigma, N_{\Sigma/X}(-L - L')) = h^1(L, (\text{pr}_L)_*(N_{\Sigma/X}(-\sigma(L)))(-p'))$. By Lemma 6.3 (iii), $h^1(\Sigma, N_{\Sigma/X}(-L - L')) = 0$, so also $h^1(L, (\text{pr}_L)_*(N_{\Sigma/X}(-\sigma(L)))(-p')) = 0$. By Grothendieck's lemma, $(\text{pr}_L)_* N_{\Sigma/X}(-\sigma(L))$ is a direct sum of line bundles of degree ≥ 0 , i.e., it is generated by global sections. Therefore Axiom (iv) is satisfied. Finally, $\sigma^* \mathcal{O}_\Sigma(\sigma(L)) \cong \mathcal{O}_L$, so Axiom (v) is satisfied. \square

Proposition 6.5. *If either,*

- (i) $d = 1$ and $n \geq 4$, or
- (ii) $d \geq 2$ and $n + 1 \geq d^2$,

then $\text{pr}_{F(\mathcal{X})} : W^o \rightarrow F(\mathcal{X})$ is dominant. Therefore, for a general pair $([X], [L]) \in F(\mathcal{X})$, L is a twistable line on X .

Proof. By Lemma 6.3 (v), it suffices to prove that W^o is nonempty.

(i): If $d = 1$ and $n \geq 4$, then for any quadric surface $\Sigma \subset \mathbb{P}^n$ and any hyperplane $X \subset \mathbb{P}^n$ containing $\text{span}(\Sigma)$, ∂ is an isomorphism so $([X], [\Sigma], [L])$ is in W^o for any line $L \subset \Sigma$.

(ii): Next suppose that $d \geq 2$. Denote by I_d the set,

$$I_d = \{(i, j) \in \mathbb{Z} \times \mathbb{Z} : 0 \leq i, j \leq d-1, i+j \geq 3\},$$

which has $d^2 - 4$ elements. Denote by,

$$(Y_0, Y_1, Y_2, Y_3) \cup (X_{i,j})_{(i,j) \in I_d} \cup (Z_m : m = 1, \dots, n+1-d^2),$$

a basis of $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$, i.e., a basis of homogeneous coordinates on \mathbb{P}^n . Denote by $\Sigma \subset \mathbb{P}^n$ the smooth quadric surface with ideal,

$$I_\Sigma = \langle Y_0 Y_3 - Y_1 Y_2 \rangle + \langle X_{i,j} \mid (i, j) \in I_d \rangle + \langle Z_m \mid m = 1, \dots, n+1-d^2 \rangle.$$

This is the image of the closed immersion $f : \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^n$,

$$([U_0 : U_1], [V_0 : V_1]) \mapsto [U_0 V_0 : U_0 V_1 : U_1 V_0 : U_1 V_1 : 0 : \dots : 0].$$

For each $(i, j) \in I_d$, denote $k = \min(i, j)$, denote $i' = i - k$, and denote $j' = j - k$. Denote by $X \subset \mathbb{P}^n$ the hypersurface with defining equation,

$$F = (Y_0 Y_3 - Y_1 Y_2) Y_3^{d-2} + \sum_{(i,j) \in I_d} Y_0^k Y_1^{i'} Y_2^{j'} Y_3^{d+k-i-j-1} X_{i,j}.$$

Clearly $\Sigma \subset X$. The claim is that $\partial F : \mathbb{C}^{n+1} \rightarrow H^0(\Sigma, \mathcal{O}_\Sigma(d-1))$ is surjective. By construction,

$$\frac{\partial F}{\partial Y_0} \mapsto U_1^{d-1} V_1^{d-1}, \quad \frac{\partial F}{\partial Y_1} \mapsto U_1^{d-1} V_0 V_1^{d-2}, \quad \frac{\partial F}{\partial Y_2} \mapsto U_0 U_1^{d-2} V_1^{d-1}, \quad \frac{\partial F}{\partial Y_3} \mapsto U_0 U_1^{d-2} V_0 V_1^{d-2}.$$

For each $(i, j) \in I_d$,

$$\frac{\partial F}{\partial X_{i,j}} \mapsto U_0^i U_1^{d-1-i} V_0^j V_1^{d-1-j}. \quad (12)$$

Since the partial derivatives of the form $\frac{\partial F}{\partial Y_i}$ give the terms $U_0^i U_1^{d-1-i} V_0^j V_1^{d-1-j}$ with $(i, j) = (0, 0), (0, 1), (1, 0)$, and $(1, 1)$, and since these are precisely the pairs (i, j) not contained in I_d , ∂F is surjective. Therefore, for every line $L \subset \Sigma$, $([X], [\Sigma], [L])$ is in W^o . \square

Together with Remark 6.1, Lemma 6.4 and Proposition 6.5 imply the following corollary.

Corollary 6.6. *If $X \subset \mathbb{P}^n$ is a general hypersurface of degree d and either,*

- (i) $d = 1$ and $n \geq 3$, or
- (ii) $d \geq 2$ and $n + 1 \geq d^2$,

then Hypothesis 4.10 holds.

7. BASE CASE OF THE INDUCTION FOR HYPERSURFACES

In this section it is proved that if $n \geq d^2 + d + 1$ and if $X \subset \mathbb{P}^n$ is a general hypersurface of degree d , then there exists a morphism $\bar{\zeta}_1 : \mathbb{P}^1 \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ that is both very twisting and very positive. This provides the base case for the induction argument of Section 5.

Remark 7.1. There is one exceptional case, $d = 1$. It will be proved that for $n \geq 7$, there is a morphism $\bar{\zeta}_1 : \mathbb{P}^1 \rightarrow \overline{\mathcal{M}}_{0,1}(\mathbb{P}^{n-1}, 1)$ that is both very twisting and very positive. As in Remark 6.1, repeatedly projecting from a point produces a morphism $\bar{\zeta}_1 : \mathbb{P}^1 \rightarrow \overline{\mathcal{M}}_{0,1}(\mathbb{P}^{n-1}, 1)$ that is very twisting and very positive for all $n \geq 3$.

The techniques in this section are the same as those of Section 6. Proposition 6.5 is proved by finding a single degree d polynomial F on \mathbb{P}^n , vanishing on some quadric surface Σ , and such that

$$\partial_{F,\Sigma} : \mathbb{C}^{n+1} \rightarrow H^0(\Sigma, \mathcal{O}_\Sigma(d-1))$$

is surjective. In this section, the role of $L \subset X$ is replaced by a rational normal curve $C_0 \subset X$ of degree $k \leq n$ (in the end, only the case $k = 2d$ will be needed). The role of the quadric surface is replaced by a rational normal scroll Σ of degree $2k - 1$ such that $C_0 \subset \Sigma \subset X$. The cohomology vanishing results of the last section are replaced by the vanishing of $h^i(\Sigma, N_{\Sigma/X}(-C_0 - 2L))$ for $i > 0$, where L is a line of the ruling of Σ . The computation in this section will be to find a single degree d polynomial F on \mathbb{P}^n , vanishing on Σ , and such that the image, W , of the map,

$$\partial_{F,\Sigma} : \mathbb{C}^{n+1} \rightarrow H^0(\Sigma, \mathcal{O}_\Sigma(d-1)),$$

has the property that the induced map,

$$W \otimes H^0(\Sigma, \mathcal{O}_\Sigma((k-3)L)) \rightarrow H^0(\Sigma, \mathcal{O}_\Sigma(d-1) \otimes \mathcal{O}_\Sigma((k-3)L)),$$

is surjective. A similar polynomial F to that of the last section satisfies this condition.

7.1. Generating linear systems on \mathbb{F}_1 . In the last section, the relevant surface was the Hirzebruch surface $\mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$ embedded as a quadric surface. In this section, the relevant surface is the Hirzebruch surface \mathbb{F}_1 embedded as a rational normal scroll of degree $2k - 1$. The projective model of \mathbb{F}_1 used here is,

$$\mathbb{F}_1 = \{([T_0 : T_1], [T_0U : T_1U : V]) \in \mathbb{P}^1 \times \mathbb{P}^2 \mid T_0(T_1U) = T_1(T_0U)\}.$$

In the equation above, “ T_0U ” and “ T_1U ” are just names of homogeneous coordinates on \mathbb{P}^2 (although the term U does have a meaning described below). Denote by $\text{pr}_{\mathbb{P}^1} : \mathbb{F}_1 \rightarrow \mathbb{P}^1$ and $\text{pr}_{\mathbb{P}^2} : \mathbb{F}_1 \rightarrow \mathbb{P}^2$ the projection morphisms. Denote by $\mathcal{O}_{\mathbb{F}_1}(F)$ the invertible sheaf $\text{pr}_{\mathbb{P}^1}^* \mathcal{O}_{\mathbb{P}^1}$ and by $\mathcal{O}_{\mathbb{F}_1}(E + F)$ the invertible sheaf $\text{pr}_{\mathbb{P}^2}^* \mathcal{O}_{\mathbb{P}^2}$. The invertible sheaf $\mathcal{O}_{\mathbb{F}_1}(E)$ is associated to the *directrix* $E \subset \mathbb{F}_1$. (This explains the terminology T_0U and T_1U ; U is a nonzero element of $H^0(\mathbb{F}_1, \mathcal{O}_{\mathbb{F}_1}(E))$, and T_0U and T_1U are the products of U with the two global sections T_0 and T_1 of $H^0(\mathbb{F}_1, \mathcal{O}_{\mathbb{F}_1}(F))$.)

The invertible sheaves $\mathcal{O}_{\mathbb{F}_1}(E + F)$ and $\mathcal{O}_{\mathbb{F}_1}(F)$ generate the Picard group of \mathbb{F}_1 ; thus motivating the notation,

$$\mathcal{O}(a, b) := \mathcal{O}_{\mathbb{F}_1}(a(E + F) + bF).$$

The divisors $E + F$ and F are each nef, but not ample. Therefore they generate the nef cone; an invertible sheaf $\mathcal{O}(a, b)$ is nef iff $0 \leq a, b$, and it is ample iff $0 < a, b$.

Let $\mathcal{O}(a, b)$ be a nef invertible sheaf, and let $W \subset H^0(\mathbb{F}_1, \mathcal{O}(a, b))$ be a linear system. Let $c \geq 0$ be an integer.

Definition 7.2. The linear system W is a *c-generating linear system* if the associated map,

$$\mu_{W,c} : W \otimes H^0(\mathbb{F}_1, \mathcal{O}(0, c)) \rightarrow H^0(\mathbb{F}_1, \mathcal{O}(a, b + c)),$$

is surjective.

When is W a *c-generating linear system*?

Notation 7.3. For each integer i , denote $r(i) = \lfloor \frac{b+i-1}{c+1} \rfloor$. Denote by $\beta_d, \beta_r, \alpha_d$ and α_r the unique integers such that $0 \leq \alpha_r, \beta_r < c+1$ and $b-1 = \beta_d(c+1) + \beta_r$, $a+b-1 = \alpha_d(c+1) + \alpha_r$. Denote by $N(a, b, c)$ the integer,

$$N(a, b, c) = \sum_{i=0}^a (r(i)+2) = 2a+2 + \begin{cases} \beta_d(c+1 - \beta_r) + \alpha_d(1 + \alpha_r) + (\alpha_d - \beta_d - 1)(c+1) & \alpha_d > \beta_d, \\ \beta_d(\alpha_r - \beta_r + 1) & \alpha_d = \beta_d. \end{cases}$$

Denote by $W_0(a, b, c) \subset H^0(\mathbb{F}_1, \mathcal{O}(a, b))$ the linear system,

$$W_0(a, b, c) = \text{span}\{U^i V^{a-i} T_0^{(b+i)-j(c+1)} T_1^{j(c+1)} \mid i = 0, \dots, a, j = 0, \dots, r(i)\} \\ + \text{span}\{U^i V^{a-i} T_1^{b+i} \mid i = 0, \dots, a\}.$$

Lemma 7.4. *The linear system $W_0(a, b, c)$ is a c-generating linear system of dimension $N(a, b, c)$.*

Proof. For each pair of nonnegative integers a', b' there is a decreasing filtration on $H^0(\mathbb{F}_1, \mathcal{O}(a', b'))$,

$$F^i H^0(\mathbb{F}_1, \mathcal{O}(a', b')) = H^0(\mathbb{F}_1, \mathcal{O}(a', b')(-iE)) \cong H^0(\mathbb{F}_1, \mathcal{O}(a' - i, b')).$$

For any linear system $W \subset H^0(\mathbb{F}_1, \mathcal{O}(a, b))$, there is an induced filtration $F^i W = F^i \cap W$. The multiplication map $\mu_{W,c}$ respects the filtrations on W and on $H^0(\mathbb{F}_1, \mathcal{O}(a, b + c))$. Hence $\mu_{W,c}$ is surjective if every associated graded map,

$$\text{gr}^i \mu_{W,c} : \text{gr}^i W \otimes H^0(\mathbb{F}_1, \mathcal{O}(0, c)) \rightarrow \text{gr}^i H^0(\mathbb{F}_1, \mathcal{O}(a, b + c)),$$

is surjective. Now $\dim(W)$ is the sum of all terms $\dim(\text{gr}^i W)$. For each i , what is the minimum possible dimension of a vector subspace $W^i \subset \text{gr}^i H^0(\mathbb{F}_1, \mathcal{O}(a, b))$ such that the associated map,

$$\text{gr}^i \mu_{W^i,c} : W^i \otimes H^0(\mathbb{F}_1, \mathcal{O}(0, c)) \rightarrow \text{gr}^i H^0(\mathbb{F}_1, \mathcal{O}(a, b + c)),$$

is surjective?

The associated graded pieces of $\mathcal{O}(a', b')$ are,

$$\text{gr}^i H^0(\mathbb{F}_1, \mathcal{O}(a', b')) \cong \begin{cases} H^0(E, \mathcal{O}_E(b' + i)), & 0 \leq i \leq a' \\ \{0\}, & i > a' \end{cases}.$$

Let $W^i \subset H^0(E, \mathcal{O}_E(b + i))$ be a linear system such that the multiplication map,

$$\text{gr}^i \mu_{W^i,c} : W^i \otimes H^0(E, \mathcal{O}_E(c)) \rightarrow H^0(E, \mathcal{O}_E(b + c + i)),$$

is surjective. Counting dimensions on the left and right side of the equation, $\dim(W^i) \cdot (c+1) \geq (b + c + i + 1)$, i.e.,

$$\dim(W^i) \geq \left\lfloor \frac{b+i-1}{c+1} \right\rfloor + 2 = r(i) + 2.$$

The linear system $W^i = \text{gr}^i W_0(a, b, c)$, which is generated by the set of monomials

$$\{U^i V^{a-i} T_0^{(b+i)-j(c+1)} T_1^{j(c+1)} \mid j = 0, \dots, r(i)\} \cup \{U^i V^{a-i} T_1^{b+i}\},$$

has the property that $W^i \otimes H^0(E, \mathcal{O}_E(c)) \rightarrow H^0(E, \mathcal{O}_E(b+c+i))$ is surjective. And $\dim(W^i) = r(i) + 2$. So $W_0(a, b, c)$ is a c -generating linear system of dimension,

$$2a + 2 + \sum_{i=0}^a \left\lfloor \frac{b+i-1}{c+1} \right\rfloor = N(a, b, c).$$

Moreover, this is the minimum dimension among c -generating linear systems for which each map $\text{gr}^i \mu_{W^i, c}$ is surjective. \square

Notation 7.5. Denote by S the Cox homogeneous coordinate ring of \mathbb{F}_1 , i.e.,

$$S := \bigoplus_{(a,b) \in \mathbb{Z}^2} H^0(\mathbb{F}_1, \mathcal{O}(a, b)) = \mathbb{C}[T_0, T_1, U, V].$$

This is a \mathbb{Z}^2 -graded ring, where $\deg(T_0) = \deg(T_1) = (0, 1)$, $\deg(V) = (1, 0)$ and $\deg(U) = (1, -1)$. For every $(a, b) \in \mathbb{Z}^2$, $S_{(a,b)} = H^0(\mathbb{F}_1, \mathcal{O}(a, b))$. Denote by $\deg : S \rightarrow \mathbb{Z} \cup \{-\infty\}$ the total degree defined by $\deg(M) = a + b$ for all elements $M \in S_{(a,b)}$. Denote by \prec the graded lexicographical monomial order on S that refines the grading by total degree by $U > V > T_0 > T_1$. For every linear system $W \subset H^0(\mathbb{F}_1, \mathcal{O}(a, b))$ denote by $\text{IN}(W)$ the linear system generated by the initial terms of W .

Lemma 7.6. *If the linear system $\text{IN}(W)$ contains $W_0(a, b, c)$, then W is a c -generating linear system.*

Proof. The linear system of initial terms of $\text{Image}(\mu_{W,c})$ satisfies

$$\text{IN}(W) \cdot S_{(0,c)} \subset \text{IN}(\text{Image}(\mu_{W,c})).$$

Since $\text{IN}(W)$ contains $W_0(a, b, c)$,

$$W_0(a, b, c) \cdot S_{(0,c)} \subset \text{IN}(\text{Image}(\mu_{W,c})).$$

By Lemma 7.4, $W_0(a, b, c) \cdot S_{(0,c)} = S_{(a,b+c)}$. Hence $\text{IN}(\text{Image}(\mu_{W,c})) = S_{(a,b+c)}$, and therefore $\text{Image}(\mu_{W,c}) = S_{(a,b+c)}$. \square

Remark 7.7. The most important case is $a = d - 1$, $b = (d - 1)(k - 1)$ and $c = k - 3$ for positive integers $d \geq 1$ and $k \geq 3$ (d will be the degree of the hypersurface $X \subset \mathbb{P}^n$, and k will be the degree of the curve $C_0 \subset X$). In particular, if $d \geq 2$ and $k \geq 2d$, then $b - 1 = (d - 1)(k - 2) + d - 2$, $a + b - 1 = (d - 1)(k - 2) + 2d - 3$ and $0 \leq d - 2$, $2d - 3 \leq k - 3$. Hence $r(i) = d - 1$ for $i = 0, \dots, a$ and $N(a, b, c) = d(d + 1) = d^2 + d$. Moreover, $\min_{k \geq 3} N(d - 1, (d - 1)(k - 1), (k - 3)) = d^2 + d$. This is the origin of the term “ $d^2 + d$ ” in Theorem 1.1.

7.2. Cohomology Results.

Notation 7.8. Let N_d, \mathbb{P}^{N_d} , and $\mathcal{X} \subset \mathbb{P}^{N_d} \times \mathbb{P}^n$ be as in Notation 6.2. Let k be any integer with $3 \leq k \leq \frac{n}{2}$ (only the case $k = 2d$ will be used later). Let $\mathcal{R}^k(\mathbb{P}^n) \subset \text{Hilb}_{\mathbb{P}^n}^{kt+1}$ denote the open subscheme parametrizing curves $C_0 \subset \mathbb{P}^n$ that are projectively equivalent to a degree k rational normal curve $C_0 \subset \mathbb{P}^k \subset \mathbb{P}^n$. Let $\mathcal{R}^k(\mathcal{X}) \subset \mathbb{P}^{N_d} \times \mathcal{R}^k(\mathbb{P}^n)$ denote the parameter space for pairs $([X], [C_0])$ such that $C_0 \subset X$. Let $Q(t) = \frac{1}{2}(t + 1)((2k - 1)t + 2)$ denote the Hilbert polynomial of a rational normal scroll of degree $2k - 1$ in \mathbb{P}^{2k} . Let $\mathcal{U} \subset \text{Hilb}_{\mathbb{P}^n}^{Q(t)}$ denote the open subscheme parametrizing closed subschemes $\Sigma \subset \mathbb{P}^n$ that are projectively equivalent to a rational normal scroll of degree $2k - 1$ in $\mathbb{P}^{2k} \subset \mathbb{P}^n$ and that are abstractly isomorphic to \mathbb{F}_1 . Let $\mathcal{V} \subset \mathcal{U} \times \mathcal{R}^k(\mathbb{P}^n)$ denote the parameter space of pairs $([\Sigma], [C_0])$ such that $C_0 \subset \Sigma$ and such that, via the isomorphism of $\Sigma \cong \mathbb{F}_1$, the invertible sheaf of C_0 is $\mathcal{O}(1, 0)$. Let $\mathcal{W} \subset \mathbb{P}^{N_d} \times \mathcal{U} \times \mathcal{R}^k(\mathbb{P}^n)$ denote the parameter space for triples $([X], [\Sigma], [C_0])$ where $([\Sigma], [C_0])$ is in \mathcal{V} and where $\Sigma \subset X$.

Observe that $\mathcal{R}^k(\mathbb{P}^n)$ is a homogeneous space of PGL_{n+1} , and therefore is smooth and connected. Observe that the projection $\mathcal{R}^k(\mathcal{X}) \rightarrow \mathcal{R}^k(\mathbb{P}^n)$ is a projective bundle of relative dimension $N_d - (kd + 1)$. Observe that the projection map $\mathcal{V} \rightarrow \mathcal{U}$ factors as an open subset (with nonempty fibers) of a projective bundle over \mathcal{U} of relative dimension 2 (more precisely, every fiber is isomorphic to the \mathbb{A}^2 of irreducible curves in the linear system $|\mathcal{O}(1, 0)|$). Observe that the projection map $\mathcal{W} \rightarrow \mathcal{V}$ is a projective bundle of relative dimension $N_d - Q(d)$.

For each triple $([X], [\Sigma], [C_0]) \in \mathcal{W}$, define $\partial_{X, \Sigma} : \mathbb{C}^{n+1} \rightarrow H^0(\Sigma, \mathcal{O}_\Sigma(d-1))$ as in Section 6. More precisely, denote $\mathcal{E} = \mathcal{O}_W^{\oplus(n+1)}$. Denote by \mathcal{G} the unique quotient of $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d-1)) \otimes_{\mathbb{C}} \mathcal{O}_U$ that is locally free and whose fiber at each point Σ is the quotient,

$$H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(d-1)) \rightarrow H^0(\Sigma, \mathcal{O}_\Sigma(d-1)).$$

Denote by \mathcal{F} the locally free \mathcal{O}_W -modules

$$\mathrm{pr}_{\mathbb{P}^{N_d}}^*(\mathcal{O}_{\mathbb{P}^{N_d}}(1)) \otimes \mathrm{pr}_U^* \mathcal{G}.$$

Then there is a map of \mathcal{O}_W -modules $\partial : \mathcal{E} \rightarrow \mathcal{F}$ whose fiber at each point $([X], [\Sigma], [C_0])$ is the map $\partial_{X, \Sigma}$. Denote by $\mathcal{W}^\circ \subset \mathcal{W}$ the open subscheme parametrizing points $([X], [\Sigma], [C_0])$ such that,

$$\mathrm{Image}(\partial_{X, \Sigma}) \subset H^0(\Sigma, \mathcal{O}_{\mathbb{P}^n}(d-3)|_\Sigma),$$

is a $(k-3)$ -generating linear system.

Let $\Sigma \subset \mathbb{P}^{2k} \subset \mathbb{P}^n$ be a rational normal surface scroll of degree $2k-1$, and let $f : \mathbb{F}_1 \rightarrow \Sigma$ be an isomorphism. For each pair of nonnegative integers (a, b) , denote by $N(a, b)$ the locally free $\mathcal{O}_{\mathbb{F}_1}$ -module,

$$N(a, b) = f^*(N_{\Sigma/\mathbb{P}^n} \otimes \mathcal{O}_{\mathbb{P}^n}(-1)|_\Sigma) \otimes \mathcal{O}(a, b),$$

and denote by $N'(a, b) \subset N(a, b)$ the subsheaf,

$$N'(a, b) = f^*(N_{\Sigma/\mathbb{P}^{2k}} \otimes \mathcal{O}_{\mathbb{P}^n}(-1)|_\Sigma) \otimes \mathcal{O}(a, b).$$

Lemma 7.9. (i) $N'(0, 0)$ is generated by global sections and $h^i(\mathbb{F}_1, N'(0, 0)) = 0$ for $i > 0$,
(ii) $N(0, 0)$ is generated by global sections and $h^i(\mathbb{F}_1, N(0, 0)) = 0$ for $i > 0$,
(iii) for every pair of nonnegative integers (a, b) and for every coherent sheaf \mathcal{F} on \mathbb{F}_1 that is generated by global sections and such that $h^i(\mathbb{F}_1, \mathcal{F}) = 0$ for $i > 0$, $\mathcal{F}(a, b) := \mathcal{F} \otimes \mathcal{O}(a, b)$ is generated by global sections and $h^i(\mathbb{F}_1, \mathcal{F}(a, b)) = 0$ for $i > 0$.

In particular, for every pair of nonnegative integers (a, b) , $N(a, b)$ (resp. $N'(a, b)$) is generated by global sections and $h^i(\mathbb{F}_1, N(a, b)) = 0$ for $i > 0$ (resp. $h^i(\mathbb{F}_1, N'(a, b)) = 0$ for $i > 0$).

Proof. (i): The morphism $\mathrm{pr}_{\mathbb{P}^1} : \mathbb{F}_1 \rightarrow \mathbb{P}^1$ is isomorphic over \mathbb{P}^1 to projection from the projective bundle,

$$\mathbb{F}_1 \cong \mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(-(k-1)) \oplus \mathcal{O}_{\mathbb{P}^1}(-k)).$$

Under this isomorphism the invertible sheaf $\mathcal{O}(1, k-1)$ on Σ corresponds to the invertible sheaf $\mathcal{O}(1)$ on $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(-(k-1)) \oplus \mathcal{O}_{\mathbb{P}^1}(-k))$ where $\mathcal{O}(1)$ is the universal invertible quotient of $\mathrm{pr}_{\mathbb{P}^1}^*(\mathcal{O}_{\mathbb{P}^1}(k-1) \oplus \mathcal{O}_{\mathbb{P}^1}(k))$. Up to projective equivalence, the morphism $f : \mathbb{F}_1 \rightarrow \mathbb{P}^{2k}$ is the closed immersion given by the complete linear system of $\mathcal{O}(1)$; in particular, $f^* \mathcal{O}_{\mathbb{P}^{2k}}(1) \cong \mathcal{O}(1)$. Using this isomorphism, there is a short exact sequence of $\mathcal{O}_{\mathbb{F}_1}$ -modules,

$$0 \rightarrow \mathrm{pr}_{\mathbb{P}^1}^* T_{\mathbb{P}^1} \rightarrow \mathrm{pr}_{\mathbb{P}^1}^*(\mathcal{O}_{\mathbb{P}^1}(1)^{\oplus(2k-1)}) \otimes f^* \mathcal{O}_{\mathbb{P}^{2k}}(1) \rightarrow f^* N_{\Sigma/\mathbb{P}^{2k}} \rightarrow 0.$$

Twisting by $f^* \mathcal{O}_{\mathbb{P}^{2k}}(-1)$, $N'(0, 0)$ is a quotient of $\mathrm{pr}_{\mathbb{P}^1}^*(\mathcal{O}_{\mathbb{P}^1}(1)^{\oplus(2k-1)})$. Hence $N'(0, 0)$ is generated by global sections. Also,

$$(\mathrm{pr}_{\mathbb{P}^1})_*(\mathrm{pr}_{\mathbb{P}^1}^* T_{\mathbb{P}^1} \otimes f^* \mathcal{O}_{\mathbb{P}^{2k}}(-1)) = \{0\}, \quad R^1(\mathrm{pr}_{\mathbb{P}^1})_*(\mathrm{pr}_{\mathbb{P}^1}^* T_{\mathbb{P}^1} \otimes f^* \mathcal{O}_{\mathbb{P}^{2k}}(-1)) = \{0\}.$$

Twisting the short exact sequence by $f^*\mathcal{O}_{\mathbb{P}^{2k}}(-1)$ and forming the associated long exact sequence of higher direct images, $R^1(\mathrm{pr}_{\mathbb{P}^1})_*(f^*N_{\Sigma/\mathbb{P}^{2k}}(-1)) = \{0\}$, and $(\mathrm{pr}_{\mathbb{P}^1})_*(f^*N_{\Sigma/\mathbb{P}^{2k}}(-1)) \cong \mathcal{O}_{\mathbb{P}^1}(1)^{\oplus(2k-1)}$. Computing the cohomology of $N'(0,0)$ via the Leray spectral sequence associated to $\mathrm{pr}_{\mathbb{P}^1} : \mathbb{F}_1 \rightarrow \mathbb{P}^1$, $h^i(\mathbb{F}_1, N'(0,0)) = 0$ for $i > 0$.

(ii): There is a short exact sequence,

$$0 \longrightarrow N'(0,0) \longrightarrow N(0,0) \longrightarrow \mathcal{O}(0,0)^{\oplus(n-2k)} \longrightarrow 0.$$

By (i), $h^i(\mathbb{F}_1, \mathcal{O}_{\mathbb{F}_1}) = 0$ for $i > 0$. Therefore $N(0,0)$ is generated by global sections and $h^i(\mathbb{F}_1, N(0,0)) = 0$ for $i > 0$.

(iii): Let \mathcal{F} be a coherent sheaf on \mathbb{F}_1 such that \mathcal{F} is generated by global sections and such that $h^i(\mathbb{F}_1, \mathcal{F}) = 0$ for $i > 0$. It will be proved by double induction on (a, b) that for every pair of nonnegative integers (a, b) , $\mathcal{F}(a, b)$ is generated by global sections and $h^i(\mathbb{F}_1, \mathcal{F}(a, b)) = 0$ for $i > 0$.

The base case is $b = 0$ and is established by induction on a . For $a = 0$, the result follows by hypothesis. Let $a > 0$ and, by way of induction, suppose the result is proved for $a - 1$. Let $D \subset \mathbb{F}_1$ be a general member of the linear system $|\mathcal{O}(1,0)|$. Then D is a smooth curve isomorphic to \mathbb{P}^1 . Since D is general, there is a short exact sequence,

$$0 \longrightarrow \mathcal{F}(a-1,0) \longrightarrow \mathcal{F}(a,0) \longrightarrow \mathcal{F}(a,0)|_D \longrightarrow 0.$$

The sheaf $\mathcal{F}|_D$ is generated by global sections, and $\mathcal{O}_{\mathbb{F}_1}(a(e+f))|_D \cong \mathcal{O}_{\mathbb{P}^1}(a)$. Hence also $\mathcal{F}(a,0)|_D$ is generated by global sections. By the induction assumption, $h^1(\mathbb{F}_1, \mathcal{F}(a-1,0)) = 0$. By the long exact sequence of cohomology associated to the short exact sequence, every global section of $\mathcal{F}(a,0)|_D$ is the image of a global section of $\mathcal{F}(a,0)$. Hence $\mathcal{F}(a,0)$ is generated by global sections. A coherent sheaf on \mathbb{P}^1 that is generated by global sections has no higher cohomology. Combined with the induction assumption and the long exact sequence in cohomology associated to the short exact sequence above, $h^i(\mathbb{F}_1, \mathcal{F}(a,0)) = 0$ for $i > 0$. Therefore, for every $a > 0$, $\mathcal{F}(a,0)$ is generated by global sections and $h^i(\mathbb{F}_1, \mathcal{F}(a,0)) = 0$ for $i > 0$.

Suppose that $b > 0$ and suppose the result is proved for $b - 1$. Let $L \subset \mathbb{F}_1$ be a general fiber of $\mathrm{pr}_{\mathbb{P}^1}$. Then L is smooth and isomorphic to \mathbb{P}^1 . Since L is general, there is a short exact sequence,

$$0 \longrightarrow \mathcal{F}(a,b-1) \longrightarrow \mathcal{F}(a,b) \longrightarrow \mathcal{F}(a,b)|_L \longrightarrow 0.$$

Via the isomorphism $L \cong \mathbb{P}^1$, $\mathcal{O}_{\mathbb{F}_1}(a(e+f) + bf)|_L \cong \mathcal{O}_{\mathbb{P}^1}(a)$. By almost identical arguments to those above, $\mathcal{F}(a,b)$ is generated by global sections and $h^i(\mathbb{F}_1, \mathcal{F}(a,b)) = 0$ for $i > 0$. \square

Let $([X], [\Sigma], [C_0])$ be a point in \mathcal{W}^o and let $f : \mathbb{F}_1 \rightarrow \Sigma$ be an isomorphism. For each pair of nonnegative integers (a, b) , denote $N_X(a, b) = f^*(N_{\Sigma/X} \otimes \mathcal{O}_{\mathbb{P}^n}(-1)|_{\Sigma}) \otimes \mathcal{O}(a, b)$.

Lemma 7.10. (i) *The hypersurface X is smooth along Σ .*

- (ii) *For each pair of nonnegative integers (a, b) , $h^i(\mathbb{F}_1, N_X(a, b + k - 3)) = 0$ for $i > 0$.*
- (iii) *For every line of ruling $L \subset \Sigma$ and every nonnegative integer a , $h^1(L, N_{L/X}(a - 1)) = 0$.*
- (iv) *For every nonnegative integer a , $h^1(C_0, N_{C_0/X}(a - 2)) = 0$.*
- (v) *The projection morphism $\mathrm{pr}_{\mathbb{P}^{N_d}} : \mathcal{W} \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [\Sigma], [C_0])$.*
- (vi) *For every line of ruling $L \subset \Sigma$, the projection morphism $\mathrm{pr}_{\mathbb{P}^{N_d}} : F(\mathcal{X}) \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [L])$.*
- (vii) *The projection morphism $\mathrm{pr}_{\mathbb{P}^{N_d}} : \mathcal{R}^k(\mathcal{X}) \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [C_0])$.*
- (viii) *The projection morphism $\pi : \mathcal{W} \rightarrow \mathcal{R}^k(\mathcal{X})$ is smooth at $([X], [\Sigma], [C_0])$.*

Proof. (i): Since the partial derivatives of a defining equation of X span a c -generating linear series, in particular they generate the sheaf $\mathcal{O}_\Sigma(d-1)$. Hence, there is no point of Σ at which all the partial derivatives vanish. By the Jacobian criterion, X is smooth along Σ .

(ii): There is a short exact sequence,

$$0 \longrightarrow N_{\Sigma/X} \longrightarrow N_{\Sigma/\mathbb{P}^n} \longrightarrow N_{X/\mathbb{P}^n}|_\Sigma \longrightarrow 0.$$

Denote $\alpha = a + (d-1)$ and $\beta = b + (d-1)(k-1)$ (these are different than $\alpha_d, \alpha_r, \beta_d$ and β_r). There is a short exact sequence,

$$0 \longrightarrow N_X(a, b) \longrightarrow N(a, b) \longrightarrow \mathcal{O}(\alpha, \beta) \longrightarrow 0.$$

Since $a, b \geq 0$, by Lemma 7.9, $h^i(\mathbb{F}_1, N(a, b)) = 0$ for $i \geq 0$. By direct computation, $h^i(\mathbb{F}_1, \mathcal{O}(\alpha, \beta)) = 0$ for $i \geq 0$. Hence $h^2(\mathbb{F}_1, N_X(a, b)) = 0$, and $h^1(\mathbb{F}_1, N_X(a, b)) = 0$ iff the following map is surjective,

$$H^0(\mathbb{F}_1, N(a, b)) \rightarrow H^0(\mathbb{F}_1, \mathcal{O}(\alpha, \beta)).$$

There is a commutative diagram,

$$\begin{array}{ccc} H^0(\mathbb{F}_1, N(a, b)) \otimes H^0(\mathbb{F}_1, \mathcal{O}(a', b')) & \longrightarrow & H^0(\mathbb{F}_1, N(a+a', b+b')) \\ \downarrow & & \downarrow \\ H^0(\mathbb{F}_1, \mathcal{O}(\alpha, \beta)) \otimes H^0(\mathbb{F}_1, \mathcal{O}(a', b')) & \longrightarrow & H^0(\mathbb{F}_1, \mathcal{O}(\alpha+a', \beta+b')) \end{array}.$$

By direct computation the bottom horizontal arrow is surjective if $a', b' \geq 0$. Hence, if the left vertical arrow is surjective, then also the right vertical arrow is surjective; i.e., if $h^1(\mathbb{F}_1, N_X(a, b)) = 0$ then also $h^1(\mathbb{F}_1, N_X(a+a', b+b')) = 0$. Thus (ii), is reduced to the case $a = 0, b = k-3$. In this case the commutative diagram above factors the following commutative diagram,

$$\begin{array}{ccc} H^0(\mathbb{F}_1, T_{\mathbb{P}^n}) \otimes H^0(\mathbb{F}_1, \mathcal{O}(0, k-3)) & \longrightarrow & H^0(\mathbb{F}_1, N(0, k-3)) \\ \downarrow & & \downarrow \\ H^0(\mathbb{F}_1, \mathcal{O}(d-1, (d-1)(k-1))) \otimes H^0(\mathbb{F}_1, \mathcal{O}(0, k-3)) & \longrightarrow & H^0(\mathbb{F}_1, \mathcal{O}(\alpha, \beta)) \end{array}.$$

By definition, the composition,

$$H^0(\mathbb{F}_1 T_{\mathbb{P}^n}) \otimes H^0(\mathbb{F}_1, \mathcal{O}(0, k-3)) \rightarrow H^0(\mathbb{F}_1, \mathcal{O}(\alpha, \beta)),$$

is surjective iff the triple $([X], [\Sigma], [C_0])$ is in \mathcal{W}^o . Since $([X], [\Sigma], [C_0])$ is in \mathcal{W}^o , the right vertical arrow is surjective, i.e., $h^1(\mathbb{F}_1, N_X(0, k-3)) = 0$.

(iii): There is a short exact sequence,

$$0 \longrightarrow N_{L/\Sigma}(a-1) \longrightarrow N_{L/X}(a-1) \longrightarrow N_{\Sigma/X}|_L(a-1) \longrightarrow 0.$$

Since $N_{L/\Sigma} \cong \mathcal{O}_L$, for all $a \geq 0$, $h^1(L, N_{L/\Sigma}(a-1)) = 0$. Therefore it suffices to prove $h^1(L, N_{\Sigma/X}|_L(a-1)) = 0$. Since $\mathcal{O}(a-1, b)|_L \cong \mathcal{O}_L(a-1)$, there is a short exact sequence,

$$0 \longrightarrow N_X(a, k-3) \longrightarrow N_X(a, k-2) \longrightarrow N_{\Sigma/X}|_L(a-1) \longrightarrow 0.$$

By (ii), for $a \geq 0$ the higher cohomology of the first two terms vanishes. By the long exact sequence in cohomology associated to this short exact sequence, $h^1(L, N_{\Sigma/X}|_L(a-1)) = 0$ for $a \geq 0$.

(iv): The proof is almost identical to the proof of (iii).

(v): By [16, Prop. 2.14.2], the obstruction space for the relative Hilbert scheme $\text{Hilb}_{\mathcal{X}/\mathbb{P}^n}^{Q(t)}$ at the point $([X], [\Sigma])$ is contained in $H^1(\Sigma, N_{\Sigma/X})$. If $([X], [\Sigma], [C_0])$ is in \mathcal{W}^o , then by (ii), $h^1(\Sigma, N_{\Sigma/X}) =$

$h^1(\mathbb{F}_1, N_X(1, k-1)) = 0$. By [16, Thm. 2.10], $\text{Hilb}_{\mathcal{X}/\mathbb{P}^{N_d}}^{Q(t)} \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [\Sigma])$. The projection $\mathcal{W}^o \rightarrow \text{Hilb}^{Q(t)}(\mathcal{X}/\mathbb{P}^{N_d})$ is an open subset of a projective bundle, and so is smooth. Therefore the composite morphism $\mathcal{W}^o \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [\Sigma], [C_0])$.

(vi): The proof is very similar to the proof of (v) and uses the vanishing, $h^1(L, N_{L/X}) = 0$, which was proved in (iii).

(vii): The proof is very similar to the proof of (v) and uses the vanishing, $h^1(C_0, N_{C_0/X}) = 0$, which was proved in (iv).

(viii): Since $\mathcal{W}^o \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [\Sigma], [C_0])$ and since $\mathcal{R}^k \rightarrow \mathbb{P}^{N_d}$ is smooth at $([X], [C_0])$, to prove that $\pi : \mathcal{W}^o \rightarrow \mathcal{R}^k(\mathcal{X})$ is smooth at $([X], [\Sigma], [C_0])$, it suffices to check that the derivative map $d\pi : T_{\mathcal{W}^o/\mathbb{P}^{N_d}} \rightarrow \pi^*T_{\mathcal{R}^k(\mathcal{X})/\mathbb{P}^{N_d}}$ is surjective at $([X], [\Sigma], [C_0])$. This reduces to the statement that $H^0(\Sigma, N_{\Sigma/X}) \rightarrow H^0(C_0, N_{\Sigma/X}|_{C_0})$ is surjective. The cokernel is contained in $H^1(\mathbb{F}_1, N_X(0, k-1))$. By (iii), $h^1(C_0, N_{\Sigma/X}|_{C_0}) = 0$, therefore the derivative $d\pi$ is surjective at $([X], [\Sigma], [C_0])$. \square

Let $([X], [\Sigma], [C_0])$ be a point of \mathcal{W} . Denote by $\sigma : C_0 \rightarrow \Sigma$ the inclusion and denote by $\text{pr}_{C_0} : \Sigma \rightarrow C_0$ the unique projection morphism such that σ is a section of pr_{C_0} (via the isomorphism $\Sigma \cong \mathbb{F}_1$, pr_{C_0} corresponds to $\text{pr}_{\mathbb{P}^1}$). Denote by $g : \Sigma \rightarrow X$ the inclusion. There is a family of stable maps $\zeta : C_0 \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$,

$$\zeta = ((\text{pr}_{C_0} : \Sigma \rightarrow C_0, \sigma), g : \Sigma \rightarrow X).$$

Lemma 7.11. *If $([X], [\Sigma], [C_0])$ is in \mathcal{W}^o , then $\zeta : C_0 \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ is very twisting and very positive.*

Proof. Very twisting: First the axioms of Definition 4.3 are verified. Since $g \circ \sigma : C_0 \rightarrow X$ is an embedding, Axiom (i) of Definition 4.3 is satisfied. By Lemma 7.10 (vii), the dimension of the obstruction group of $\overline{\mathcal{M}}_{0,0}(X, k)$ at $[g \circ \sigma : C_0 \rightarrow X]$ is 0, i.e., Axiom (ii) is satisfied.

The proof that Axiom (iii) holds is identical to the argument for Axiom (iii) in the proof of Lemma 6.4, with Lemma 6.3 (iv) replaced by Lemma 7.10 (iii).

As in the proof of Lemma 6.4, $\zeta^*T_{\text{ev}} \cong (\text{pr}_{C_0})_*N_X(0, k-1)$. Hence ζ^*T_{ev} is ample iff $h^1(C_0, \zeta^*T_{\text{ev}}(-2)) = 0$. By a Leray spectral sequence argument similar to the one in the proof of Lemma 6.4, $h^1(C_0, \zeta^*T_{\text{ev}}(-2)) = h^1(\mathbb{F}_1, N_X(0, k-3))$, which, Lemma 7.10 (ii), equals 0. Therefore ζ^*T_{ev} is an ample bundle, i.e., Axiom (iv) is satisfied.

Finally, observe that $\sigma^*\mathcal{O}_\Sigma(\sigma) \cong \mathcal{O}_{C_0}(1)$ is ample, i.e., Axiom (v) is satisfied. Thus ζ is a very twisting family.

Very positive: Next the axioms of Definition 4.11 are verified. Axioms (i), (ii) and (iii) follow from Axioms (i), (ii) and (iii) of Definition 4.3, as proved above. There is a short exact sequence,

$$0 \longrightarrow \zeta^*T_{\text{ev}} \longrightarrow \zeta^*T_{\overline{\mathcal{M}}_{0,1}(X,1)} \longrightarrow (g \circ \sigma)^*T_X \longrightarrow 0.$$

It is proved above that ζ^*T_{ev} is ample. Moreover, by Lemma 7.10 (iv), $N_{C_0/X}$ is ample. Of course T_{C_0} is ample. Therefore $T_X|_{C_0}$ is ample by Lemma 2.10 (ii). Since the first and last term in the short exact sequence are ample, by Lemma 2.10 (ii), $\zeta^*T_{\overline{\mathcal{M}}_{0,1}(X,1)}$ is ample. Since $\zeta^*\text{pr}^*T_{\overline{\mathcal{M}}_{0,0}(X,1)}$ is a quotient of $\zeta^*T_{\overline{\mathcal{M}}_{0,1}(X,1)}$, by Lemma 2.10 (i), $\zeta^*\text{pr}^*T_{\overline{\mathcal{M}}_{0,0}(X,1)}$ is ample; i.e., Axiom (iv) is satisfied.

Finally, $\sigma^* \mathcal{O}_\Sigma(\sigma) \cong \mathcal{O}_{C_0}(1)$, which is ample, i.e., Axiom (v) is satisfied. Thus ζ is a very positive family. \square

Proposition 7.12. (i) *If $d \geq 3$ and $n \geq d^2 + d + 1$, then for $k = 2d$, $\mathcal{W}^\circ \rightarrow \mathcal{R}^k(\mathcal{X})$ is dominant, and $\mathcal{R}^k(\mathcal{X}) \rightarrow \mathbb{P}^{N_d}$ is dominant.*
(ii) *If $d = 1$ or 2 and if $n \geq 7$, then for $k = 3$, $\mathcal{W}^\circ \rightarrow \mathcal{R}^3(\mathcal{X})$ is dominant, and $\mathcal{R}^3(\mathcal{X}) \rightarrow \mathbb{P}^{N_d}$ is dominant.*

Proof. By item (8) of lemma 7.10, it suffices to prove that \mathcal{W}° is nonempty. We have to find a pair $([X], [\Sigma])$ such that for $a = d - 1$, $b = (d - 1)(k - 1)$ and for $c = k - 3$, we have that the image of the derivative map

$$d_{X, \Sigma} : H^0(\mathbb{P}^n, T_{\mathbb{P}^n}(-1)) \rightarrow H^0(\mathbb{F}_1, \mathcal{O}(a, b)) \quad (13)$$

is a c -generating linear system.

Recall that $S = \mathbb{C}[T_0, T_1, U, V]$ is the \mathbb{Z}^2 -graded Cox homogeneous coordinate ring of \mathbb{F}_1 . Denote by A_d the set of $d^2 + d$ monomials that occur in the linear system $W_0(a, b, c)$, i.e.,

$$A_d = \left\{ U^i V^{d-1-i} T_0^{((d-1)(k-1)+i)-j(k-2)} T_1^{j(k-2)} \mid i = 0, \dots, d-1, j = 1, \dots, r(i) \right\} \\ \cup \left\{ U^i V^{d-1-i} T_1^{(d-1)(k-1)+i} \mid i = 0, \dots, d-1 \right\}$$

where $r(i) = d - 1 + \left\lfloor \frac{d-2+i}{k-2} \right\rfloor$.

(i), $d \geq 4$: Suppose that $d \geq 4$ and $n \geq d^2 + d + 1$. Denote by B_d the set of $4d - 1$ monomials,

$$B_d = \begin{array}{l} \{ U^{d-1} \quad T_0^{(d-1)k-(k-2)j} \quad T_1^{(k-2)j} \mid j = 0, \dots, d-1 \} \\ \cup \{ U^{d-2} \quad V \quad T_0^{(d-1)k-1-(k-2)j} \quad T_1^{(k-2)j} \mid j = 0, \dots, d-1 \} \\ \cup \{ U^{d-3} \quad V^2 \quad T_0^{(d-1)k-2-(k-2)j} \quad T_1^{(k-2)j} \mid j = 1, \dots, d-1 \} \\ \cup \{ U^{d-4} \quad V^3 \quad T_0^{(d-1)k-3-(k-2)j} \quad T_1^{(k-2)j} \mid j = 1, \dots, d-1 \} \\ \cup \{ \quad \quad \quad V^{d-1} \quad T_0^{(d-1)(k-1)} \end{array}$$

Denote by C_d the set of $d^2 - 3d + 1$ monomials $C_d = A_d - B_d$. Denote by

$$\{Y_0, \dots, Y_k\} \cup \{Z_0, \dots, Z_{k-1}\} \cup \{X_M \mid M \in C_d\} \cup \{V_l \mid l = 1, \dots, n - (d^2 + d + 1)\},$$

a basis of $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$, i.e., a basis of homogeneous coordinates on \mathbb{P}^n .

Denote by $f : \mathbb{F}_1 \rightarrow \mathbb{P}^{2k} \subset \mathbb{P}^n$ the morphism mapping $([T_0 : T_1], [T_0 U : T_1 U : V]) \in \mathbb{F}_1$ to the point in \mathbb{P}^n with coordinates $X_M = 0$, $M \in C_d$, with $V_l = 0$, $l = 1, \dots, n - (d^2 + d + 1)$, and with

$$Y_0 = U T_0^k, \dots, Y_i = U T_0^{k-i} T_1^i, \dots, Y_k = U T_1^k, \\ Z_0 = V T_0^{k-1}, \dots, Z_j = V T_0^{k-1-j} T_1^j, \dots, Z_{k-1} = V T_1^{k-1}.$$

This is an embedding whose image $\Sigma = f(\mathbb{F}_1)$ is a rational normal scroll of degree $2k - 1$.

The pullback map $H^0(\mathbb{P}^{2k}, \mathcal{O}_{\mathbb{P}^{2k}}(1)) \rightarrow H^0(\mathbb{F}_1, \mathcal{O}(1, k - 1))$ is surjective by construction. And the natural map,

$$\text{Sym}^{d-1} H^0(\mathbb{F}_1, \mathcal{O}(1, k - 1)) \rightarrow H^0(\mathbb{F}_1, \mathcal{O}(d - 1, (d - 1)(k - 1))),$$

is surjective. Therefore the pullback map

$$H^0(\mathbb{P}^{2k}, \mathcal{O}_{\mathbb{P}^{2k}}(d - 1)) \rightarrow H^0(\mathbb{F}_1, \mathcal{O}(d - 1, (d - 1)(k - 1)))$$

is surjective. For each monomial $M \in C_d$, choose a polynomial $G_M(Y_0, \dots, Y_{2k})$ such that $f^* G_M = M$.

coordinate	$\text{IN}(\partial_{X,\Sigma}(\text{coordinate}))$	lower order terms in $\partial_{X,\Sigma}(\text{coordinate})$
$Z_{j+1},$ $j = 0, \dots, d-1$	$U^{d-1}T_0^{(k-1)d-(k-2)j}T_1^{(k-2)j}$	$-U^{d-1}T_0^{(k-1)d-(k-2)(j+1)-1}T_1^{(k-2)(j+1)+1}$
Z_d	$U^{d-1}T_0^{(k-1)d-(k-2)(d-1)}T_1^{(k-2)(d-1)}$	$-U^{d-3}V^2T_0^{(k-1)(d-2)+(d-3)}T_1^{k-1}$ $-UV^{d-2}T_0^{(k-1)(d-2)+d}T_1^d$
$Y_{j+1},$ $j = 0, \dots, d-1$	$-U^{d-2}VT_0^{(k-1)d-1-(k-2)j}T_1^{(k-2)j}$	$+U^{d-2}VT_0^{(k-1)d-(k-2)(j+1)-2}T_1^{(k-2)(j+1)+1}$
Y_d	$-U^{d-2}VT_0^{(k-1)d-1-(k-2)(d-1)}T_1^{(k-2)(d-1)}$	$+U^{d-4}V^2T_0^{(k-1)(d-2)+(d-4)}T_1^{k-1}$ $-V^{d-1}T_0^{(k-1)(d-1)-d}T_1^d$
$Z_{d+j},$ $j = 1, \dots, d-2$	$U^{d-3}V^2 \times$ $T_0^{(k-1)(d-1)+(d-3)-(k-2)j}T_1^{(k-2)j}$	$-U^{d-2}V^2 \times$ $T_0^{(k-1)(d-1)+(d-4)+(k-2)(j+1)}T_1^{(k-2)(j+1)}$
Z_{k-1}	$U^{d-3}V^2 \times$ $T_0^{(k-1)(d-1)+(d-3)-(k-2)(d-1)}T_1^{(k-2)(d-1)}$	
$Y_{d+j},$ $j = 1, \dots, d-2$	$-U^{d-4}V^3 \times$ $T_0^{(k-1)(d-1)+(d-4)-(k-2)j}T_1^{(k-2)j}$	$U^{d-4}V^3 \times$ $T_0^{(k-1)(d-1)+(d-5)+(k-2)(j+1)}T_1^{(k-2)(j+1)}$
Y_{k-1}	$U^{d-4}V^3 \times$ $T_0^{(k-1)(d-1)+(d-4)-(k-2)(d-1)}T_1^{(k-2)(d-1)}$	
Y_k	$V^{d-1}T_0^{(k-1)(d-1)}$	
$X_M,$ $M \in C_d$	M	
Z_0	$-U^{d-1}T_0^{(k-1)(d-1)+(d-2)}T_1$	$UV^{d-2}T_0^{(k-1)(d-2)}T_1^k$
Y_0	$U^{d-2}VT_0^{(k-1)(d-1)+(d-3)}T_1$	

FIGURE 1. The map $\partial_{X,\Sigma}$

Consider the hypersurface $X \subset \mathbb{P}^n$ with defining equation

$$\begin{aligned}
F = & \sum_{j=0}^{d-2} (Y_j Z_{j+1} - Y_{j+1} Z_j) & Y_0^{d-2-j} Y_{k-3}^i & + \\
& \sum_{j=2}^{d-2} (Y_{d-1+j} Z_{d+j} - Y_{d+j} Z_{d-1+j}) & Y_k^{j-2} Y_3^{d-2-l} Z_3 Z_4 & + \\
& (Y_{d-1} Z_d - Y_d Z_{d-1}) & Y_{k-3}^{d-4} Y_{k-4}^2 & + \\
& (Y_d Z_{d+1} - Y_{d+1} Z_d) & Y_1^{d-4} Z_7 Z_8 & + \\
& (Y_{k-2} Z_{k-1} - Y_{k-1} Z_{k-2}) & Y_k^{d-4} Z_{k-1} Z_5 & + \\
& (Y_k Z_0 - Z_d Y_d) & Z_0^{d-2} & + \\
& \sum_{M \in C_d} G_M(Y_0, \dots, Y_{2k}) X_M.
\end{aligned}$$

Observe that F is contained in the homogeneous ideal of Σ , i.e., $\Sigma \subset X$. The derivative map $\partial_{X,\Sigma}$ acts on the coordinates Y_j, Z_j, X_M as in Figure 1. Each of the monomials in A_d occurs as the initial term of $\partial_{X,\Sigma}$ acting on one coordinate. For every coordinate except Y_0 and Z_0 , the initial term of $\partial_{X,\Sigma}$ is one of the monomials in A_d . By Lemma 7.6, the image of $\partial_{X,\Sigma}$ is a $(k-3)$ -generating linear system.

(i), $d = 3$: Suppose that $d = 3$ and $n \geq 3^2 + 3 + 1 = 13$. Denote by B_3 the set of $4d - 1 = 11$ monomials,

$$B_d = \begin{array}{l} \{ U^2 \quad T_0^{12-(k-2)j} \quad T_1^{(k-2)j} \quad |j = 0, 1, 2 \} \\ \cup \{ U \quad V \quad T_0^{11-(k-2)j} \quad T_1^{(k-2)j} \quad |j = 0, 1, 2 \} \\ \cup \{ \quad \quad V^2 \quad T_0^{10-(k-2)j} \quad T_1^{(k-2)j} \quad |j = 0, 1, 2 \} \\ \cup \{ U \quad V \quad \quad \quad T_1^{11} \quad \quad \quad \} \\ \cup \{ \quad \quad V^2 \quad \quad \quad T_1^{10} \quad \quad \quad \}. \end{array}$$

Denote by C_3 the singleton set consisting of the monomial $M = U^2 T_1^{12}$. Denote by

$$\{Y_0, \dots, Y_6\} \cup \{Z_0, \dots, Z_5\} \cup \{X_M | M \in C_3\} \cup \{V_l | l = 1, \dots, n - 13\},$$

a basis of $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$, i.e., a basis of homogeneous coordinates on \mathbb{P}^n .

Denote by $f : \mathbb{F}_1 \rightarrow \mathbb{P}^{12} \subset \mathbb{P}^n$ the morphism mapping $([T_0 : T_1], [T_0 U : T_1 U : V]) \in \mathbb{F}_1$ to the point in \mathbb{P}^n with coordinates $X_M = 0, M \in C_d$, with $V_l = 0, l = 1, \dots, n - 13$, and with

$$\begin{aligned} Y_0 &= U T_0^6, \dots, Y_i = U T_0^{6-i} T_1^i, \dots, Y_k = U T_1^6, \\ Z_0 &= V T_0^5, \dots, Z_j = V T_0^{5-j} T_1^j, \dots, Z_5 = V T_1^5. \end{aligned}$$

This is an embedding whose image $\Sigma = f(\mathbb{F}_1)$ is a rational normal scroll of degree 11.

Consider the hypersurface $X \subset \mathbb{P}^n$ with defining equation

$$F = \begin{array}{llll} \sum_{j=0}^1 & (Y_j Z_{j+1} - Y_{j+1} Z_j) & Y_0^{1-j} Y_3^j & + \\ & (Y_2 Z_3 - Y_3 Z_2) & Y_6 & + \\ & (Y_5 Z_5 - Y_6 Z_4) & Z_5 & + \\ & (Y_3 Z_3 - Y_6 Z_0) & Z_4 & + \\ & (Y_2 Z_2 - Y_4 Z_0) & Z_0 & + \\ & Y_6^2 X_M. & & \end{array}$$

Observe that F is contained in the homogeneous ideal of Σ , i.e., $\Sigma \subset X$. It is straightforward to compute the action of $\partial_{X, \Sigma}$ on the coordinates Y_j, Z_j, X_M . Every monomial in A_3 occurs as the initial term of $\partial_{X, \Sigma}$ acting on one coordinate. For every coordinate except Y_0 and Z_0 , the initial term of $\partial_{X, \Sigma}$ is one of the monomials in A_3 . By Lemma 7.6, the image of $\partial_{X, \Sigma}$ is a $(k-3)$ -generating linear system.

(ii), $d = 2$: Suppose that $d = 2$ and $n \geq 2^2 + 2 + 1 = 7$. Let $k = 3$. Denote by B_2 the set of 6 monomials,

$$B_2 = \begin{array}{l} \{ U \quad T_0^{3-j} \quad T_1^j \quad |j = 0, 1, 2 \} \\ \cup \{ \quad \quad V \quad T_0^{2-j} \quad T_1^j \quad |j = 0, 1, 2 \}. \end{array}$$

Denote by C_2 the singleton set consisting of the monomial $M = U T_1^3$. Denote by

$$\{Y_0, Y_1, Y_2, Y_3, Z_0, Z_1, Z_2, X_M\} \cup \{V_l | l = 1, \dots, n - 7\},$$

a basis of $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$, i.e., a basis of homogeneous coordinates on \mathbb{P}^n .

Denote by $f : \mathbb{F}_1 \rightarrow \mathbb{P}^6 \subset \mathbb{P}^n$ the morphism mapping $([T_0 : T_1], [T_0 U : T_1 U : V]) \in \mathbb{F}_1$ to the point in \mathbb{P}^n with coordinates $X_M = 0, M \in C_d$, with $V_l = 0, l = 1, \dots, n - 7$, and with

$$\begin{aligned} Y_0 &= U T_0^3, Y_1 = U T_0^2 T_1, Y_2 = U T_0 T_1^2, Y_3 = U T_1^3, \\ Z_0 &= V T_0^2, Z_1 = V T_0 T_1, Z_2 = V T_1^2. \end{aligned}$$

This is an embedding whose image $\Sigma = f(\mathbb{F}_1)$ is a rational normal scroll of degree 6.

Consider the hypersurface $X \subset \mathbb{P}^n$ with defining equation

$$F = (Y_3Z_1 - Y_2Z_2) + (Y_0Z_1 - Y_1Z_0) + Y_3X_M.$$

Observe that F is contained in the homogeneous ideal of Σ , i.e., $\Sigma \subset X$. It is straightforward to compute the action of $\partial_{X,\Sigma}$ on the coordinates Y_j, Z_j, X_M . Every monomial in A_2 occurs as the initial term of $\partial_{X,\Sigma}$ acting on one coordinate. For every coordinate except Y_0 , the initial term of $\partial_{X,\Sigma}$ is one of the monomials in A_2 . By Lemma 7.6, the image of $\partial_{X,\Sigma}$ is a $(k-3)$ -generating linear system.

(ii), $d = 1$. Denote by

$$\{Y_0, Y_1, Y_2, Y_3, Z_0, Z_1, Z_2, X_M\} \cup \{V_l | l = 1, \dots, n-7\},$$

a basis of $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(1))$, i.e., a basis of homogeneous coordinates on \mathbb{P}^n .

Denote by $f : \mathbb{F}_1 \rightarrow \mathbb{P}^6 \subset \mathbb{P}^n$ the morphism mapping $([T_0 : T_1], [T_0U : T_1U : V]) \in \mathbb{F}_1$ to the point in \mathbb{P}^n with coordinates $X_M = 0, M \in C_d$, with $V_l = 0, l = 1, \dots, n-7$, and with

$$\begin{aligned} Y_0 &= UT_0^3, Y_1 = UT_0^2T_1, Y_2 = UT_0T_1^2, Y_3 = UT_1^3, \\ Z_0 &= VT_0^2, Z_1 = VT_0T_1, Z_2 = VT_1^2. \end{aligned}$$

This is an embedding whose image $\Sigma = f(\mathbb{F}_1)$ is a rational normal scroll of degree 6.

Consider the hypersurface $X \subset \mathbb{P}^n$ with defining equation $F = X_M$. Observe that F is contained in the homogeneous ideal of Σ , i.e., $\Sigma \subset X$. The image of $\partial_{X,\Sigma}(X_M)$ is a generator for $S_{0,0}$; i.e., the image of $\partial_{X,\Sigma}$ is a $(k-3)$ -generating linear system. \square

Together with Remark 7.1, Lemma 7.11 and Proposition 7.12 imply the following corollary.

Corollary 7.13. *If $X \subset \mathbb{P}^n$ is a general hypersurface of degree d and if $n \geq d^2 + d + 1$, then there exists a 1-morphism $\zeta : C_0 \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$ that is both very twisting and very positive.*

8. PROOF OF THE MAIN THEOREM

As explained at the end of Section 1, if $d < \frac{n+1}{2}$, then for a general hypersurface $X \subset \mathbb{P}^n$ of degree d , Hypothesis 1.5, Hypothesis 1.6, and Hypothesis 1.7 are satisfied. By Corollary 6.6, if $d \geq 2$ and $n+1 \geq d^2$, or if $d = 1$ and $n \geq 3$, then for a general hypersurface $X \subset \mathbb{P}^n$ of degree d , Hypothesis 4.10 is satisfied. Finally, if $n \geq d^2 + d + 1$ then by Corollary 7.13 there exists a very twisting, very positive family $\zeta : C_0 \rightarrow \overline{\mathcal{M}}_{0,1}(X, 1)$. Therefore (ζ, ζ) is an inducting pair.

By Theorem 5.13, for every $e \geq 1$ there exists an inducting pair $(\zeta_1, \overline{\zeta}_e)$. In particular, there exists a very positive 1-morphism $\overline{\zeta}_e : C \rightarrow \overline{\mathcal{M}}_{0,1}(X, e)$. As shown in the proof of Theorem 5.13, it may be assumed that C is smooth and that the image of C is contained in the smooth locus of the fine moduli locus. By Lemma 4.14 (i), $\overline{\zeta}_e$ is a very free morphism. By [11, Prop. 7.4], $\overline{\mathcal{M}}_{0,0}(X, e)$ is an irreducible variety. Therefore, by [16, Thm. IV.3.7], $\overline{\mathcal{M}}_{0,0}(X, e)$ is rationally connected.

REFERENCES

- [1] D. Abramovich and A. Vistoli. Compactifying the space of stable maps. *Journal of the American Mathematical Society*, 15:27–75, 2002. arXiv: math.AG/9908167.
- [2] V. Alexeev. Moduli spaces $m_{g,n}(w)$ for surfaces. In *Higher dimensional complex varieties (Trento, 1994)*, pages 1–22. de Gruyter, 1996.
- [3] K. Behrend. Gromov-Witten invariants in algebraic geometry. *Inventiones Mathematica*, 127:601–617, 1997. arXiv: math.AG/9601011.

- [4] K. Behrend and B. Fantechi. The intrinsic normal cone. *Inventiones Mathematica*, 128:45–88, 1998. arXiv: math.AG/961010.
- [5] K. Behrend and Y. Manin. Stacks of stable maps and Gromov-Witten invariants. *Duke Math Journal*, 85:1–60, 1996. arXiv: math.AG/9506023.
- [6] S. Bosch, W. Lütkebohmert, and M. Raynaud. *Néron Models*, volume 21 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, 1990.
- [7] O. Debarre. *Higher-dimensional algebraic geometry*. Universitext. Springer-Verlag, 2001.
- [8] W. Fulton. *Intersection Theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge*. Springer-Verlag, 2 edition, 1998.
- [9] W. Fulton and R. Pandharipande. Notes on stable maps and quantum cohomology. In *Algebraic geometry – Santa Cruz 1995*, pages 45–96. American Mathematical Society, 1995. arXiv: math.AG/9608011.
- [10] A. Grothendieck and J. Dieudonné. *Éléments de géométrie algébrique. IV*, volume 20, 24, 28, 32 of *Publications Mathématiques*. Institute des Hautes Études Scientifiques., 1964-1967.
- [11] J. Harris, M. Roth, and J. Starr. Rational curves on hypersurfaces of low degree. submitted Crelle’s Journal, preprint arXiv: math.AG/0203088.
- [12] R. Hartshorne. *Residues and Duality, Lecture notes of a seminar on the work of A. Grothendieck*, volume 20 of *Lecture Notes in Math*. Springer-Verlag, 1966.
- [13] R. Hartshorne. *Algebraic Geometry*, volume 52 of *Graduate Texts in Mathematics*. Springer-Verlag, 1977.
- [14] J.-P. Jouanolou. *Théorèmes de Bertini et Applications*, volume 42 of *Progress in Mathematics*. Birkhäuser, 1983.
- [15] B. Kim and R. Pandharipande. The connectedness of the moduli space of maps to homogeneous spaces. In *Symplectic geometry and mirror symmetry*, chapter 5. World Scientific, 2001. arXiv: math.AG/0003168.
- [16] J. Kollár. *Rational Curves on Algebraic Varieties*, volume 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge*. Springer-Verlag, 1996.
- [17] J. Kollár and S. Mori. *Birational Geometry of Algebraic Varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, 1998.
- [18] S. Lang. On quasi-algebraic closure. *Ann. Math.*, 55:373–390, 1952.
- [19] R. Pandharipande. Intersections of \mathbb{Q} -divisors on Kontsevich’s moduli space. *Trans. Amer. Math. Soc.*, 351:1481–1505, 1999.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE MA 02138
E-mail address: `harris@math.harvard.edu`

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE MA 02139
E-mail address: `jstarr@math.mit.edu`