

MAT 545 FALL 2025
HOMEWORK 2

1. (i) Let X be a smooth (or complex) manifold with the open covering $\{U_\alpha\}_{\alpha \in A}$ and the assignment for each nonempty ordered intersection $U_\alpha \cap U_\beta$ a smooth (holomorphic) map $g_{\alpha\beta} : U_\alpha \cap U_\beta \rightarrow \text{GL}(r, \mathbb{R})$ (or $\text{GL}(r, \mathbb{C})$), satisfying

$$g_{\alpha\beta} \cdot g_{\beta\gamma} \cdot g_{\gamma\alpha} = I \quad \text{on} \quad U_\alpha \cap U_\beta \cap U_\gamma \neq \emptyset, \quad \text{and} \quad g_{\alpha\alpha} = I,$$

where I is the identity operator on $\mathbb{R}^r$ (or $\mathbb{C}^r$). Let E be smooth (or holomorphic) rank r real (or complex) vector bundle over X , defined by

$$E = \bigsqcup_{\alpha \in A} U_\alpha \times \mathbb{K}^r / \sim \quad (\mathbb{K} = \mathbb{R} \text{ or } \mathbb{C}),$$

where $(x, u) \in U_\alpha \times \mathbb{K}^r$ is equivalent to $(y, v) \in U_\beta \times \mathbb{K}^r$ iff $x = y$ and $u = g_{\alpha\beta}(x)v$. Prove that transition functions for E are $g_{\alpha\beta}$.

- (ii) Let X be connected complex manifold and $\mathcal{E}$ be a locally free sheaf of $\mathcal{O}_X$ -modules over X . Prove that there is a holomorphic vector bundle over X whose sheaf of holomorphic sections is $\mathcal{E}$.

2. Let X be a complex manifold with the holomorphic atlas $\{(U_\alpha, \varphi_\alpha)\}_{\alpha \in A}$, $\dim_{\mathbb{C}} X = n$. Show that the functions $g_{\alpha\beta} = J(\varphi_\alpha \circ \varphi_\beta^{-1}) \circ \varphi_\beta$, where $J(f)$ is the Jacobian matrix of a holomorphic map f of a domain D in $\mathbb{C}^n$ to $\mathbb{C}^n$, are transition functions of the holomorphic tangent bundle $\mathcal{T}_X$ of X .

3. Show that the complex manifold is canonically oriented.

4. Let $D \subset \mathbb{C}^n$ be a polydisk, and suppose that $\omega \in \mathcal{A}^{1,1}(D)$ satisfies $d\omega = 0$ and is real-valued. Prove that there is a smooth function $f : D \rightarrow \mathbb{R}$ with the property that $\omega = i\bar{\partial}\partial f$.

5. Consider the Bochner-Martinelli kernel

$$k_{\text{BM}}(z) = c_n \sum_{k=1}^n (-1)^{k-1} \frac{\bar{z}_k}{|z|^{2n}} dz_1 \wedge \cdots \wedge dz_n \wedge d\bar{z}_1 \wedge \cdots \wedge \widehat{d\bar{z}_k} \wedge \cdots \wedge d\bar{z}_n,$$

where $c_n = (-1)^{\frac{n(n-1)}{2}} (n-1)! / (2\pi i)^n$, and $\widehat{d\bar{z}_k}$ means that the factor $d\bar{z}_k$ is omitted.

- (a) Prove that $dk_{\text{BM}} = \bar{\partial}k_{\text{BM}} = 0$ on $\mathbb{C}^n \setminus \{0\}$.
(b) Let $S^{2n-1} \subset \mathbb{R}^{2n} \simeq \mathbb{C}^n$ be the $(2n-1)$ -dimensional unit sphere.

Show that

$$\int_{S^{2n-1}} k_{\text{BM}} = 1.$$

(c) Let $\pi : \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}^n$ be the map $\pi(z, \zeta) = z - \zeta$, and let

$$K(z, \zeta) = \Pi_z^{0,0} \Pi_\zeta^{n,n-1} \pi^* k_{\text{BM}}$$

(note that projector $\Pi_z^{0,0}$ is actually redundant since k_{BM} has type $(n, n-1)$). Let $D \subset \mathbb{C}^n$ be a bounded domain with a C^1 boundary and let $f \in C^1(\bar{D})$. Prove Bochner-Martinelli formula:

$$f(z) = \int_{\partial D} K(z, \zeta) f(\zeta) + \int_D K(z, \zeta) \wedge \partial f(\zeta).$$

6. Let

$$K^{p,q}(z, \zeta) = \Pi_z^{p,q} \Pi_\zeta^{n-p,n-q-1} \pi^* k_{\text{BM}}$$

(the projector $\Pi_z^{p,q}$ is actually redundant), and let $\varphi \in \mathcal{A}_c^{p,q}(\mathbb{C}^n)$ be a differential form of type (p, q) with compact support. Prove Koppelman formula:

$$\varphi(z) = \bar{\partial}_z \int_{\mathbb{C}^n} K^{p,q-1}(z, \zeta) \wedge \varphi(\zeta) + \int_{\mathbb{C}^n} K^{p,q}(z, \zeta) \wedge \bar{\partial} \varphi(\zeta).$$

(Hint: Use relation $\bar{\partial}_\zeta K^{p,q} = -\bar{\partial}_z K^{p,q-1}$, which is derived from part (a) of problem 5).

7. Let $B = \{z \in \mathbb{C}^n : |z| < 1\}$ be the unit ball in $\mathbb{C}^n$ centered at 0, $S^{2n-1} = \partial B$ and let $f \in \mathcal{O}(\bar{B})$. Using the following steps, prove Leray-Fantappiè formula

$$f(z) = \int_{S^{2n-1}} f(\zeta) S(z, \zeta),$$

where

$$S(z, \zeta) = \frac{(n-1)!}{(2\pi i)^n} \sum_{k=1}^n (-1)^{k-1} \frac{\bar{\zeta}_k}{(1 - \bar{\zeta} \cdot z)^n} d\bar{\zeta}_1 \wedge \cdots \wedge \widehat{d\bar{\zeta}_k} \wedge \cdots \wedge d\bar{\zeta}_n \wedge d\zeta_1 \wedge \cdots \wedge d\zeta_n$$

and $\bar{\zeta} \cdot z = \bar{\zeta}_1 z_1 + \cdots + \bar{\zeta}_n z_n$.

(a) Put (as in class)

$$\begin{aligned} \mathcal{K}(z, w, \zeta) &= c_n \sum_{k=1}^n (-1)^{k-1} \frac{(w_k - \bar{\zeta}_k)}{((z - \zeta) \cdot (w - \bar{\zeta}))^n} (dz_1 - d\zeta_1) \wedge \cdots \wedge (dz_n - d\zeta_n) \wedge \\ &\quad (dw_1 - d\bar{\zeta}_1) \wedge \cdots \wedge (dw_{k-1} - d\bar{\zeta}_{k-1}) \wedge (dw_{k+1} - d\bar{\zeta}_{k+1}) \wedge \cdots \wedge (dw_n - d\bar{\zeta}_n) \end{aligned}$$

and

$$\mathcal{K}_0(z, w, \zeta) = \Pi_z^{0,0} \Pi_w^{0,0} \Pi_\zeta^{n,n-1} \mathcal{K}(z, w, \zeta),$$

so that

$$K(z, \zeta) = \mathcal{K}_0(z, \bar{z}, \zeta)$$

— a pullback of $\mathcal{K}_0$ under the map $\pi : \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}^n \times \mathbb{C}^n \times \mathbb{C}^n$ given by $(z, \zeta) \mapsto (z, \bar{z}, \zeta)$. For $f \in \mathcal{O}(\bar{B})$ and $z, w \in B$ put

$$g(z, w) = \int_{\partial B} \mathcal{K}_0(z, w, \zeta) f(\zeta).$$

Show that $g \in \mathcal{O}(B \times B)$.

- (b) Using Bochner-Martinelli formula, for $f \in \mathcal{O}(\bar{B})$ and $z \in B$ show that $f(z) = g(z, w)|_{w=\bar{z}}$.
 - (c) Prove that $\partial_w g(z, w) = 0$, so $g(z, w)$ does not actually depend on w . Get Leray-Fantappiè formula by putting $w = 0$ in the integral formula in part (a).
8. Prove that $f \in \mathcal{O}(B)$ admits a power series expansion in B , which converges absolutely and uniformly for z in every ball centered at 0 of radius $r < 1$.