

# THE PHASE-PARAMETER RELATION FOR ANALYTIC FAMILIES OF UNIMODAL MAPS

ARTUR AVILA

ABSTRACT. We prove the phase-parameter relation (in the sense of [AM1]) for analytic families of unimodal maps. This result enables very precise estimates on the dynamics of typical unimodal maps. Using the phase-parameter relation, we show that typical analytic unimodal maps admit a quasiquadratic renormalization. This reduces the study of typical unimodal maps to the quasiquadratic case which had been studied in [AM2]. As another application, we conclude that for typical analytic unimodal maps the exponent of the polynomial recurrence of the critical orbit is exactly 1. We also show that those ideas lead to a new proof of the Theorem of Shishikura on the Lebesgue measure of non-renormalizable parameters in the boundary of the Mandelbrot set.

## CONTENTS

1. Introduction	1
2. Preliminaries	4
3. Complex dynamics	6
4. Puzzle and parapuzzle geometry	9
5. Unimodal maps	11
6. Construction of the special family	14
7. Phase-parameter relation for the special family	20
8. Lamination in spaces of unimodal maps	21
9. Proof of Theorem A	24
10. Proof of Theorem B	26
11. Proof of corollaries	29
Appendix A. Hybrid classes	31
Appendix B. Non-renormalizable parameters in the Mandelbrot set	33
References	36

## 1. INTRODUCTION

A unimodal map  $f$  is a smooth (at least  $C^2$ ) self map of an interval. Let us say that  $f$  is regular if it has a quadratic critical point, is hyperbolic and its critical point is not periodic or preperiodic (this definition is such that regular maps coincide with structurally stable maps). The set of regular maps is (open and) dense in all topologies by a result of Kozłowski [K2].

The most studied family of unimodal maps is the quadratic family  $p_\lambda = \lambda - x^2$ ,  $-1/4 \leq \lambda \leq 2$ . In [AM1] it was shown that for a typical non-regular quadratic map

---

Partially supported by Faperj and CNPq, Brazil.

$p_{\lambda_0}$ , the phase space of  $p_{\lambda_0}$  near 0 and the parameter space near  $\lambda_0$  are related by some metric rules called the *Phase-Parameter relation* (notice that it is crucial that the phase and parameter of the quadratic family have the same dimension). The proof of [AM1] was tied to the combinatorial theory of the Mandelbrot set, so it can only work for quadratic maps (or, at most, full unfolded families of quadratic-like maps).

Let us say that an analytic family of unimodal maps is non-trivial if regular parameters are dense (in particular non-trivial analytic families are dense in any topology). The first main result of this paper is the following:

**Theorem A.** *Let  $f_\lambda$  be a one-parameter non-trivial analytic family of unimodal maps. Then  $f_\lambda$  satisfies the Phase-Parameter relation at almost every parameter.*

The Phase-Parameter relation has many remarkable consequences for the study of the dynamical behavior of typical parameters. Our second main result is an application of the Phase-Parameter relation:

**Theorem B.** *Let  $f_\lambda$  be a non-trivial analytic family of unimodal maps (any number of parameters). Then almost every parameter is either regular or has a renormalization which is topologically conjugate to a quadratic polynomial.*

This result allows one to reduce the study of typical unimodal maps to the special case of unimodal maps which are quasiquadratic (persistently topologically conjugate to a quadratic polynomial).

**1.1. Application: statistical properties of typical unimodal maps.** Typical quasiquadratic maps had been previously studied in [AM2]. Their main result is that the dynamics of typical quasiquadratic maps have an excellent statistical description (in terms of physical measures, decay of correlations and stochastic stability), thus answering the Palis Conjecture (see [AM2] for details) in the unimodal quasiquadratic case.

For regular maps, the good statistical description comes for free. For a non-regular map  $f$ , it is related to essentially two properties regarding its critical point  $c$ : the Collet-Eckmann condition<sup>1</sup> and subexponential recurrence<sup>2</sup>.

Thus, [AM2] achieves the good statistical description via a dichotomy: typical quasiquadratic maps are either regular or Collet-Eckmann and subexponentially recurrent. This is done in both the analytic case as in the smooth case ( $C^k$ ,  $k = 3, \dots, \infty$ ). For typical non-regular analytic unimodal maps, it is proved even more, that the critical point is polynomially recurrent<sup>3</sup>.

Our Theorem B allows to immediately obtain the analytic case in our more general setting (see Theorem 11.1 for a more precise statement):

**Corollary C.** *Let  $f_\lambda$  be a non-trivial analytic family of unimodal maps (in any finite number of parameters). Then almost every non-regular parameter is Collet-Eckmann and its critical point is polynomially recurrent.*

This allows us not only to generalize the smooth case of [AM2] besides quasiquadratic maps, but to reduce the differentiability requirements, including the  $C^2$  case in the description (see Theorem 11.3 for a more precise statement):

---

<sup>1</sup>A unimodal map  $f$  is Collet-Eckmann if  $|Df^n(f(c))| > C\lambda^n$  for some constants  $C > 0$  and  $\lambda > 1$ .

<sup>2</sup>That is, for every  $\alpha > 0$ ,  $|f^n(c) - c| > e^{-\alpha n}$  for  $n$  sufficiently big.

<sup>3</sup>That is, there exists  $\gamma > 0$  such that  $|f^n(c) - c| > n^{-\gamma}$  for every  $n$  sufficiently big.

**Corollary D.** *In generic smooth ( $C^k$ ,  $k = 2, \dots, \infty$ ) families of unimodal maps (any number of parameters), almost every parameter is regular, or has a renormalization which is conjugate to a quadratic map, Collet-Eckmann and its critical point is subexponentially recurrent.*

(Theorem C of [ALM] considered only families which are at least  $C^3$ .)

*Remark 1.1.* The dichotomies in Corollaries C and D implies that the dynamics of typical non-regular unimodal maps have the same excellent statistical description of the quasiquadratic case studied by [AM2], see also Remark 11.1 for a list of references. In particular, our Corollaries C and D give an answer to the Palis Conjecture in the general unimodal case.

The Phase-Parameter relation allows to obtain very precise estimates on the dynamics of typical parameters. For instance, the statistical analysis of [AM1] allows one to compute the exact exponent of the polynomial recurrence<sup>4</sup>. (In [AM2], it was impossible to estimate the exponent even in the quasiquadratic case.)

**Corollary E.** *Let  $f_\lambda$  be a non-trivial analytic family of unimodal maps (any number of parameters). Then almost every parameter is either regular or has a polynomially recurrent critical point with exponent 1.*

We call the attention of the reader to [AM3] where much more refined statistical applications of Theorem A are obtained (which are inaccessible with the methods of [AM2]).

**1.2. Complex parameters.** A very natural question raised by the description of typical parameters in the real quadratic family is if the results generalize to complex parameters. It is widely expected that the description should be actually simpler: almost every complex parameter should be hyperbolic. However, only partial results are available.

In this direction, let us remark that the argument of the proof of Theorem B can be also applied in the complex setting, and leads to a new proof of the following result of Shishikura (unpublished):

**Theorem F.** *The set of non-hyperbolic, non-infinitely renormalizable complex quadratic parameters has zero Lebesgue measure.*

We discuss this application in Appendix B.

**1.3. Outline of the proof of Theorem A.** The proof of Theorem A can be divided in four parts. The crucial step of this paper is step (2) below, which allows to integrate the work of [AM1] (step (1)), and [ALM] (step (3)).

(1) We study the complex phase-parameter relation for certain families of complex return type maps, which model complex extensions of the return maps  $R_n : I_n \rightarrow I_n$  to the principal nest of a unimodal map  $f$ . This analysis is based on the scheme of [AM1] which is based on [L3]. The class of families which we will study are so called full families, and we prove a complex analogous of the Phase-Parameter relation for them.

(2) We show that through any given analytic unimodal map  $f$  (assumed finitely renormalizable with a recurrent critical point), there exists an analytic family  $\tilde{f}_\lambda$

---

<sup>4</sup>The exponent of the polynomial recurrence of the critical point  $c$  of a unimodal maps  $f$  is the infimum of all  $\gamma > 0$  such that for  $n$  sufficiently big  $|f^n(c) - c| > n^{-\gamma}$ .

which gives rise to a full family of complex return type maps. This step is based on the perturbation idea of [ALM] which is used in that paper to show existence of a transverse direction to the topological class of  $f$ . Using the argument of [ALM] we can also show that this analytic family is transverse to the topological class of  $f$ . Using step (1), we conclude that the Phase-Parameter relation is valid at  $f$  for this special family  $\tilde{f}_\lambda$ .

(3) We show that if the Phase-Parameter relation is valid for one transverse family at  $f$ , then it is valid for all transverse families at  $f$ . This step is heavily based on the results of [ALM]: in order to compare the parameter space of both families, one uses the local holonomy of the lamination associated to the partition of spaces of unimodal maps in topological classes.

(4) Using a simple generalization of [ALM] we conclude that a non-trivial family of unimodal maps is transverse to the topological class of almost every non-regular parameter. This concludes the proof of Theorem A.

**1.4. Structure of the paper.** In §2 we give some basic background on quasi-conformal maps and holomorphic motions. In §3 we recall the work of [AM1] to deal with the so-called  $R$ -chains (sequences of families of return-type maps related by renormalization). In §3.5 we state the Complex Phase-Parameter relation for chains, which follows from [AM1]. In §4 we discuss the results of Lyubich in [L2] and [L3], and consider extensions of [L3] to more general chains than quadratic ones. In §5 we introduce the basic theory of unimodal maps. In §6 we construct a special analytic family of unimodal maps which induce a full family of return type maps, and in §7 we state and prove the Phase-Parameter relation for the special family. In §8 we introduce the results of [ALM] on the lamination structure of topological classes of unimodal maps and state some straightforward generalizations (some details are given in Appendix A). In §9 and §10 we prove Theorems A and B, and in §11 we show the relation to the corollaries. In Appendix B we give a proof of Theorem F.

**Acknowledgements:** Most of the results of this paper were announced in [Av2], and formed my thesis together with [AM1], [AM2] and [ALM]. I would like to thank Wellington de Melo, Mikhail Lyubich and Carlos Gustavo Moreira who collaborated with me in those works, and to Viviane Baladi and Jean-Christophe Yoccoz for useful conversations.

## 2. PRELIMINARIES

**2.1. General notation.** Let  $\Omega$  be the set of finite sequences (possibly empty) of non-zero integers  $\underline{d} = (j_1, \dots, j_m)$ .

A *Jordan curve*  $T$  is a subset of  $\mathbb{C}$  homeomorphic to a circle. A *Jordan disk* is a bounded open subset  $U$  of  $\mathbb{C}$  such that  $\partial U$  is a Jordan curve.

We let  $\mathbb{D}_r(w) = \{z \in \mathbb{C} \mid |z - w| < r\}$ . Let  $\mathbb{D}_r = \mathbb{D}_r(0)$ , and  $\mathbb{D} = \mathbb{D}_1$ . If  $r > 1$ , let  $A_r = \{z \in \mathbb{C} \mid 1 < |z| < r\}$ . An *annulus*  $A$  is a subset of  $\mathbb{C}$  such that there exists a conformal map from  $A$  to some  $A_r$ . In this case,  $r$  is uniquely defined and we denote the *modulus* of  $A$  as  $\text{mod}(A) = \ln(r)$ .

**2.2. Graphs and sections.** Let us fix a complex Banach space  $\mathbb{E}$ . If  $\Lambda \subset \mathbb{E}$ , a *graph* of a continuous map  $\phi : \Lambda \rightarrow \mathbb{C}$  is the set of all  $(z, \phi(z)) \in \mathbb{E} \oplus \mathbb{C}$ ,  $z \in \Lambda$ .

Let  $\mathbf{0} : \mathbb{E} \rightarrow \mathbb{E} \oplus \mathbb{C}$  be defined by  $\mathbf{0}(z) = (z, 0)$ .

Let  $\pi_1 : \mathbb{E} \oplus \mathbb{C} \rightarrow \mathbb{E}$ ,  $\pi_2 : \mathbb{E} \oplus \mathbb{C} \rightarrow \mathbb{C}$  be the coordinate projections. Given a set  $\mathcal{X} \subset \mathbb{E} \oplus \mathbb{C}$  we denote its fibers  $X[z] = \pi_2(\mathcal{X} \cap \pi_1^{-1}(z))$ .

A *fiberwise map*  $\mathcal{F} : \mathcal{X} \rightarrow \mathbb{E} \oplus \mathbb{C}$  is a map such that  $\pi_1 \circ \mathcal{F} = \pi_1$ . We denote its fibers  $F[z] : X[z] \rightarrow \mathbb{C}$  such that  $\mathcal{F}(z, w) = (z, F[z](w))$ .

Let  $B_r(\mathbb{E})$  be the ball of radius  $r$  around 0.

**2.3. Quasiconformal maps and quasisymmetric maps.** Let  $U \subset \mathbb{C}$  be a domain. A map  $h : U \rightarrow \mathbb{C}$  is *K-quasiconformal* (K-qc) if it is a homeomorphism onto its image and for any annulus  $A \subset U$ ,  $\text{mod}(A)/K \leq \text{mod}(h(A)) \leq K \text{mod}(A)$ . The minimum such  $K$  is called the *dilatation*  $\text{Dil}(h)$  of  $h$ .

A homeomorphism  $h : \mathbb{R} \rightarrow \mathbb{R}$  is said to be  $\gamma$ -quasisymmetric if it has a real-symmetric extension  $h : \mathbb{C} \rightarrow \mathbb{C}$  which is quasiconformal with dilatation bounded by  $\gamma$ . If  $X \subset \mathbb{R}$ , we will also say that  $h : X \rightarrow \mathbb{R}$  is  $\gamma$ -qs if it has a  $\gamma$ -qs extension.

**2.4. Holomorphic motions.** Let  $\Lambda$  be a connected open set of a Banach space  $\mathbb{E}$ . A *holomorphic motion*  $h$  over  $\Lambda$  is a family of holomorphic maps defined on  $\Lambda$  whose graphs (called *leaves* of  $h$ ) do not intersect. The *support* of  $h$  is the set  $\mathcal{X} \subset \mathbb{C}^2$  which is the union of the leaves of  $h$ .

We have naturally associated maps  $h[z] : \mathcal{X} \rightarrow X[z]$ ,  $z \in \Lambda$  defined by  $h[z](x, y) = w$  if  $(z, w)$  and  $(x, y)$  belong to the same leaf. The *transition* (or holonomy) maps  $h[z, w] : X[z] \rightarrow X[w]$ ,  $z, w \in \Lambda$ , are defined by  $h[z, w](x) = h[w](z, x)$ .

Given a holomorphic motion  $h$  over a domain  $\Lambda$ , a holomorphic motion  $h'$  over a domain  $\Lambda' \subset \Lambda$  whose leaves are contained in leaves of  $h$  is called a *restriction* of  $h$ . If  $h$  is a restriction of  $h'$  we also say that  $h'$  is an *extension* of  $h$ .

Let  $K : [0, 1) \rightarrow \mathbb{R}$  be defined by  $K(r) = (1 + \rho)/(1 - \rho)$  where  $0 \leq \rho < 1$  is such that the hyperbolic distance in  $\mathbb{D}$  between 0 and  $\rho$  is  $r$ .

**$\lambda$ -Lemma** ([MSS], [BR]) *Let  $h$  be a holomorphic motion over a hyperbolic domain  $\Lambda \subset \mathbb{C}$  and let  $z, w \in \Lambda$ . Then  $h[z, w]$  extends to a quasiconformal map of  $\mathbb{C}$  with dilatation bounded by  $K(r)$ , where  $r$  is the hyperbolic distance between  $z$  and  $w$  in  $\Lambda$ .*

*In the general case ( $\Lambda$  not one-dimensional), the same estimate holds with the Kobayashi distance instead of the hyperbolic distance. In particular, if  $h$  is a holomorphic motion over  $B_r(\mathbb{E})$ , and if  $z, w \in B_{r/2}(\mathbb{E})$  then  $h[z, w] = O(\|z - w\|)$ .*

If  $h = h_U$  is a holomorphic motion of an open set, we define  $\text{Dil}(h)$  as the supremum of the dilatations of the maps  $h[z, w]$ .

A *completion* of a holomorphic motion means an extension of  $h$  to the whole complex plane:  $X[z] = \mathbb{C}$  for all  $z \in \Lambda$ . The problem of existence of completions is considerably different in one-dimension or higher:

**Extension Lemma** ([Sl]) *Any holomorphic motion over a simply connected domain  $\Lambda \subset \mathbb{C}$  can be completed.*

**Canonical Extension Lemma** ([BR]) *Let  $h$  be a holomorphic motion over  $B_r(\mathbb{E})$ . Then the restriction of  $h$  to  $B_{r/3}(\mathbb{E})$  can be completed in a canonical way.*

**2.4.1. Symmetry.** Let us assume that  $\mathbb{E}$  is the complexification of a real-symmetric space  $\mathbb{E}^{\mathbb{R}}$ , that is, there is a anti-linear isometric involution  $\text{conj}$  fixing  $\mathbb{E}^{\mathbb{R}}$ . Let us use  $\text{conj}$  to denote also the map  $(z, w) \rightarrow (\text{conj } z, \overline{w})$  in  $\mathbb{E} \oplus \mathbb{C}$ .

A set  $X \subset \mathbb{E}, \mathbb{E} \oplus \mathbb{C}$  is called real-symmetric if  $\text{conj}(X) = X$ . Let  $\Lambda \subset \mathbb{E}$  be a real-symmetric domain. A holomorphic motion  $h$  over  $\Lambda$  is called real-symmetric if the image of any leaf by  $\text{conj}$  is also a leaf.

The systems we are interested on are real, so they naturally possess symmetry. In many cases, we will consider a real-symmetric holomorphic motion associated to the system, which will need to be completed using the Extension Lemma (in one-dimension) or the Canonical extension lemma (in higher dimensions).

Since the Canonical Extension Lemma is canonical, it can be used to produce real-symmetric holomorphic motions out of real-symmetric holomorphic motions. On the other hand, the Extension Lemma adds ambiguity on the procedure, since the extension is not unique. In particular, this could lead to loss of symmetry. In order to avoid this problem, we will choose a little bit more carefully our extensions. Indeed, in [AM1] it is remarked that the proof of the Extension Lemma actually shows:

**Real Extension Lemma.** *Any real-symmetric holomorphic motion can be completed to a real-symmetric holomorphic motion.*

So we can make the following:

**Symmetry assumption.** *Extensions of real-symmetric motions will always be taken real-symmetric.*

More precise reference

**2.4.2. Notation warning.** We will use the following conventions. Instead of talking about the sets  $X[z]$ , fixing some  $z \in \Lambda$ , we will say that  $h$  is the motion of  $X$  over  $\Lambda$ , where  $X$  is to be thought of as a set which depends on the point  $z \in \Lambda$ . In other words, we usually drop the brackets from the notation.

We will also use the following notation for restrictions of holomorphic motions: if  $Y \subset X$ , we denote  $\mathcal{Y} \subset \mathcal{X}$  as the union of leaves through  $Y$ .

**2.5. Codimension-one laminations.** Let  $F$  be a Banach space. A codimension-one *holomorphic lamination*  $\mathcal{L}$  on an open subset  $\mathcal{W} \subset F$  is a family of disjoint codimension-one Banach submanifolds of  $F$ , called the *leaves* of the lamination such that for any point  $p \in \mathcal{W}$ , there exists a holomorphic local chart  $\Phi : \mathcal{W} \rightarrow \mathcal{V} \oplus \mathbb{C}$  where  $\mathcal{V}$  is a neighborhood in some complex Banach space  $\mathbb{E}$ , such that for any leaf  $L$  and any connected component  $L_0$  of  $L \cap \mathcal{W}$ , the image  $\Phi(L_0)$  is a graph of a holomorphic function  $\mathcal{V} \rightarrow \mathbb{C}$ .

The neighborhood  $\mathcal{W}$  in the above definition is called a *flow box*, and the connected components  $L_0$  are called *local leaves* in this flow box.

It is clear that the local theory of codimension-one laminations is the theory of holomorphic motions. For instance, the  $\lambda$ -Lemma imply that holonomy maps of codimension-one laminations have quasiconformal extensions, and gives bounds on the dilatation of those extensions.

### 3. COMPLEX DYNAMICS

In this section, we will deal exclusively with one-dimensional holomorphic motions over some Jordan domain  $\Lambda \subset \mathbb{C}$ .

**3.1.  $R$ -maps and  $L$ -maps.** Let  $U$  be a Jordan disk and  $U^j$ ,  $j \in \mathbb{Z}$  be a family of Jordan disks with disjoint closures such that  $\overline{U^j} \subset U$ . A holomorphic map  $R : \cup U^j \rightarrow U$  surjective in each component is called a  $R$ -map (return type map) if for  $j \neq 0$ ,  $R|_{U^j}$  extends to a homeomorphism onto  $\overline{U}$  and  $R|_{U^0}$  extends to a double covering map onto  $\overline{U}$  ramified at 0.

Given a  $R$ -map  $R$  we induce an  $L$ -map (landing type map) as follows. For  $\underline{d} \in \Omega$ ,  $\underline{d} = (j_1, \dots, j_m)$ , we define  $U^{\underline{d}} = \{z \in U | R^{k-1}(z) \in U^{j_k}, 1 \leq k \leq m\}$  and we let  $R^{\underline{d}} = R^m|_{U^{\underline{d}}}$ . Let  $W^{\underline{d}} = (R^{\underline{d}})^{-1}(U^0)$ . The  $L$ -map associated to  $R$  is defined as  $L(R) : \cup W^{\underline{d}} \rightarrow U^0$ ,  $L(R)|_{W^{\underline{d}}} = R^{\underline{d}}$ .

**3.1.1. Renormalization.** Given a  $R$ -map  $R$  such that  $R(0) \in \cup W^{\underline{d}}$  we can define the (generalized in the sense of Lyubich) *renormalization*  $N(R)$  by  $N(R) = L(R) \circ R$  where defined in  $U^0$ : its domain is the  $R$ -puzzle  $(V, V^j)$  such that  $V = U^0$  and the  $V^j$  are connected components of  $(R|_{U^0})^{-1}(\cup W^{\underline{d}})$ .

**3.2. Tubes and tube maps.** A *proper motion* of a set  $X$  over  $\Lambda$  is a holomorphic motion of  $X$  over  $\Lambda$  such that the map  $\mathbf{h}[z] : \Lambda \times X[z] \rightarrow \mathcal{X}$  defined by  $\mathbf{h}[z](w, x) = (w, h[z](w, x))$  has an extension to  $\overline{\Lambda} \times X[z]$  which is a homeomorphism.

An *equipped tube*  $h_T$  is a holomorphic motion of a Jordan curve  $T$ . Its support is called a *tube*. We say that an equipped tube is *proper* if it is a proper motion. Its support is called a *proper tube*. The *filling* of a tube  $\mathcal{T}$  is the set  $\mathcal{U} \subset \Lambda \times \mathbb{C}$  such that  $U[z]$  is the bounded component of  $\mathbb{C} \setminus T[z]$ ,  $z \in \Lambda$ .

A *special motion* is a holomorphic motion  $h = h_{X \cup T}$  such that  $\mathcal{X}$  is contained in the filling  $\mathcal{U}$  of  $\mathcal{T}$ ,  $h|_T$  is an equipped proper tube and the closure of any leaf through  $X$  does not intersect the closure of  $\mathcal{T}$ .

If  $\mathcal{T}$  is a tube over  $\Lambda$ , and  $\mathcal{U}$  is its filling, a fiberwise map  $\mathcal{F} : \mathcal{U} \rightarrow \mathbb{C}^2$  is called a *tube map* if it admits a continuous extension to  $\overline{\mathcal{U}}$ .

**3.2.1. Tube pullback.** Let  $\mathcal{F} : \mathcal{V} \rightarrow \mathbb{C}^2$  be a tube map such that  $\mathcal{F}(\partial\mathcal{V}) = \partial\mathcal{U}$ , where  $\mathcal{U}$  is the filling of a tube over  $\Lambda$  and let  $h$  be a holomorphic motion supported on  $\overline{\mathcal{U}} \cap \pi_1^{-1}(\Lambda)$ .

Let  $\Gamma$  be a (parameter) open set such that  $\overline{\Gamma} \subset \Lambda$  and  $W$  be a (phase) open set which moves holomorphically by  $h$  over  $\Lambda$  and such that  $\overline{W} \subset U$ . Assume that  $W$  contains critical values of  $\mathcal{F}|(\mathcal{V} \cap \pi_1^{-1}(\Gamma))$ , that is, if  $\lambda \in \Gamma$ ,  $z \in V[\lambda]$  and  $DF[\lambda](z) = 0$  then  $F[\lambda](z) \in W[\lambda]$ .

Let us consider a leaf of  $h$  through  $z \in U \setminus \overline{W}$ , and let us denote by  $\mathcal{E}(z)$  its preimage by  $\mathcal{F}$  intersected with  $\pi_1^{-1}(\Gamma)$ . Each connected component of  $\mathcal{E}(z)$  is a graph over  $\Gamma$ , moreover,  $\overline{\mathcal{E}}(z) \subset \mathcal{U}$ . So the set of connected components of  $\mathcal{E}(z)$ ,  $z \in U \setminus \overline{W}$  is a holomorphic motion over  $\Gamma$ .

We define a new holomorphic motion over  $\Gamma$ , called *the lift of  $h$  by  $(\mathcal{F}, \Gamma, W)$* , as an extension to the closure of  $V$  of the holomorphic motion whose leaves are the connected components of  $\mathcal{E}(z)$ ,  $z \in U \setminus \overline{W}$  (the lift is not uniquely defined). It is clear that this holomorphic motion is a special motion of  $V$  over  $\Gamma$  and its dilatation over  $F^{-1}(U \setminus \overline{W})$  is bounded by  $K(r)$  where  $r$  is the hyperbolic diameter of  $\Gamma$  in  $\Lambda$ .

**3.2.2. Diagonal and phase-parameter holonomy maps.** Let  $h$  be an equipped proper tube supported on  $\mathcal{T}$ . A *diagonal* of  $\mathcal{T}$  is a holomorphic section  $\Psi : \Lambda \rightarrow \mathbb{C}^2$  (so that  $\pi_1 \circ \Psi = \pi_1$ ), admitting a continuous extension to  $\Lambda$ , and such that  $\Psi(\Lambda)$  is contained on the filling of  $\mathcal{T}$  and for  $\lambda \in \Lambda$ ,  $h[\lambda] \circ \Psi|_{\partial\Lambda}$  has degree one onto  $T[\lambda]$ .

Let  $h = h_{X \cup T}$  be a special motion and let  $\Phi$  be a diagonal of  $h|T$ . It is a consequence of the Argument Principle (see [L3]) that the leaves of  $h|X$  intersect  $\Phi(\Lambda)$  in a unique point (with multiplicity one). From this we can define a map  $\chi[\lambda] : X[\lambda] \rightarrow \Lambda$  such that  $\chi[\lambda](z) = w$  if  $(\lambda, z)$  and  $\Phi(w)$  belong to the same leaf of  $h$ . It is clear that each  $\chi[\lambda]$  is a homeomorphism onto its image, moreover, if  $U \subset X$  is open,  $\chi[\lambda]|U[\lambda]$  is locally quasiconformal, and if  $\text{Dil}(h|U) < \infty$  then  $\chi[\lambda]|U[\lambda]$  is globally quasiconformal with dilatation bounded by  $\text{Dil}(h|U)$ .

We will say that  $\chi$  is the *holonomy family* associated to the pair  $(h, \Phi)$ .

**3.3. Families of  $R$ -maps.** A  $R$ -family is a pair  $(\mathcal{R}, h)$ , where  $\mathcal{R}$  is a holomorphic map  $\mathcal{R} : \mathcal{U}^j \rightarrow \mathcal{U}$  such that the fibers  $\mathcal{R}[\lambda]$  of  $\mathcal{R}$  are  $R$ -maps, for every  $j$ ,  $\mathcal{R}|U^j$  is a tube map, and  $h$  is a holomorphic motion  $h = h_{\overline{U}}$  such that  $h|(\partial U \cup \cup_j \partial U_j)$  is special. If additionally  $\mathcal{R} \circ \mathbf{0}$  is a diagonal to  $h$ , we say that the  $\mathcal{R}$  is *full*.

**3.3.1. From  $R$ -families to  $L$ -families.** Given a  $R$ -family  $\mathcal{R}$  with motion  $h = h_{\overline{U}}$  we induce a family of  $L$ -maps as follows. If  $\underline{d} \in |\text{omega}|$ ,  $\underline{d} = (j_1, \dots, j_m)$ , we let  $\mathcal{U}^{\underline{d}} = \{(\lambda, z) \in \mathcal{U} | R_{\lambda}^{k-1}(z) \in U^{j_k}[\lambda]\}$  and define  $\mathcal{R}^{\underline{d}} = \mathcal{R}^m|U^{\underline{d}}$ . Let  $\mathcal{W}^{\underline{d}} = (\mathcal{R}^{\underline{d}})^{-1}(\mathcal{U}^0)$ . We define  $L(\mathcal{R}) : \cup \mathcal{W}^{\underline{d}} \rightarrow \mathcal{U}^0$  by  $L(\mathcal{R})|W^{\underline{d}} = \mathcal{R}^{\underline{d}}$ . Notice that the  $L$ -maps which are associated with the fibers of  $\mathcal{R}$  coincide with the fibers of  $L(\mathcal{R})$ .

We define a holomorphic motion  $L(h)$  in the following way. The leaf through  $z \in \partial U$  is the leaf of  $h$  through  $z$ . If there is a smallest  $U^{\underline{d}}$  such that  $z \in U^{\underline{d}}$ , we let the leaf through  $z$  be the preimage by  $\mathcal{R}^{\underline{d}}$  of the leaf through  $\mathcal{R}^{\underline{d}}(z)$ . We finally extend it to  $\overline{U}$  using the Extension Lemma. The  $L$ -family associated to  $(\mathcal{R}, h)$  is a pair  $(L(\mathcal{R}), L(h))$ .

**3.3.2. Parameter partition.** Let  $(\mathcal{R}, h)$  be a full  $R$ -family. Since  $L(h)|(\mathcal{U} \cup \cup_j \overline{U^j})$  is special, we can consider the holonomy family of the pair  $(L(h)|(\mathcal{U} \cup \cup_j \overline{U^j}), \mathcal{R}(\mathbf{0}))$ , which we denote by  $\chi$ . We use  $\chi$  to partition  $\Lambda$ : let  $\Lambda^{\underline{d}} = \chi(U^{\underline{d}})$  and let  $\Gamma^{\underline{d}} = \chi(W^{\underline{d}})$ .

**3.3.3. Family renormalization.** Let  $(\mathcal{R}, h)$  be a full  $R$ -family. The  $\underline{d}$  renormalization of  $(\mathcal{R}, h)$  is the  $R$ -family  $(N^{\underline{d}}(\mathcal{R}), N^{\underline{d}}(h))$  over  $\Gamma^{\underline{d}}$  defined as follows. We take  $N^{\underline{d}}(h)$  as the lift of  $L(h)$  by  $(\mathcal{R}|U^0, \Gamma^{\underline{d}}, W^{\underline{d}})$  where defined.

It is clear that  $(N^{\underline{d}}(\mathcal{R}), N^{\underline{d}}(h))$  is full, and its fibers are renormalizations of the fibers of  $(\mathcal{R}, h)$ . Moreover,  $N^{\underline{d}}(h)$  is a special motion.

**3.3.4. Truncation and gape renormalization.** Let  $(\mathcal{R}, h)$  be a full  $R$ -family and let  $\underline{d} \in \Omega$ . We define the  $\underline{d}$  truncation of  $L(\mathcal{R})$  as  $L^{\underline{d}}(\mathcal{R}) = L(\mathcal{R})$  outside  $U^{\underline{d}}$  and  $L^{\underline{d}}(\mathcal{R}) = \mathcal{R}^{\underline{d}}$  in  $U^{\underline{d}}$ . Let  $G^{\underline{d}}(\mathcal{R}) = L^{\underline{d}}(\mathcal{R}) \circ \mathcal{R}|(\mathcal{U}_0 \cap \pi_1^{-1}(\Lambda^{\underline{d}}))$  where defined.

If  $\underline{d}$  is empty, let  $G^{\underline{d}}(h) = L(h)$ . Otherwise, let  $G^{\underline{d}}(h)$  be a holomorphic motion of  $U$  over  $\Lambda^{\underline{d}}$ , which coincides with  $L(h)$  on  $U \setminus \overline{U^0}$  and coincides with the lift of  $L(h)$  by  $(\mathcal{R}|U^0, \Lambda^{\underline{d}}, U^{\underline{d}})$  on  $U_0$ .

The  $\underline{d}$  gape renormalization of  $(\mathcal{R}, h)$  is the pair  $(G^{\underline{d}}(\mathcal{R}), G^{\underline{d}}(h))$ .

**3.4. Chains.** Assume now that we are given a full  $R$ -family, which we will denote  $\mathcal{R}_1$  (over some domain  $\Lambda_1$ , with motion  $h_1$ ) together with a parameter  $\lambda \in \mathbb{R}$ . If  $\lambda$  belongs to some renormalization domain (that is, there exists  $\underline{d}_1$  such that  $\lambda \in \Lambda_1^{\underline{d}_1}$ ), let  $\mathcal{R}_2 = N_1^{\underline{d}_1}(\mathcal{R}_1)$  (over  $\Lambda_2$ ). Assume we can continue this process constructing  $\mathcal{R}_{i+1} = N_i^{\underline{d}_i}$ ,  $n \geq 1$ . Then the sequence  $\mathcal{R}_i$  (over  $\Lambda_i$ ) will be called the  $\mathcal{R}$ -chain over  $\lambda$ . The holomorphic motion associated to  $\mathcal{R}_i$  is denoted  $h_i$  (so that  $h_{i+1} = N_i^{\underline{d}_i}(h_i)$ ),



To simplify the notation for the gape renormalization, we let  $G^{d_i}(\mathcal{R}_i) = G(\mathcal{R}_i)$  and  $G^{d_i}(h_i) = G(h_i)$ . Let  $\tau_i$  be such that  $R_i[\lambda_0](0) \in U_i^{\tau_i}$ .

**3.4.1. Holonomy maps.** Notice that for  $i > 1$ , the holomorphic motion  $h_i$  is special (since it is obtained by renormalization and so coincides with  $N^{d_{i-1}}(h_{i-1})$ ). In particular, we can consider the holonomy family associated to  $(h_i, \mathcal{R}_i \circ \mathbf{0})$ , which we denote by  $\chi_i^0 : U_i \rightarrow \Lambda_i$ .

For  $i > 1$ ,  $L(h_i)$  is also special, let  $\chi_i : U_i \rightarrow \Lambda_i$  be the holonomy family associated to  $(L(h_i), \mathcal{R}_i \circ \mathbf{0})$ .

For  $i > 2$ ,  $G(h_{i-1})$  is also special, let  $\tilde{\chi}_i : U_{i-1} \rightarrow \tilde{\Lambda}_i$  be the holonomy family of the pair  $(G(h_{i-1}), G(\mathcal{R}_{i-1}) \circ \mathbf{0})$ .

**3.4.2. Real chains.** A fiberwise map  $\mathcal{F} : \mathcal{X} \rightarrow \mathbb{C}^2$  is real-symmetric if  $\mathcal{X}$  is real-symmetric and  $\mathcal{F} \circ \text{conj} = \text{conj} \circ \mathcal{F}$ . We will say that a chain  $\{\mathcal{R}_i\}$  over a parameter  $\lambda \in \mathbb{R}$  is real-symmetric if each  $\mathcal{R}_i$  and each underlying holomorphic motion  $h_i$  is real-symmetric.

Because of the Symmetry assumption, a chain  $\{\mathcal{R}_i\}$  over a parameter  $\lambda \in \mathbb{R}$  is real-symmetric provided the first step data  $\mathcal{R}_1$  and  $h_1$  is real-symmetric. In this case, all objects related to the chain are real-symmetric, including the holonomy families  $\chi_i^0, \chi_i$  and  $\tilde{\chi}_i$ .

### 3.5. Complex Phase-Parameter relation.

**Definition 3.1.** Let us say that a  $R$ -chain  $\mathcal{R}_i$  over  $\lambda_0$  satisfies the Complex Phase-Parameter relation if for every  $\gamma > 1$ , there exists  $i_0$  such that for  $i > i_0$  the following holds:

**CPhPa1:**  $\chi_i[\lambda_0]|U_i^{\tau_i}$  admits an extension to a  $\gamma$ -qc map of  $\mathbb{C}$ ,

**CPhPa2:**  $\tilde{\chi}_i[\lambda_0]|U_i$  admits an extension to a  $\gamma$ -qc map of  $\mathbb{C}$ ,

**CPhPh1:** For  $\lambda \in \Lambda_i^{\tau_i}$ ,  $L(h_i)[\lambda_0, \lambda]|U_i$  admits an extension to a  $\gamma$ -qc map of  $\mathbb{C}$ ,

**CPhPh2:** For  $\lambda \in \Lambda_i$ ,  $G(h_{i-1})[\lambda_0, \lambda]|U_i$  admits an extension to a  $\gamma$ -qc map of  $\mathbb{C}$ .

And, moreover, if  $\mathcal{R}_i$  is real-symmetric, all the above extensions can be taken real-symmetric.

The following result is a direct consequence of the arguments of §5 of [AM1], see Remark 5.3 of that paper.

**Theorem 3.1.** *Let  $\mathcal{R}_i$  be a  $R$ -chain over  $\lambda_0$ , and assume that  $\text{mod}(\Lambda_i \setminus \overline{\Lambda_{i+1}}) \rightarrow \infty$  and  $\text{mod}(U_i[\lambda_0] \setminus \overline{U_i^0[\lambda_0]}) \rightarrow \infty$ . Then  $\mathcal{R}_i$  satisfies the complex phase-parameter relation.*

## 4. PUZZLE AND PARAPUZZLE GEOMETRY

**4.1. Puzzle estimates.** The following result is contained on (the proof of) Theorem II of [L2]:

**Theorem 4.1.** *For every  $C > 0$ , there exists  $C' > 0$  with the following property. Let  $R_i$  be a sequence of  $R$ -maps such that  $R_{i+1} = N(R_i)$  and let  $n_k - 1$  be the sequence of non-central levels. If  $\text{mod}(U_1 \setminus \overline{U_1^0}) > C'$  then  $\text{mod}(U_{n_k} \setminus \overline{U_{n_k}^0}) > C$ .*

(In Lyubich's notation,  $R$ -maps are called generalized quadratic maps.)

The following result is Theorem III of [L2]:

Define non-central returns and levels

**Theorem 4.2.** *For every  $C' > 0$ , there exists  $C'' > 0$  with the following property. Let  $R_i$  be a sequence of  $R$ -maps such that  $R_{i+1} = N(R_i)$  and let  $n_k - 1$  be the sequence of non-central levels. If  $\text{mod}(U_1 \setminus \overline{U_1^0}) > C'$  then  $\text{mod}(U_{n_k} \setminus \overline{U_{n_k}^0}) > C''k$ .*

#### 4.2. Parapuzzle estimates.

4.2.1. *The quadratic family.* Let  $p_c(z) = z^2 + c$  be the quadratic family. The following result is contained in Lemma 3.6 of [L3]:

**Theorem 4.3.** *Let us fix a non-renormalizable quadratic polynomial  $p_{c_0}$  with a recurrent critical point and no neutral periodic orbits. Then there exists a full  $R$ -family  $\mathcal{R}_1$  over some  $c_0 \in \Lambda_1$  such that if  $c \in \Lambda_1$  then  $R[c] : \cup U_1^j[c] \rightarrow U_1[c]$  is the first return map under iteration by  $p_c$ .*

The following is Theorem A of [L3]:

**Theorem 4.4.** *In the setting of Theorem 4.3, let  $\mathcal{R}_i$  be the  $R$ -chain over  $c_0$  with first step  $\mathcal{R}_1$ . If  $n_k - 1$  denotes the  $k$ -th non-central return, then  $\text{mod}(\Lambda_{n_k} \setminus \overline{\Lambda_{n_k+1}}) > Tk$ , for some constant  $T > 0$ .*

*Remark 4.1.* In Lyubich's notation he lets  $\Delta^i = \Lambda_{n_i}$  and  $\Pi^i = \Lambda_{n_i}^0$ . He states that both  $\text{mod}(\Delta^i \setminus \overline{\Delta^{i+1}})$  and  $\text{mod}(\Delta^i \setminus \overline{\Pi^i})$  grow linearly. His statement implies ours after one notices that if  $n_i + 1 = n_{i+1}$  then  $\Delta^{i+1} = \Lambda_{n_i+1}$ , otherwise  $\Pi^i = \Lambda_{n_i+1}$ .

Those two results are proved in a slightly more general setting than we state here: they are valid for so-called full unfolded families of quadratic-like maps. This version allows to state results also for finitely renormalizable quadratic polynomials (via renormalization).

4.2.2. *General case.* The following more general theorem can be proved using the ideas of Theorem A of [L3] but it is a little bit tedious to check the details (it is necessary to get deep into the construction of [L2]).

**Theorem 4.5.** *For every  $K > 1$ ,  $T > 0$ , there exists  $T' > 0$  with the following properties. Let  $(\mathcal{R}_i, h_i)$  be a  $R$ -chain over  $\lambda_0$  and let  $n_k - 1$  be the sequence of non-central levels. If  $\text{Dil}(h_1|(U_1 \setminus \overline{U_1^0}) < K$  and  $\text{mod}(U_1[\lambda] \setminus \overline{U_1^0[\lambda]}) > T$  then  $\text{mod}(\Lambda_{n_k} \setminus \overline{\Lambda_{n_k+1}}) > T'k$ .*

Since we do not need the full strength of the previous theorem, we will state and prove a weaker estimate using an inductive argument based on the obvious estimate below:

**Lemma 4.6.** *There exists a constant  $C_0$  with the following property. Let  $h$  be a holomorphic motion over  $\Lambda$  and let  $\Lambda' \subset \Lambda$  be such that  $\nu = \text{mod}(\Lambda \setminus \overline{\Lambda'}) > C_0$ . Then the dilatation of  $h$  over  $\Lambda'$  is bounded by 2.*

**Theorem 4.7.** *For every  $K > 1$ , there exists constants  $T' > 0$ ,  $T'' > 0$  with the following properties. Let  $(\mathcal{R}_i, h_i)$  be a  $R$ -chain over  $\lambda_0$  and let  $n_k - 1$  be the sequence of non-central levels. If  $\text{Dil}(h_1|(U_1 \setminus \overline{U_1^0}) < K$  and  $\text{mod}(U_1[\lambda_0] \setminus \overline{U_1^0[\lambda_0]}) > T'$  then  $\text{mod}(\Lambda_{n_k} \setminus \overline{\Lambda_{n_k+1}}) > T''k$ .*

*Proof.* Let  $\nu_i = \text{mod}(\Lambda_i \setminus \overline{\Lambda_{i+1}})$ ,  $\mu_i = \text{mod}(U_i[\lambda_0] \setminus \overline{U_{i+1}[\lambda_0]})$ ,  $k_i = \text{Dil}(h_i|U_i[\lambda_0] \setminus \overline{U_{i+1}[\lambda_0]})$ . Notice that if  $\nu_i > C_0$  then  $k_{i+1} \leq 2$ , where  $C_0$  is the constant of Lemma 4.6. Moreover, for  $i > 1$ , and in particular for  $i = n_k$ , we have  $\nu_i = \text{mod}(\chi_i^0(U_i[\lambda_0] \setminus \overline{U_{i+1}[\lambda_0]})) \geq \mu_i/k_i$ .

Let  $T = 400(2 + C_0)(2 + K)$  and let  $T' > T$  be so big that if  $\mu_1 > T'$  then  $\mu_{n_k} > T$ ,  $k \geq 1$ . Let also  $T''$  be such that if  $\mu_1 > T'$  then  $\mu_{n_k} > kT''(2 + K)$ .

Let us assume that for some  $m$ , we have  $\mu_m > T$  and  $k_m \leq (2 + K)$ , and let  $m' \geq m$  be the next non-central return.

For  $\lambda \in \Lambda_{m'+1}$ , we have  $R_m^{m'-m+1}(0) \in W_m^{\underline{d}}$  for some  $\underline{d}$ . Let  $\Upsilon$  be the component of  $R_m(0)$  of  $(R_m^{m'-m}|U_{m'})^{-1}(U_m^{\underline{d}})$  and  $\Upsilon'$  be the component of  $R_m(0)$  of  $(R_m^{m'-m}|U_{m'})^{-1}(W_m^{\underline{d}})$ .

Let  $H_m = L(h_m)$  outside of  $U_m^{\underline{d}}$  and let the leaves of  $H_m|U_m^{\underline{d}}$  be the preimages by  $R_m^{\underline{d}}$  of the leaves of  $h_m$ . If  $m = m'$ , let  $H = H_m$ . Otherwise, notice that if  $\lambda \in \Lambda_{m'-1}$ , then  $R_m^{m'-m}|U_{m'-1}^0[\lambda]$  is a  $2^{m'-m}$  branched covering map over  $U_m[\lambda]$ , and for  $\lambda \in \Lambda_{m'}$ ,

$$R_m^{m'-m}|U_{m'}[\lambda] \setminus \overline{U_{m'}^0[\lambda]}$$

is unbranched. Let  $H$  be the lift of  $H_m$  by  $(R_m^{m'-m}|U_{m'-1}^0, U_{m'}^0, \Lambda_{m'})$ . So in both cases,  $H$  is a holomorphic motion over  $\Lambda_{m'}$ .

With this definition,  $\Upsilon$  and  $\Upsilon'$  (which are apriori defined over  $\Lambda_{m'+1}$ ) move holomorphically with  $H$  (over  $\Lambda_{m'}$ ).

Let  $\chi$  be the holonomy family of the pair  $(\mathcal{R}_{m'} \circ \mathbf{0}, H|\partial U_{m'} \cup \overline{\Upsilon})$ . It is clear that  $\text{Dil}(\chi|\Upsilon)$  is bounded by  $k_m$ . In particular, we can estimate  $\nu_{m'} \geq \text{mod}(\Upsilon[\lambda_0] \setminus \overline{\Upsilon'[\lambda_0]})/k_m = \mu_m/k_m \geq \mu_m/(2 + K) > C_0$ . With  $m = 1$ , we have  $k_1 \leq K \leq 2 + K$  by hypothesis and  $m' = n_1 - 1$ , so  $\nu_{n_1-1} \geq \mu_1/(2 + K) \geq T/(2 + K) \geq C_0$  and  $k_{n_1} \leq 2 \leq 2 + K$ . With  $m = n_k$ , we have that  $m' = n_{k+1} - 1$  and  $\nu_{n_{k+1}-1} \geq \mu_{n_k}/(2 + K) \geq T/(2 + K) \geq C_0$  and  $k_{n_{k+1}} \leq 2 \leq 2 + K$ , provided  $k_{n_k} \leq 2 + K$ . By induction, we have  $k_{n_k} \leq 2 + K$  for every  $k$ , so  $\nu_{n_k} \geq \mu_{n_k}/(2 + K) > T''k$ .  $\square$

This simple inductive argument does not seem to work easily to get the full Theorem 4.5.

## 5. UNIMODAL MAPS

We refer to the book of de Melo & van Strien [MS] for the general background in one-dimensional dynamics.

We will say that a smooth (at least  $C^2$ ) map  $f : I \rightarrow I$  of the interval  $I = [-1, 1]$  is *unimodal* if  $f(-1) = -1$ ,  $f(x) = f(-x)$  and 0 is the only critical point of  $f$  and is non-degenerate, so that  $D^2f(0) \neq 0$ .

*Remark 5.1.* The introduction of normalization and symmetry assumptions are in order to avoid cumbersome notations: all results and proofs generalize to the non-symmetric case. See also Appendix C of [ALM].

*Remark 5.2.* The assumption that the critical point is non-degenerate is made just for convenience: typical unimodal maps certainly have non-degenerate critical point. If one is not willing to make this assumption, one should add the non-degeneracy condition to the Kupka-Smale definition below. In this case it would still hold that in non-trivial analytic families parameters with a degenerate critical point have zero Lebesgue measure (and are contained in a countable number of analytic submanifolds with codimension at least 1), see Lemma 10.6.

The theory of unimodal maps with fixed non-quadratic criticality is considerably different and less complete than the typical case, and the proofs of this work do not apply

Let  $\mathbb{U}^k$ ,  $k \geq 2$  be the space of  $C^k$  unimodal maps. We endow  $\mathbb{U}^k$  with the  $C^k$  topology.

Basic examples of unimodal maps are given by quadratic maps

$$(5.1) \quad q_\tau : I \rightarrow I, \quad q_\tau(x) = \tau - 1 - \tau x^2,$$

where  $\tau \in [1/2, 2]$  is a real parameter.

A map  $f \in \mathbb{U}^2$  is said to be Kupka-Smale if all periodic orbits are hyperbolic. It is said to be hyperbolic if it is Kupka-Smale and the critical point is attracted to a periodic attractor. It is said to be regular if it is hyperbolic and its critical point is not periodic or preperiodic. It is well known that regular maps are structurally stable.

A  $k$ -parameter  $C^r$  (or analytic) family of unimodal maps is a  $C^r$  (or analytic) map  $F : \bar{\Lambda} \times I \rightarrow I$  such that  $f_\lambda \in \mathbb{U}^2$ , where  $f_\lambda(x) = F(\lambda, x)$  where  $\Lambda \in \mathbb{R}^k$  is a bounded open connected domain with smooth ( $C^\infty$ ) boundary. We denote  $\mathbb{UF}^r(\Lambda)$  the space of  $C^r$  families of unimodal maps, endowed with the  $C^r$  topology. Notice that  $\mathbb{UF}^r(\Lambda)$  is a separable Baire space.

We will not introduce a topology in the space of analytic families of unimodal maps.

**5.1. Combinatorics and hyperbolicity.** Let  $f \in \mathbb{U}^2$ . A symmetric interval  $T \subset I$  is said to be nice if the iterates of  $\partial T$  never return to  $\text{int } T$ . A nice interval  $T \neq I$  is said to be a restrictive (or periodic) interval of period  $m$  for  $f$  if  $f^m(T) \subset T$  and  $m$  is minimal with this property. In this case, the map  $A \circ f^m \circ A^{-1} : I \rightarrow I$  is again unimodal for some affine map  $A : T \rightarrow I$ : this map is usually called a renormalization of  $f$  if  $m > 1$  or a unimodal restriction if  $m = 1$ .

If  $T \subset I$  is a nice interval, the domain of the first return map  $R_T$  to  $T$  consists of a (at most) countable union of intervals which we denote  $T^j$ . We reserve the index 0 for the component of 0:  $0 \in T^0$ , if 0 returns to  $T$ . From the nice condition,  $R_T|_{T^j}$  is a diffeomorphism if  $0 \notin T^j$ , and is an even map if  $0 \in T^j$ . We call  $T^0$  the central domain of  $R_T$ . The return  $R_T$  is said to be central if  $R_T(0) \in T^0$ .

The following well known result shows that nice intervals allow to study arbitrarily small neighborhoods of 0.

**Lemma 5.1.** *Let  $f \in \mathbb{U}^2$  be non-regular Kupka-Smale and assume the critical orbit is infinite. Then for every  $\epsilon > 0$ , there exists a nice interval  $[-p, p] \subset T$  with  $p$  preperiodic.*

Reference

Under the Kupka-Smale condition, the dynamics outside a nice interval is hyperbolic, and in particular persistent:

**Lemma 5.2.** *Let  $f \in \mathbb{U}^2$  and let  $T \subset I$  be a nice interval. If all periodic orbits contained in  $I \setminus \text{int } T$  are hyperbolic (in particular if  $f$  is Kupka-Smale), then*

(1) *The set of points  $X \subset I$  which never enter  $\text{int } T$  splits in two forward invariant sets: an open set  $U$  attracted by a finite number of periodic orbits and a closed set  $K$  such that  $f|_K$  is uniformly expanding:  $|Df^n(x)| > C\lambda^n$ , for  $x \in K$  and for some constants  $C > 0$ ,  $\lambda > 1$ . Moreover, preperiodic points are dense in  $K$ .*

(2) *There exists a neighborhood  $\mathcal{V} \subset \mathbb{U}^2$  of  $f$  and a continuous family of homeomorphisms  $H[g] : I \rightarrow I$ ,  $g \in \mathcal{V}$  such that  $g \circ H[g]|_{I \setminus T} = h[g] \circ f$ , and  $h[f] = \text{id}$ .*

Reference

The following is an easy consequence of Lemma 5.2.

**Lemma 5.3.** *Let  $f_\lambda$ ,  $\lambda \in (-\epsilon, \epsilon)$  be a  $C^2$  family of unimodal maps, and let  $T$  be a nice interval with preperiodic boundary for  $f = f_{\lambda_0}$ . Assume that there exists an interval  $0 \in J$  and a family  $T[\lambda]$  of intervals with preperiodic boundary, such that  $T[0] = T$  and for  $\lambda \in J$ , all non-hyperbolic periodic orbits of  $f_\lambda$  intersect  $\text{int } T[\lambda]$ . Then there exists a continuous family of homeomorphisms  $H[\lambda] : I \rightarrow I$ ,  $\lambda \in J$  such that  $H[\lambda](T) = T[\lambda]$  and  $f_\lambda \circ H[\lambda](I \setminus T) = H[\lambda] \circ f$  and  $H[0] = \text{id}$ .*

**5.1.1. Principal nest.** We say that  $f$  is infinitely renormalizable if there exists arbitrarily small restrictive intervals  $T \subset I$ . Otherwise we say that  $f$  is finitely renormalizable.

Let  $\mathcal{F} \subset \mathbb{U}^2$  be the class of Kupka-Smale finitely renormalizable maps whose critical point is recurrent, but not periodic. If  $f \in \mathcal{F}$ , the first return map  $f^m : T \rightarrow T$  to its smallest restrictive interval has a orientation reversing fixed point which we call  $p$ . Let  $I_1 = [-p, p]$ . Define a nested sequence of intervals  $I_i$  as follows. Assuming  $I_i$  defined, let  $R_i$  be the first return map to  $I_i$  and let  $I_{i+1}$  be the central domain  $I_i^0$  of  $R_i$ .

The sequence  $I_i$  is called the *principal nest* of  $f$ . A level of the principal nest is called central if  $R_i$  is a central return.

**5.2. Negative Schwarzian derivative.** The Schwarzian derivative of a map  $C^3$  map  $f : I \rightarrow I$  is defined by

$$Sf = \frac{D^3 f}{Df} - \frac{3}{2} \left( \frac{D^2 f}{Df} \right)^2$$

in the complement of the critical points of  $f$ . If  $Sf$  and  $Sg$  are simultaneously positive (or negative) then  $S(g \circ f)$  is positive (or negative).

If  $f$  is a unimodal map the condition of negative Schwarzian derivative is very useful and can be exploited in several ways. Quadratic maps have negative Schwarzian derivative. Moreover, one can often reduce to this situation as is shown by the following well known estimate:

**Lemma 5.4.** *If  $f \in \mathbb{U}^3$  is infinitely renormalizable, then if  $T \subset I$  is a small enough periodic nice interval, the first return map to  $T$  has negative Schwarzian derivative.*

Recently, Kozłowski showed that the assumption of negative Schwarzian can be often removed. The next result follows from Lemma 5.2 and [GSS] (which is based on the work of Kozłowski [K1]).

**Lemma 5.5.** *Let  $f \in \mathcal{F} \cap \mathbb{U}^3$ . There exists  $i > 0$ , an analytic diffeomorphism  $s : I \rightarrow I$  and a neighborhood  $\mathcal{V} \subset \mathbb{U}^3$  of  $f$ , such that there exists a continuation  $I_i[g]$ ,  $g \in \mathcal{V}$  of  $I_i$  ( $H[g](I_i) = I_i[g]$  in the notation of Lemma 5.2) such that the first return map to  $s(I_i[g])$  by  $s \circ g \circ s^{-1} : I \rightarrow I$  has negative Schwarzian derivative.*

**5.3. Decay of geometry.** The following result is due to Lyubich in the case of negative Schwarzian derivative and holds in general due to the work of Kozłowski:

**Lemma 5.6.** *Let  $f \in \mathcal{F}$  be at least  $C^3$ , and let  $n_k - 1$  denote the sequence of non-central levels in the principal nest of  $f$ . Then  $|I_{n_k+1}|/|I_{n_k}| < C\lambda^k$  for some constants  $C > 0$ ,  $\lambda < 1$ .*

**5.4. Quasiquadratic maps.** A map  $f \in \mathbb{U}^3$  is *quasiquadratic* if any nearby map  $g \in \mathbb{U}^3$  is topologically conjugate to some quadratic map. By the theory of Milnor-Thurston and Guckenheimer [MS], a map  $f \in \mathbb{U}^3$  with negative Schwarzian derivative and  $Df(-1) \geq 1$  is quasiquadratic, so the quadratic maps are quasiquadratic. The following result gives conditions for a unimodal map to be quasiquadratic:

**Theorem 5.7** (see Lemma 2.13 of [ALM]). *Let  $f \in \mathbb{U}^3$  be a Kupka-Smale unimodal map which is topologically conjugate to a quadratic map. Then  $f$  is quasiquadratic.*

*Remark 5.3.* The previous result is the reason that the quasiquadratic condition considers only  $C^3$  maps and the  $C^3$  topology (otherwise it would not be possible to guarantee that even quadratic maps are quasiquadratic).

**Theorem 5.8** (see Remark 2.4 of [ALM]). *Let  $f \in \mathbb{U}^3$ . If  $f$  is not conjugate to a quadratic polynomial then there exists a (not necessarily hyperbolic) periodic orbit which attracts an open set. In particular, if all periodic orbits of  $f$  are repelling then  $f$  is conjugate to a quadratic polynomial.*

## 6. CONSTRUCTION OF THE SPECIAL FAMILY

**6.1. Puzzle maps.** Let  $f \in \mathbb{U}_a$  be a finitely renormalizable unimodal map with a recurrent critical point. Let us consider some nice interval  $A^0$  and let  $\{A^j\}$  be the connected components of the domain of the first landing map from  $I$  to  $A^0$ . We call the family  $\{A^j\}$  the real puzzle for  $f$  associated to  $A^0$ . The basic object used in [ALM] to analyze the dynamics of unimodal maps can be viewed as a complexification of such real puzzles, which are called simply a puzzle.

The definition of puzzle in [ALM] is too general and technical for our purposes. In this paper, we will simply describe how to construct a puzzle for  $f$  (or rather a geometric puzzle, in the language of [ALM]). Instead of giving the precise definitions of a puzzle, we will just obtain the properties that are needed for our results.

Let us fix some advanced level  $\mathbf{n}$  of the principal nest of  $f$  and assume that  $|I_{\mathbf{n}}|/|I_{\mathbf{n}-1}|$  is very small. Let us fix the following notation: let  $A^0 = I_{\mathbf{n}}$  and let  $\{A^j\}$  be the real puzzle associated to  $A^0$ . We let  $A^1$  be such that  $f(0) \in A^1$ .

Given  $0 < \theta \leq \pi/2$ , and  $A \subset \mathbb{R}$ , let  $D_{\theta}(A)$  be the intersection of two round disks  $D_1$  and  $D_2$  where  $D_1 \cap \mathbb{R} = A$ ,  $\partial D_1$  intersects  $\mathbb{R}$  making an angle  $\theta$ , and  $D_2$  is the image of  $D_1$  by symmetry about  $\mathbb{R}$ . The complexification of the real puzzle  $\{A^j\}$  should be imagined as  $\{D_{\theta}(A^j)\}$  for a suitable value of  $\theta$ . Of course, since the system is non-linear, the definition can not be so simple. Nevertheless, the condition  $|I_{\mathbf{n}}|/|I_{\mathbf{n}-1}|$  small allows to bound the nonlinearity of the first landing map to  $I_{\mathbf{n}}$  and we can obtain:

**Lemma 6.1.** *Let  $0 < \phi < \psi < \gamma < \pi/2$  be fixed. For arbitrarily big  $k > 0$ , if  $|I_{\mathbf{n}}|/|I_{\mathbf{n}-1}|$  is small enough, there exists a sequence  $V^j$  of open Jordan disks such that  $D_{\phi}(A^j) \subset V^j \subset D_{\psi}(A^j)$  and  $V^0 = D_{(\phi+\psi)/2}(A^0)$  with the following properties:*

- (1) *If  $j \neq 0$  and  $f(A^j) \subset A^k$  then  $f : V^j \rightarrow V^k$  is a diffeomorphism;*
- (2) *If  $f(A^0) \cap A^j \neq \emptyset$ , then  $\text{mod } f(V^0) \setminus \overline{D_{\gamma}(A^j)} > k$ .*

**6.2. A special Banach space of perturbations.** Let  $A^1 = [l, r]$  with  $l < r$ , and let  $N = [-l, l]$ . Domains  $V^j$  which do not intersect  $A^1$  or  $N$  will play no role in the construction to follow. Let  $V$  be the union of all  $V^j$  such that  $A^j \subset N \cup A^1$ .

One of the main problems of [ALM] is to obtain a direction  $v$  (or infinitesimal perturbation) which is transverse to the topological class of  $f$ . The idea is to

consider a perturbation which does not affect much  $f$  in  $N$ , but causes a bump near the critical value, localized in  $A^1$ . There are several difficulties related to this scheme, the first of which is that such a bump can only be reasonably controlled up to its first derivative. Another difficulty is that we want an analytic perturbation, so it cannot vanish in  $N$  and be a bump at  $A^1$ . The solution involves the consideration of a certain Banach spaces of smooth ( $C^1$ ) functions in  $N \cup A^1$  which are analytic in  $\text{int } N \cup \text{int } A^1$ , which allows to construct perturbations that, while badly behaved in the real line (can be only controlled up to the first derivative), are well behaved with respect to the complex puzzle structure.

While the proof in [ALM] involves two steps, construction of a transverse smooth vector field and approximation of this vector field by polynomials, which need two different Banach spaces, we will realize the same construction with just one Banach space. This is important to estimate the asymmetric roles of perturbations concentrated in  $N$  and  $A^1$ . The proof of our main perturbation estimate (Lemma 6.4) is a mixture of two estimates of [ALM], Lemmas 7.4 (for perturbations localized in  $A^1$ ) and 7.9 (for perturbations supported on  $N \cup A^1$ ) of that paper.

Let  $Z = D_\gamma(A^1) \cup D_\gamma(N)$ , and let  $\Upsilon$  be the space of all vector fields  $v$  holomorphic on  $Z$  and whose derivative admits a continuous extension to  $\overline{Z}$ , which vanish up to the first derivative in  $\partial A^1$  and its forward iterates (this is a finite set) and such that  $v|_{D_\gamma(N)}$  is a symmetric (odd) vector field. We use the norm  $\|v\| = \sup_{\{z \in Z\}} |Dv|$ .

Let  $\Upsilon = \Upsilon_1 \oplus \Upsilon_2$ , where  $v \in \Upsilon_1$  if  $v|_{D_\gamma(N)} = 0$  and  $v \in \Upsilon_2$  if  $v|_{D_\gamma(A^1)} = 0$ .

The reader should think of vector fields  $v \in \Upsilon$  as acting as perturbations on  $f$  by  $v \rightarrow f \circ (\text{id} + v)$ . Let  $f_v = f \circ (\text{id} + v)$ . One of the main advantages of the definition of  $\Upsilon$  is that, for small  $v \in \Upsilon$ , “the puzzle persists”, that is, there exists a continuation  $V^v$  of the set  $V$  inside  $Z$ , whose connected components behave, under iteration by  $f_v$ , in the same way that the connected components of  $V$  behaved under iteration by  $f$ .

To make this more precise, let us say that  $v \in \Upsilon$  is admissible if there exists a holomorphic motion  $h^v$  over  $\mathbb{D}$ , defined by the family of transition maps  $h[0, \lambda] \equiv h_\lambda^v : \mathbb{C} \rightarrow \mathbb{C}$ ,  $\lambda \in \mathbb{D}$  such that:

- (1)  $h_\lambda^v|_{\mathbb{C} \setminus Z} = \text{id}$ ,  $h_\lambda^v|_{\partial f(V^0)} = \text{id}$ ;
- (2)  $f_{\lambda v} \circ h_\lambda^v|_{V \setminus V^0} = h_\lambda \circ f$ ,  $f_{\lambda v} \circ h_\lambda^v|_{\partial V^0} = f$ .

The holomorphic motion  $h^v$  will be said to be *compatible* with  $v$ .

The following is a restatement of Lemma 7.9 of [ALM].

**Lemma 6.2.** *There exists  $\epsilon > 0$  such that if  $v$  belongs to  $\{v \in \Upsilon \mid \|v\| < \epsilon\}$  then  $v$  is admissible.*

We also need the following simple estimate (see the proof of Lemma 7.4 of [ALM]):

**Lemma 6.3.** *Let  $0 < \theta < \gamma < \pi/2$ . There exists  $\epsilon' > 0$  such that if  $A$  is an interval and  $v$  is holomorphic on  $D_\gamma(A)$  whose derivative extends continuously to  $\overline{D_\gamma(A)}$  satisfying  $|Dv| < \epsilon'$  then  $\text{id} + v : D_\gamma(A) \rightarrow \mathbb{C}$  is a diffeomorphism and  $D_\theta(A) \subset (\text{id} + v)(D_\gamma(A))$ .*

Now we can prove:

**Lemma 6.4.** *There exists constants  $\epsilon_1 > 0$ ,  $\epsilon_2 > 0$ , where  $\epsilon_1$  depends only on  $\psi$  and  $\gamma$  such that if  $v$  belongs to  $\{v \in \Upsilon_1 \mid \|v\| < \epsilon_1\} \times \{v \in \Upsilon_2 \mid \|v\| < \epsilon_2\}$  then  $v$  is admissible.*

*Proof.* Let  $n_1$  be such that  $f^{n_1}(V^1) = V^0$  and let  $\theta = (\psi + \gamma)/2$ .

Let  $v \in \{v \in \Upsilon \mid \|v\| < \epsilon\}$ . By Lemma 6.2, there exists a holomorphic motion  $h^v$  compatible with  $v$ . We claim that if  $0 < \epsilon_2 < \epsilon$  and  $\|v\| < \epsilon_2$  then for  $\lambda \in \mathbb{D}$ ,  $h_\lambda^v(V^1) \subset D_\theta(A^1)$ .

Indeed, if this is not the case, there would be a sequence  $z_k \in \partial D_\theta(A^1)$ ,  $v_k \in \Upsilon$ ,  $v_k \rightarrow 0$ , such that  $f_{v_k}^{n_1+1}(z_k) \in f(V^0)$ . It clearly follows that  $z_k \rightarrow \partial A^1 = \{l, r\}$ , let us say that  $z_k \rightarrow l$ . It is clear that

$$f_{v_k}^{n_1+1}(z_k) = f_{v_k}^{n_1+1}(l) + Df_{v_k}^{n_1+1}(l)z_k + o(z_k) = f^{n_1+1}(l) + Df^{n_1+1}(l)z_k + o(z_k).$$

In particular, the sequence  $f_{v_k}^{n_1+1}(z_k)$  converges to  $f^{n_1+1}(l)$  along a direction which makes angle  $\theta$  with the real line (since  $Df^{n_1+1}(l) \in \mathbb{R} \setminus \{0\}$ ), so  $f_{v_k}^{n_1+1}(z_k) \notin f(V^0)$  for  $k$  big, which is a contradiction.

Let  $\epsilon_1$  be as in Lemma 6.3. If  $v = v_1 + v_2$ , with  $v_i \in \Upsilon_i$  and  $\|v_i\| < \epsilon_i$ , let  $h_\lambda^v : \mathbb{C} \setminus (D_\gamma(A^1) \setminus V^1)$  be given by  $h_\lambda^v|(\mathbb{C} \setminus D_\gamma(A^1)) = h_\lambda^{v_2}$  and  $h_\lambda^v|V^1 = ((\text{id} + \lambda v_1)|D_\gamma(A^1))^{-1} \circ h_\lambda^{v_1}$ . Any extension of  $h_\lambda^v$  to  $\mathbb{C}$  is clearly compatible with  $v$ .  $\square$

We will also need the following easy lemma:

**Lemma 6.5.** *If  $|I_n|/|I_{n-1}|$  is sufficiently small, then for  $w$  in  $\{w \in \Upsilon_1 \mid \|w\| < \epsilon_1\} \times \{w \in \Upsilon_2 \mid \|w\| < \epsilon_2\}$  and  $\lambda \in \mathbb{D}$ , then  $(f_{\lambda w}|h_\lambda^w(V^0))^{-1}(D_\gamma(V^1)) \subset \mathbb{D}_{\rho|A^0|}(0)$ , where  $\rho \rightarrow 0$  as  $|I_n|/|I_{n-1}| \rightarrow 0$ .*

*Proof.* Let  $U = h_\lambda(V^0)$  and  $U^0 = (f_{\lambda w}|W)^{-1}(D_\gamma(A^1))$ . Notice that  $f_{\lambda w}(0) = f(0) \in D_\gamma(V^1)$ . Thus,  $f_{\lambda w}|(U \setminus \overline{U^0})$  is a double covering of  $f(U_0) \setminus \overline{D_\gamma(A^1)}$ . By Lemma 6.1, if  $|I_n|/|I_{n-1}|$  is small then  $\text{mod}(f(U_0) \setminus \overline{D_\gamma(A^1)})$  is large, and so  $\text{mod}(U \setminus \overline{U^0})$  is also big. Since the derivative of  $\text{id} + \lambda w$  is smaller than  $\max\{1 + \epsilon_1, 1 + \epsilon_2\}$ , we see that the diameter of  $U$  is at most  $2|A^0|$ , so the diameter of  $U^0$  can be bounded by  $\rho|A^0|/2$  with small  $\rho$  as required.  $\square$

**6.3. Analytic vector fields.** We will be specially concerned with special types of  $w$  which generate analytic families of unimodal maps. The following lemma is obvious:

**Lemma 6.6.** *If  $w$  has an analytic extension  $w : I \rightarrow I$  of  $C^1$  of norm less than one, such that  $w(-1) = w(1) = 0$ , then  $f_{\lambda w}$ ,  $\lambda \in (-1, 1)$  is an analytic family of unimodal maps.*

The following is a consequence of the Mergelyan Polynomial Approximation theorem:

**Lemma 6.7.** *Let  $w \in \Upsilon$ . Then there exists a sequence  $w_m \in \Upsilon$  such that the  $C^1$  norm of  $w_m|I$  is less than or equal to  $\|w\|$ ,  $w_m(-1) = w_m(1) = 0$  and  $w_m \rightarrow w$  in  $\Upsilon$ . If  $w$  is real-symmetric then we can also choose  $w$  real-symmetric.*

The proof of the following lemma is obvious using Lemma 5.2.

**Lemma 6.8.** *If  $w$  belongs to  $\{w \in \Upsilon_1 \mid \|w\| < \epsilon_1\} \times \{w \in \Upsilon_2 \mid \|w\| < \epsilon_2\}$  and has an analytic extension to  $I$  with  $w(-1) = w(1) = 0$ , the following holds. For  $\epsilon > 0$  small and  $\lambda \in (-\epsilon, \epsilon)$ ,*

- (1) *the boundary points of  $I_n$  are preperiodic points for  $f_{\lambda w}$ ,*
- (2) *there exists homeomorphisms  $H[\lambda]$ , depending continuously on  $\lambda$  such that  $f_{\lambda w} \circ H[\lambda]|I \setminus I_n = H[\lambda] \circ f$ ,*



- (3)  $H[\lambda](I_n) = I_n$ ,
- (4) *the connected components of  $((\text{id} + \lambda w)|h_\lambda^w(V^0))^{-1}(\cup V^j)$  intersect the real line at the connected components of the domain of the first return map to  $I_n$  under iteration by  $f_{\lambda w}$ .*

**6.4. A special perturbation.** Let us consider an affine map  $Q : A^1 \rightarrow I$ , and let

$$\tilde{v}_n(z) = (1 - z^2)(1 - e^{-2n}) + \frac{2}{n}(e^{-n(1+z)} + e^{-n(1-z)} - e^{-2n} - 1)$$

and let  $v_n \in \Upsilon_1$  be such that  $v_n|_{D_\gamma(A^1)} = Q^*\tilde{v}_n\epsilon_1/8$ . Notice that  $\|v_n\| < \epsilon_1$ .

**6.4.1. Infinitesimal transversality.** The importance of the sequence  $v_m$  in [ALM] is that it is eventually transverse to the topological class of  $f$  in a certain sense which we will precise later.

A quasiconformal vector field  $\alpha$  of  $\overline{\mathbb{C}}$  is a continuous vector field with locally integrable distributional derivatives  $\overline{\partial}\alpha$  and  $\partial\alpha$  in  $L^1$  and  $\overline{\partial}\alpha \in L^\infty$ .

Let us say that  $w$  is *formally transverse at  $f$*  if there is no quasiconformal vector field  $\alpha$  of  $\mathbb{C}$ , such that  $w(z) = f^*\alpha - \alpha$ ,  $z \in \text{orb}_f(0)$ .

The following summarizes Lemmas 7.6, 7.7 and 7.8 of [ALM].

**Lemma 6.9.** *Let  $v_m$  be defined as above. If  $|I_n|/|I_{n-1}|$  is sufficiently small, then for  $m$  sufficiently big,  $v_m$  is formally transverse at  $f$ .*

The following is due to (a version of) the so-called Key estimate of [ALM] (more precisely we use Corollary 7.14 of [ALM]):

**Lemma 6.10.** *The set of vector fields  $w \in \Upsilon$  which are not formally transverse at  $f$  is a closed subspace of  $\Upsilon$ .*

In particular, if  $m$  is sufficiently big and  $w$  is close to  $v_m$  then  $w$  is formally transverse at  $f$ .

**6.4.2. Macroscopic transversality.** The following result can be interpreted as the macroscopic counterpart to the infinitesimal transversality of  $v_m$ .

Let  $r > 0$  be minimal with  $f^{r+1}(0) \in V^1$ .

**Lemma 6.11.** *There exists a constant  $\tau_0 > 0$  depending only on  $\epsilon_1$  and  $\phi$ , such that if  $|I_n|/|I_{n-1}|$  is sufficiently small the following holds. Let  $v_m$  be defined as above and let  $r > 0$  be minimal with  $f^{r+1}(0) \in V^1$ . Then for  $m$  sufficiently big, there exists a domain  $\hat{\Theta} \subset \mathbb{D}$  such that the map  $\theta : \hat{\Theta} \rightarrow \mathbb{C}$  given by  $\theta(\lambda) = f_\lambda^r(0)$  is a diffeomorphism onto  $\mathbb{D}_{\tau_0|I_n|}$ .*

*Proof.* Since  $\|v_m\| < \epsilon_1$ , there exists a holomorphic motion  $h^{v_m}$  which is compatible with  $v_m$ .

Let  $\Psi : \mathbb{D} \rightarrow \mathbb{C}$ ,  $\Psi(\lambda) = (\text{id} + \lambda v_m)(f(0))$ . It is clearly a diffeomorphism over a round disk  $D_m$  centered on 0. Let  $d_m$  be the hyperbolic distance between  $f(0)$  and  $\partial D_m$  in  $D_{\pi/2}(A^1)$ . It is easy to estimate directly  $d_m$  by below in terms of  $\epsilon_1$  and  $m$ . In particular, for  $m$  big,  $d_m > \tilde{\tau} > 0$  where  $\tilde{\tau}$  depends only on  $\epsilon_1$ , not on the position of  $f(0)$  in  $A^1$ .

(To see this, notice that  $D\Psi(0) = v_m(f(0))$ , and the norm of  $v_m(f(0))$  in the hyperbolic metric of  $D_{\pi/2}(A^1)$  at  $f(0)$  is at least  $\epsilon_1/10$  for  $m$  big. Let  $P : D_{\pi/2}(A^1) \rightarrow \mathbb{D}$  be a Moebius transformation taking  $f(0)$  to 0. The norm of  $D(P \circ \Psi)(0)$  in the hyperbolic metric of  $\mathbb{D}$  at 0 is at least  $\epsilon_1/10$ , so the Euclidean norm of  $D(P \circ \Psi)(0)$  is at least  $\epsilon_1/10$ . By the Koebe 1/4 Theorem,  $P(D_m)$  contains a round disk of

radius  $\epsilon_1/40$  around 0, thus the hyperbolic distance from  $\partial P(D_m)$  to 0 in  $\mathbb{D}$  is at least  $\epsilon_1/40$ .)

Now let  $Q$  be the connected component of  $f(0)$  on  $f^{-(r-1)}(V^0)$ , so that  $f^{r-1} : Q \rightarrow V^0$  is a diffeomorphism. The hyperbolic distance between  $\partial D \cap Q$  and  $f(0)$  in  $Q$  is bounded from below by  $\tilde{\tau}$  by the Schwarz Lemma (if  $\partial D \cap Q = \emptyset$ , we let this distance be  $\infty$ ). It follows that  $f^{r-1}(Q \cap D)$  contains a  $\tilde{\tau}$  hyperbolic neighborhood of  $f^r(0)$  on  $V^0$ . Now, if  $|I_n|/|I_{n-1}|$  is very small, then  $|I_{n+1}|/|I_n|$  is also very small, so  $f^r(0)$  (which is contained in  $I_{n+1}$ ) is  $\tilde{\tau}/2$  close to 0 in the hyperbolic metric of  $V^0 \supset D_\phi(A^0)$ .

As a consequence,  $f^{r-1}(Q \cap D)$  contains a  $\tilde{\tau}/2$  hyperbolic neighborhood of 0 in  $V^0$ , and since  $V^0 \supset D_\phi(A^0)$ , it must contain  $\mathbb{D}_{\tau|A^0|}$ , where  $\tau$  depends on  $\epsilon_1$  and  $\phi$ .  $\square$

**6.4.3. Construction of a full  $R$ -family.** Let  $\tau_0$  be the constant of Lemma 6.11 and let  $|I_n|/|I_{n-1}|$  be such that Lemma 6.5 holds with  $\rho < \tau_0/4$ .

Let  $m$  be big and let us fix  $v = v_m$  such that Lemmas 6.11 and 6.9 are valid, and let  $\hat{\Theta}$  be the domain of Lemma 6.11.

Let  $w \in \{w \in \Upsilon_1 \mid \|w\| < \epsilon_1\} \times \{w \in \Upsilon_2 \mid \|w\| < \epsilon_2\}$ .

Let  $U[0] = V^0$  and let the family  $\{U^j[0]\}$  denote the connected components of  $(f|V^0)^{-1}(\cup V^j)$ , letting  $0 \in U^0[0]$ .

Let us consider a holomorphic motion  $\tilde{H}$  over  $\mathbb{D}$  given by the transition maps  $\tilde{H}[0, \lambda] = H_\lambda : \mathbb{C} \rightarrow \mathbb{C}$  such that:

$$\tilde{H}_\lambda | \mathbb{C} \setminus U[0] = h_{\lambda w}$$

$$f_{\lambda w} \circ \tilde{H}_\lambda | U[0] \setminus U^0[0] = h_{\lambda w} \circ f.$$

Let  $U[\lambda] = \tilde{H}_\lambda(U[0])$ ,  $U^j[\lambda] = \tilde{H}_\lambda(U^j[0])$ .

Let  $R[\lambda]$  be the first return map from  $U^j[\lambda]$  to  $U_0$ . It is clear that  $(R[\lambda], H_\lambda)$  has a structure of a (non-full)  $R$ -family over  $\mathbb{D}$ . Let us consider the landing family  $(L[\lambda], H_\lambda)$  associated to  $(R[\lambda], \tilde{H}_\lambda)$ .

Let  $W^\mathcal{L}[0]$  be the domain of  $L[0]$  containing  $R[0](0)$ . Notice that  $L[\lambda]|W^\mathcal{L}[\lambda]$  extends to a diffeomorphism  $R^\mathcal{L}[\lambda]$  onto  $U[\lambda]$ . For  $\tau < \tau_0$ , let  $\Delta_\tau[\lambda]$  be the preimage of  $\mathbb{D}_{\tau|A^0|}(0)$  by this diffeomorphism.

If  $w = v$  then  $R^\mathcal{L}[\lambda] = R^\mathcal{L}[0]$  for all  $\lambda$ , since  $v$  is supported on  $D_\gamma(A^1)$ .

In particular,  $R^\mathcal{L}[\lambda] = R^\mathcal{L}[0]$  and  $\Delta_\tau[\lambda] = \Delta_\tau[0]$  for all  $\lambda$ . So  $\lambda \mapsto R[\lambda](0)$  is a map which restricts (in some domain  $0 \in \mathbb{O}^v$ ) to a diffeomorphism onto  $\Delta_\tau[0]$ . It follows that taking  $\tau = \tau_0/2$ , there exists a domain  $0 \in \mathbb{O}^w$  where  $\lambda \mapsto R[\lambda](0)$  is a diffeomorphism onto  $\Delta_\tau[0]$ , for any  $w$  close to  $v$  (of course,  $\mathbb{O}^w$  depends on  $w$ ).

But for  $w \in \Upsilon$  close to  $v$ , for all  $\lambda \in \mathbb{D}$ ,  $U^0[\lambda]$  is contained in  $D_{\rho|A^0|}$ , so  $W^\mathcal{L}[\lambda]$  is contained in  $\Delta_{\tau_0/2}[0]$  with space, for all  $\lambda \in \mathbb{D}$ . By the argument principle, letting  $\Theta$  be the connected component of 0 of the set of  $\lambda \in \hat{\Theta}$  with  $R[\lambda](0) \in W^\mathcal{L}[\lambda]$ , the map  $S : \overline{\Theta} \rightarrow \overline{W^\mathcal{L}[0]}$  such that  $S(\lambda) = H_\lambda^{-1}(R[\lambda](0))$  is a homeomorphism. We also have that the diameter of  $\Theta$  is very small if  $\rho$  is small (in particular if  $|I_n|/|I_{n-1}|$  is small).

Let  $U_1[0] = U^0[0]$  and let  $\{U_1^j[0]\}$  be the connected components of the preimage by  $R[0]|U^0[0]$  of  $\cup W^\mathcal{L}[0]$ , and let  $0 \in U_1^0$ .

Let  $h$  be a holomorphic motion over  $\Theta$  given by transition maps  $h[0, \lambda] = h_\lambda : \mathbb{C} \rightarrow \mathbb{C}$  such that

$$h_\lambda | \mathbb{C} \setminus U_1 = H_\lambda,$$

$$R[\lambda] \circ H_\lambda|_{U_1 \setminus U_1^0} = H_\lambda \circ R[0].$$

Let  $U_1[\lambda] = h_\lambda(U_1[0])$  and  $U_1^j[\lambda] = h_\lambda(U_1^j[0])$ .

Our construction shows clearly that the first return map  $R_1[\lambda]$  from  $\cup U_1^j[\lambda]$  to  $U_1[\lambda]$  is a  $R$ -map for  $\lambda \in \Theta$ , and so  $(R_1[\lambda], h_\lambda)$  is a  $R$ -family.

Our choice of  $\Theta$  implies that  $R_1[\lambda]$  is a full  $R$ -family.

Let us summarize the construction:

**Lemma 6.12.** *If  $|I_n|/|I_{n-1}|$  is small enough, there exists a real-symmetric vector field  $v \in \Upsilon$  and a neighborhood  $v \in \mathcal{V} \subset \Upsilon$  such that for any  $w \in \mathcal{V}$  real-symmetric there exists a real-symmetric domain  $0 \in \Theta$  and domains  $U_1[\lambda]$ ,  $U_1^j[\lambda]$ ,  $\lambda \in \Theta$  and a real-symmetric holomorphic motion  $h$  over  $\Theta$  defined by the transition maps  $h[0, \lambda] \equiv h_\lambda : \mathbb{C} \rightarrow \mathbb{C}$  such that*

$$(1) \quad h_\lambda(U_1[0]) = U_1[\lambda], \quad h_\lambda(U_1^j[0]) = U_1^j[\lambda];$$

$$(2) \quad \text{For } \lambda \in \Theta \cap \mathbb{R}, \quad h_\lambda(I_n) = I_n;$$

(3) *The first return map from  $\cup U_1^j[\lambda]$  to  $U_1[\lambda]$  under iteration by  $f_{\lambda w}$  is a  $R$ -map  $R_1[\lambda]$ ;*

$$(4) \quad (R_1[\lambda], h_\lambda) \text{ form a full } R\text{-family.}$$

And moreover, if  $w$  has an analytic extension to  $I$  such that  $w(-1) = w(1) = 0$  then:

(5) *For  $\lambda \in \Theta \cap \mathbb{R}$ ,  $I_{n+1}[\lambda] \equiv U_1[\lambda] \cap \mathbb{R}$  is the component of  $0$  of the first return map to  $I_n$  under iteration by  $f_{\lambda w}$ ;*

(6) *For  $\lambda \in \Theta \cap \mathbb{R}$ ,  $I_{n+1}^j[\lambda] \equiv U_1^j[\lambda] \cap \mathbb{R}$  are the domains of the first return map to  $I_{n+1}[\lambda]$ .*

The construction of the  $R$ -family gives us also a good control of its geometry.

**Lemma 6.13.** *In the setting of Lemma 6.12,  $\text{Dil}(h_\lambda|\mathbb{C} \setminus \overline{U_1^0[0]}) < 1 + \epsilon$ , and  $\text{mod}(U_1[0] \setminus \overline{U_1^0[0]}) > C$ , where  $\epsilon \rightarrow 0$  and  $C \rightarrow \infty$  if  $|I_n|/|I_{n-1}| \rightarrow 0$ .*

*Proof.* Indeed,  $\text{Dil}(h_\lambda|\mathbb{C} \setminus \overline{U_1^0[0]}) < 1 + \epsilon$  is bounded by the hyperbolic diameter of  $\Theta$  on  $\mathbb{D}$ , which is small if  $|I_n|/|I_{n-1}| \rightarrow 0$  is big. On the other hand,  $\text{mod}(U_1[0] \setminus \overline{U_1^0[0]}) \geq \text{mod}(U[0] \setminus \overline{U^0[0]})/2 \geq \text{mod}(f(V^0) \setminus \overline{V^1})/4 > k/4$ , which is big if  $I_n \setminus I_{n-1}$  is small by Lemma 6.1.  $\square$

**6.5. Remarks on the infinitesimal transversality of the special perturbation.** We would like to point out that the “macroscopic transversality” of  $v_m$  is very much related to its infinitesimal transversality. The argument is as follows:

(1)  $v_m$  can be  $C^1$  extended to  $I$  as an odd vector field which vanishes on  $[r, 1]$ ,  $[-1, -r]$  and  $[-l, l]$ . This vector field is not  $C^2$  but its  $C^1$  norm is small ( $\epsilon_1$ ).

(2) (Macroscopic transversality implies a  $C^1$  connecting lemma) Notice that the interval  $(f_{v_m}^r(0), f_{v_m}^r(0))$  contains the interval  $I_{n+1}$  (with lots of space). We conclude that the family  $f + \lambda v_m$ ,  $\lambda \in (-1, 1)$  must go through a parameter  $\lambda$  where  $f_{\lambda v_m}^r(0) = 0$ , and so changes the combinatorics of  $f$ .

(3) Using the Key Estimate of [ALM], we see that if  $v_m$  is not formally transverse at  $f$  then it is actually tangent to the topological class of  $f$  in the following sense. There exists a (real-symmetric) holomorphic motion  $h$  over  $\mathbb{D}$  whose transition maps  $h[0, \lambda] \equiv h_\lambda : \mathbb{C} \rightarrow \mathbb{C}$  are such that  $f_\lambda = h_\lambda \circ f \circ h_\lambda^{-1}$  is a family of so-called “puzzle maps” (which behave as unimodal maps) such that

$$\frac{d}{d\lambda} f_\lambda|_{\lambda=0} = \frac{d}{d\lambda} f_{\lambda v_m}|_{\lambda=0} = Df v_m$$

(the maps  $h_\lambda$  are characterized by  $\bar{\partial}h_\lambda/\partial h_\lambda = \lambda\partial\alpha$  for a specially chosen quasiconformal vector field  $\alpha$  satisfying  $v_m = f^*\alpha - \alpha$  on the critical orbit). (This family can be considered the Beltrami path through  $f$  in the direction of  $v$ .)

(4) The family  $f_\lambda$  is tangent to  $f_{\lambda v_m}$  at  $\lambda = 0$  and both families have big extensions (to  $\mathbb{D}$ ). In particular, they must be close together in a slightly smaller disk, where we can detect the change of combinatorics: there is a parameter  $\lambda \in \mathbb{D}$  such that  $f_\lambda^r(0) = 0^5$ .

(5) In particular, the family  $f_\lambda$  must change combinatorics, but this is a contradiction, since it consists of maps topologically conjugate to  $f$ . So we conclude that  $v_m$  is formally transverse at  $f$ . Notice that our argument is that a “reasonably efficient<sup>6</sup>” tangent path to  $v_m$  closes macroscopically the critical orbit.

(6) (Infinitesimal analytic connecting lemma) Although  $v_m$  is only  $C^1$  in the interval, we can approximate it in the topology of  $\Upsilon$  by polynomials  $w$  which will be still formally transverse to  $f$ . Those vector fields  $w$  are transversal to the topological class of  $f$ : they close “infinitesimally” the critical orbit.

## 7. PHASE-PARAMETER RELATION FOR THE SPECIAL FAMILY

Let  $f \in \mathcal{F}$  and let  $R_i : \cup I_i^j \rightarrow I_i$  be the first return map. For  $\underline{d} \in \Omega$ ,  $\underline{d} = (j_1, \dots, j_m)$ , let  $I_i^{\underline{d}} = \{x \in I_i \mid R_i^k(x) \in I_i^{j_{k+1}}, 0 \leq k < m\}$ , and let  $R_i^{\underline{d}} = R_i^m|_{I_i^{\underline{d}}}$ . Let  $C_i^{\underline{d}} = (R_i^{\underline{d}})^{-1}(I_i^0)$ . The map  $L_i : \cup C_i^{\underline{d}} \rightarrow I_i^0$  is the first landing map from  $I_i$  to  $I_{i+1}$ .

**Definition 7.1.** Let us say that a family  $f_\lambda$  of unimodal maps satisfies the Topological Phase-Parameter relation at a parameter  $\lambda_0$  if  $f = f_{\lambda_0} \in \mathcal{F}$ , and there exists  $i_0 > 0$  and a sequence of nested intervals  $J_i$ ,  $i \geq i_0$  such that:

- (1)  $J_i$  is the maximal interval containing 0 such that for all  $\lambda \in J_i$  there exists a homeomorphism  $H_i[\lambda] : I \rightarrow I$  such that  $f_\lambda \circ H_i[\lambda](I \setminus I_{i+1}) = H_i[\lambda] \circ f$ .
- (2) There exists a homeomorphism  $\Xi_i : I_i \rightarrow J_i$  such that  $\Xi_i(C_i^{\underline{d}})$  is the set of  $\lambda$  such that the first return of 0 to  $H_i[\lambda](I_i)$  under iteration by  $f_\lambda$  belongs to  $H_i[\lambda](C_i^{\underline{d}})$ .

**Definition 7.2.** Let  $f_\lambda$  be a family of unimodal maps. We say that  $f_\lambda$  has Decay of Parameter Geometry at  $\lambda_0$  if  $f = f_{\lambda_0} \in \mathcal{F}$ , it satisfies the Topological Phase-Parameter relation at  $\lambda_0$  and  $|J_{n_k+1}|/|J_{n_k}| < C\lambda^k$  for some constants  $C > 0$ ,  $\lambda < 1$ , where  $n_k - 1$  counts the non-central levels of the principal nest of  $f$ .

**Theorem 7.1.** *Let  $f \in \mathcal{F}$  be analytic. There exists a polynomial vector field  $w$  such that the family  $f_{\lambda w} = f \circ (\text{id} + \lambda w)$ ,  $\lambda \in (-\epsilon, \epsilon)$  is an analytic family of unimodal maps which satisfies the Topological Phase-Parameter relation and has Decay of Parameter Geometry at 0.*

*Proof.* Let  $w$  and  $\mathbf{n}$  be as in Lemma 6.12. Denote by  $(\mathcal{R}_1, h_1)$  the  $R$ -family of that lemma. Since  $f \in \mathcal{F}$ , the critical point is recurrent and we can clearly construct a  $R$ -chain  $(\mathcal{R}_i, h_i)$  over  $\lambda = 0$ . It is clear that the real trace of  $R_i[0] : \cup U_i^j[0] \rightarrow U_i[0]$  is the first return map to  $I_{\mathbf{n}+i}$ . Let  $J_{\mathbf{n}+i} = \Lambda_i \cap \mathbb{R}$ , let  $\Xi_{\mathbf{n}+i} = \chi_i[0]|_{I_{\mathbf{n}+i}}$ . It is clear that  $|J_{n_k+1}|/|J_{n_k}|$  decays exponentially by Lemma 6.13 and Theorem 4.7,

<sup>5</sup>More precisely, we use that the holomorphic map  $\lambda \mapsto f_\lambda^r(0)$  has the same derivative at 0 as the almost linear map  $\lambda \mapsto f_{\lambda v_m}^r(0)$ , and a simple estimate shows that there exists a parameter  $\lambda \in \mathbb{D}$  such that  $f_{\lambda v_m}^r(0) = 0$ .

<sup>6</sup>In the sense of admitting a controlled extension to a big domain, as the Beltrami path we constructed.

where  $n_k - 1$  counts the non-central levels of the principal nest of  $f$ . In particular,  $|J_n| \rightarrow 0$ .

In order to conclude the result, we just have to show the existence of the continuous family of homeomorphisms  $H_i[\lambda]$ , for  $i$  sufficiently big. Notice that if  $\lambda \in J_{n+i}$ , if  $p \in I_{n+i}[\lambda]$  is a periodic orbit for  $f_\lambda$  which never enters  $I_{n+i}^0[\lambda]$  then  $p$  is hyperbolic by the Schwarz Lemma. So, if  $\lambda \in J_{n+i}$ , the only non-hyperbolic periodic orbits for  $f_\lambda$  must be entirely contained in  $I \setminus I_{n+1}$ . But since  $f|I \setminus I_{n+1}$  is hyperbolic, there exists  $\epsilon > 0$  such that if  $\lambda \in (-\epsilon, \epsilon)$ , all periodic orbits in  $I \setminus I_{n+1}[\lambda]$  of  $f_\lambda$  are hyperbolic (by Lemma 5.2). In particular, if  $i$  is sufficiently big,  $J_i \subset (-\epsilon, \epsilon)$ , and all periodic orbits of  $f_\lambda$  in  $I \setminus I_{i+1}[\lambda]$  are hyperbolic. The result follows by Lemma 5.3.  $\square$

Let  $K_i = I_i \setminus \cup C_i^d$ . Notice that  $H_i$  and  $\Xi_i$  are only uniquely defined in  $K_i$ . Condition (2) of the Topological Phase-Parameter relation can be equivalently formulated as the existence of a homeomorphism  $\Xi_i : I_i \rightarrow J_i$  such that the first return of the critical point (under iteration by  $f_\lambda$ ) to  $H_i[\lambda](I_i)$  belongs to  $H_i[\lambda](K_i)$  if and only if  $\lambda \in \Xi_i(K_i)$ .

Let us now estimate the metric properties of  $H_i|K_i$  and  $\Xi_i|K_i$ .

Let  $\tilde{I}_{i+2} = (R_i|I_i^0)^{-1}(I_i^d)$ , where  $\underline{d}$  is such that  $(R_i|I_i^0)^{-1}(C_i^{\underline{d}}) = I_{i+1}$ .

Let  $\tau_i$  such that  $R_i(0) \in I_i^{\tau_i}$ . Let  $K_i^\tau = K_i \cap I_i^{\tau_i}$  and  $\tilde{K}_i = I_i \setminus (\cup I_i^j \cup \tilde{I}_{i+1})$ .

Let  $J_i^j = \Xi_i(I_i^j)$ .

**Definition 7.3.** Let  $f_\lambda$  be a family of unimodal maps. We say that  $f_\lambda$  satisfies the Phase-Parameter relation at  $\lambda_0$  if  $f = f_{\lambda_0}$  is simple,  $f_\lambda$  satisfies the Topological Phase-Parameter relation at  $\lambda_0$  and for every  $\gamma > 1$ , there exists  $i_0 > 0$  such that:

**PhPa1:**  $\Xi_i|_{K_i \cap I_i^{\tau_i}}$  is  $\gamma$ -qs,

**PhPa2:**  $\Xi_i|_{\tilde{K}_i}$  is  $\gamma$ -qs,

**PhPh1:**  $H_i[\lambda]|_{K_i}$  is  $\gamma$ -qs if  $\lambda \in J_i^{\tau_i}$ ,

**PhPh2:** the map  $H_i[\lambda]|_{\tilde{K}_i}$  is  $\gamma$ -qs if  $\lambda \in J_i$ .

**Theorem 7.2.** *In the same setting of the previous theorem, if  $f$  is simple, the family  $f_{\lambda_w}$  satisfies the Phase-Parameter relation at 0.*

*Proof.* This is an immediate consequence of Theorem 3.1 and Theorem 4.7, since the Phase-Parameter relation is just the real trace of the Complex Phase-Parameter relation (see [AM1] for details).  $\square$

More details, more precise reference

## 8. LAMINATION IN SPACES OF UNIMODAL MAPS

**8.1. Spaces of analytic unimodal maps.** Let  $a > 0$ , and let  $\Omega_a \subset \mathbb{C}$  be the set of points at distance at most  $a$  of  $I$ . Let  $\mathcal{E}_a$  be the complex Banach space of holomorphic maps  $v : \Omega_a \rightarrow \mathbb{C}$  continuous up to the boundary which are 0-symmetric (that is,  $v(z) = v(-z)$ ) and such that  $v(-1) = v(1) = 0$ , endowed with the sup-norm  $\|v\|_a = \|v\|_\infty$ . It contains the real Banach space  $\mathcal{E}_a^{\mathbb{R}}$  of "real maps"  $v$ , i.e, holomorphic maps symmetric with respect to the real line:  $v(\bar{z}) = \overline{v(z)}$ .

Let us consider the constant function  $-1 \in \Omega_a$ . The complex affine subspace  $-1 + \mathcal{E}_a$  will be denoted as  $\mathcal{A}_a$ .

Let  $\mathbb{U}_a = \mathbb{U}^2 \cap \mathcal{A}_a$ . It is clear that any analytic unimodal map belongs to some  $\mathbb{U}_a$ . Note that  $\mathbb{U}_a$  is the union of an open set in the affine subspace  $\mathcal{A}_a^{\mathbb{R}} = -1 + \mathcal{E}_a^{\mathbb{R}}$  and a codimension-one space of unimodal maps satisfying  $f(0) = 1$ .

**8.2. Laminations.** One of the main results of [ALM] is to describe the structure of the partition in topological classes of spaces of analytic unimodal maps. In that paper, they consider only the quasiquadratic case, but their work holds for the general case (due to the results of Kozłowski). The main result is that each non-hyperbolic topological class is a codimension-one analytic submanifold, which form an analytic lamination near any Kupka-Smale quasiquadratic map.

**Theorem 8.1** (Theorem A of [ALM]). *Let  $f \in \mathbb{U}_a$  be a Kupka-Smale map. There exists a neighborhood  $\mathcal{V} \subset \mathcal{A}_a$  of  $f$  endowed with a codimension-one holomorphic lamination  $\mathcal{L}$  (also called hybrid lamination) with the following properties:*

- (1) *the lamination is real-symmetric;*
- (2) *if  $g \in \mathcal{V} \cap \mathcal{A}_a^{\mathbb{R}}$  is non-regular, then the intersection of the leaf through  $g$  with  $\mathcal{A}_a^{\mathbb{R}}$  coincides with the intersection of the topological conjugacy class of  $g$  with  $\mathcal{V}$ ;*
- (3) *Each  $g \in \mathcal{V} \cap \mathcal{A}_a^{\mathbb{R}}$  belongs to some leaf of  $\mathcal{L}$ .*

(For the definition of the leaves of  $\mathcal{L}$  in the non-regular case, see Appendix A.)

**Theorem 8.2.** *In the setting of Theorem 8.1, if  $g_1, g_2 \in \mathcal{V}$  are in the same leaf of  $\mathcal{L}$  and  $\gamma_1(\lambda), \gamma_2(\lambda)$  are real analytic paths in  $\mathcal{V} \cap \mathcal{A}_a^{\mathbb{R}}$ , transverse to the leaves of  $\mathcal{V}$  and such that  $\gamma_1(\lambda_1) = g_1, \gamma_2(\lambda_2) = g_2$ , then the local holonomy map  $\psi : (\lambda_1 - \epsilon, \lambda_1 + \epsilon) \rightarrow (\lambda_2 - \epsilon', \lambda_2 + \epsilon')$  is quasiasymmetric. Moreover, for  $\delta$  sufficiently small,  $\psi|_{(\lambda_1 - \delta, \lambda_1 + \delta)}$  is  $1 + O(\|g_1 - g_2\|_a)$ -qs.*

*Proof.* This is just the  $\lambda$ -Lemma.  $\square$

Moreover, each non-regular topological class is like a Teichmüller space:

**Theorem 8.3.** *In the setting of Theorem 8.1, if  $g_1, g_2 \in \mathcal{V} \cap \mathbb{U}_a$  belong to the same leaf, then there exists  $1 + O(\|g_1 - g_2\|_a)$ -qs map  $h : I \rightarrow I$  such that  $g_2 \circ h = h \circ g_1$ .*

*Proof.* This follows from Proposition 8.9 of [ALM] and the  $\lambda$ -Lemma.  $\square$

The tangent space to topological classes has a nice characterization:

**Theorem 8.4** (Theorem 8.10 of [ALM]). *If  $f \in \mathbb{U}_a$  is a non-regular Kupka-Smale map then the tangent space to the topological class of  $f$  is given by the set of vector fields  $v \in \mathcal{E}_a$  which do not admit a representation  $v = \alpha \circ f - \alpha Df$  on the critical orbit with  $\alpha$  a qc vector field of  $\mathbb{C}$ .*

**8.3. Analytic families.** Let  $\{f_\lambda\}_{\lambda \in \Lambda}$  be an analytic family of unimodal maps. Then for  $a > 0$  sufficiently small,  $\lambda \mapsto f_\lambda$  is an analytic map from  $\Lambda$  to  $\mathbb{U}_a$ .

If  $\lambda_0 \in \Lambda$  is a Kupka-Smale parameter, transversality to the topological class of  $\lambda_0$  has the obvious meaning (using Theorem 8.1). We remark that this definition does not depend on the choice of  $\mathbb{U}_a$ .

*Remark 8.1.* Let  $B_i$  be an enumeration of all open balls contained in  $\Lambda$  of rational radius and center. The condition of non-triviality of a family  $\{f_\lambda\}$ ,  $\lambda \in \Lambda$  is an intersection of a countable number of conditions (existence of a regular parameter  $\lambda \in B_i$ ). Each of those conditions is open in  $\mathbb{U}\mathbb{F}^2(\Lambda)$ . The set of non-trivial analytic families is also dense in the  $\mathbb{U}\mathbb{F}^\infty(\Lambda)$  (this would still hold natural topology of analytic families in  $\Lambda$ , which we did not introduce), due to Theorem 8.1.

We should remark that for an analytic family of quasiquadratic maps, non-triviality is equivalent to existence of one unique regular parameter (since all non-regular topological classes are analytic submanifolds in the quasiquadratic case). In particular, non-triviality is a  $C^3$  open condition in the quasiquadratic case.

More details

**8.4. Phase-parameter relation for transverse families.** We will now show how to use the lamination of [ALM] to transfer the Phase-Parameter relation from the transversal family  $f_{\lambda w}$  to any transversal family  $f_\lambda$ . The idea is to exploit the following commutative diagram

$$\begin{array}{ccc}
 \text{Phase of } f_{\lambda w} & \xrightarrow[\text{qs conjugacy}]{\text{Theorem 8.3}} & \text{Phase of } f_\lambda \\
 \downarrow \text{Phase-Parameter for } f_{\lambda w} & & \downarrow \text{Phase-Parameter for } f_\lambda \\
 \text{Parameter of } f_{\lambda w} & \xrightarrow[\text{holonomy map of } \mathcal{L}]{\text{Theorem 8.2}} & \text{Parameter of } f_\lambda
 \end{array}$$

(notice that the estimates for all arrows are all ultimately based on the  $\lambda$ -Lemma).

**Theorem 8.5.** *Let  $f \in \mathcal{F}$ , and let  $f_\lambda$  be an analytic family of unimodal maps through  $f$  such that  $f_{\lambda_0} = f$  and  $f_\lambda$  is transverse to the topological class of  $f$  at  $\lambda = \lambda_0$ . Then the Topological Phase-Parameter relation and Decay of Parameter Geometry holds for the family  $f_\lambda$  at  $\lambda_0$ . Moreover, if  $f$  is simple, then the Phase-Parameter relation also holds.*

*Proof.* Using Theorems 7.1 and 7.2, consider a family  $f_{\lambda w}$  through  $f$  which is transverse to the hybrid class of  $f$  and which satisfies the Topological Phase-Parameter Relation/Decay of Parameter Geometry/Phase-Parameter relation. Fix  $a$  such that both  $f_{\lambda w}$  and  $f_\lambda$  are analytic paths in  $\mathbb{U}_a$ . Let  $\mathcal{L}$  be the lamination from Theorem 8.1. Since both  $f_\lambda$  and  $f_{\lambda w}$  are transverse to the topological class of  $f$  (at  $\lambda_0$  and 0), we can consider the local holonomy map of the lamination  $\mathcal{L}$ ,  $\psi : (-\epsilon, \epsilon) \rightarrow (\lambda_0 - \epsilon', \lambda_0 + \epsilon')$ .

Let  $\tilde{\Xi}_i : I_i \rightarrow \tilde{J}_i$  be the phase-parameter map for the family  $f_{\lambda w}$ , and let  $\tilde{H}_i[\lambda]$  be the phase-phase map. We obtain the phase-parameter map for  $f_\lambda$  as a composition  $\Xi_i = \psi \circ \tilde{\Xi}_i$ . Since  $|\tilde{J}_i| \rightarrow 0$ ,

$$\lim_{i \rightarrow \infty} \sup_{\lambda \in (-\epsilon, \epsilon)} \|f_{\lambda w} - f_{\psi(\lambda)}\|_a = 0.$$

In particular, by Theorem 8.2,  $\psi|_{\tilde{J}_i}$  is  $\gamma_i$ -qs with  $\lim \gamma_i = 1$ .

Since for each  $\lambda \in J_i = \psi(\tilde{J}_i)$ ,  $f_\lambda$  is qs conjugate to  $f_{\psi^{-1}(\lambda)w}$ , we see that if  $\lambda \in J_i$  then there are no non-hyperbolic periodic orbits for  $f_\lambda$  in the complement of the continuation of  $I_{i+1}$ . Using Lemma 5.2 we conclude as in Theorem 7.1 the existence of a continuous family  $H_i[\lambda]$  of phase-phase maps for the family  $f_\lambda$ . It follows that the Topological Phase-Parameter relation holds for  $f_\lambda$  at  $\lambda_0$ .

Since  $\psi$  is quasisymmetric, it is Hölder and the Decay of Parameter Geometry also follows from Theorem 7.1. If  $f$  is simple, estimates PhPa1 and PhPa2 follow from Theorem 7.2.

Let  $h_\lambda : I \rightarrow I$  be a quasisymmetric conjugacy between  $f_{\lambda w}$  and  $f_{\psi(\lambda)}$  which is  $O(\|f_{\lambda w} - f_{\psi(\lambda)}\|_a)$ -qs. This family might not be continuous, but  $H_i[\psi(\lambda)]K_i = h_\lambda \circ \tilde{H}_i[\lambda]$ , which is enough for our purposes. In particular, if  $f$  is simple, PhPh1 and PhPh2 follow from Theorem 7.2.  $\square$

*Remark 8.2.* Notice that even if we are only interested in the phase-parameter relation for individual families, this proof needs the knowledge of the behavior of topological conjugacy classes of unimodal maps in infinite dimensional spaces. For the case of the quadratic family, this is not needed: the argument of [L3] is based on the combinatorial theory of the Mandelbrot set (Douady-Hubbard, Yoccoz), which

allows to show directly that the real quadratic family gives rise to full unfolded complex return type families. In particular, our proof also gives a somewhat different approach to the phase-parameter relation on the quadratic family itself.

## 9. PROOF OF THEOREM A

Let  $f_\lambda$  be a non-trivial analytic family of unimodal maps. In view of Theorem 8.5, to conclude Theorem A it is enough to show that

- (1) Almost every non-regular parameter belongs to  $\mathcal{F}$ , that is, it is Kupka-Smale, has a recurrent critical point and is not infinitely renormalizable,
- (2) Almost every parameter in  $\mathcal{F}$  is simple,
- (3)  $f_\lambda$  is transverse to the topological class of almost every parameter.

We will take care of these issues separately below: item (1) will follow from Lemmas 9.1, 9.4, and 9.5, item (2) from Lemma 9.6 and item (3) from Lemma 9.3.

### 9.1. Transversality.

**Lemma 9.1.** *Let  $f_\lambda$  be a non-trivial analytic family of unimodal maps. Then at most countably many parameters are not Kupka-Smale or have a periodic or preperiodic critical point.*

*Proof.* Indeed, the set of parameters which are not Kupka-Smale correspond to solutions of countably many analytic equations of the type  $f_\lambda^n(p) = p$ ,  $Df_\lambda^{2n}(p) = 1$ ,  $n > 0$ . Similarly, the set of parameters with periodic or preperiodic critical point corresponds to countably many equations of the type  $f_\lambda^m(0) = f_\lambda^n(0)$ ,  $0 \leq m < n$ . So the set of parameters which are not Kupka-Smale is either countable or contains intervals. Since regular parameters are dense, the first possibility holds.  $\square$

The following result is due to Douady, see Lemma 9.1 of [ALM]:

**Lemma 9.2.** *Let  $\mathcal{L}$  be a codimension-one complex lamination on an open set  $\mathcal{V}$  of some Banach space, and let  $\gamma$  be an analytic path in  $\mathcal{V}$ . If  $\gamma$  is not contained in a leaf of  $\mathcal{L}$ , then the set of parameters where  $\gamma$  is not transverse to the leaves of  $\mathcal{L}$  consists of isolated points.*

This result immediately implies:

**Lemma 9.3.** *Let  $f_\lambda$  be a non-trivial analytic family of unimodal maps. Then the set of non-regular Kupka-Smale parameters  $\lambda_0$  such that  $f_\lambda$  is not transverse to the topological class  $f_{\lambda_0}$  at  $\lambda_0$  is countable.*

**9.2. Non-recurrent parameters.** The following result is due to Duncan Sands [S], but we will provide a quick proof based on holomorphic motions and Lemma 9.2.

**Lemma 9.4.** *Let  $f_\lambda$  be a non-trivial analytic family of unimodal maps. Then almost every parameter is regular or has a recurrent critical point.*

*Proof.* If this is not the case, there would exist  $\epsilon > 0$  and a set  $X$  of parameters  $\lambda$  of positive measure such that for  $\lambda \in X$ ,

- (1)  $\inf_{m \geq 1} |f_\lambda^m(0)| > \epsilon$  (by hypothesis).
- (2)  $f_\lambda$  is non-regular and Kupka-Smale (by Lemma 9.1).



Let us fix a density point  $\lambda_0 \in X$  of  $X$ . Using Lemma 5.1, consider a nice interval  $T = T[\lambda_0] = [-p, p] \subset (-\epsilon, \epsilon)$  for  $f_\lambda$ , with  $p$  periodic. Let  $T[\lambda]$ ,  $\lambda - \lambda_0 \in (-\delta, \delta)$ ,  $\delta > 0$  small denote the continuation of  $T$ . Let  $K[\lambda]$ ,  $\lambda - \lambda_0 \in (-\delta, \delta)$  denote the set of points in  $I \setminus T[\lambda]$  which never enter  $T[\lambda]$  and do not belong to the basin of hyperbolic attractors.

Since  $K = K[\lambda_0]$  is an expanding set by Lemma 5.2, it persists in a complex neighborhood of  $\lambda_0$ : there exists a family of homeomorphisms  $h_\lambda : K \rightarrow \mathbb{C}$ ,  $\lambda \in \mathbb{D}_{\delta'}(\lambda_0)$ ,  $\delta' < \delta$  depending continuously on  $\lambda$ , such that  $h_\lambda = \text{id}$  and  $f_\lambda \circ h_\lambda = h_\lambda \circ f_{\lambda_0}$ . It is easy to see (using Lemma 5.2) that for  $\lambda \in \mathbb{R}$ ,  $h_\lambda(K) = K[\lambda]$ . For each preperiodic orbit  $p$  of  $f$  in  $K$ , it is clear that  $\lambda \mapsto h_\lambda(p)$  is holomorphic in  $\mathbb{D}_{\delta'}(\lambda_0)$ . Since preperiodic orbits are dense in  $K$ , it follows that  $h[\lambda_0, \lambda] \equiv h_\lambda$  are actually transition maps of a holomorphic motion  $h$  over  $\mathbb{D}_{\delta'}(\lambda_0)$ .

Since  $f_\lambda$  is non-trivial,  $f_\lambda(0)$  does not belong to  $K[\lambda]$  for a dense set of  $\lambda \in (-\delta, \delta)$ , so by Lemma 9.2, the path  $\lambda \mapsto (\lambda, f_\lambda(0))$  is transverse to the leaves of  $h$  outside of countably many parameters  $\lambda$ . Perturbing  $\lambda_0$  if necessary (keeping both properties (1) and (2) above), we may assume that  $\lambda_0$  is a point of transversality. It follows that there exists a real-symmetric quasiconformal map  $\chi$  (phase-parameter holonomy map) taking a neighborhood  $V$  of  $f_{\lambda_0}(0)$  to a neighborhood of  $\lambda_0$ , and taking points in  $K \cap V$  to parameters  $\lambda \in \chi(V)$  with  $f_\lambda(0) \in K[\lambda]$ . In particular,  $\chi(K \cap V) \supset X \cap \chi(V)$ .

Since  $K$  is an expanding set, it follows that there exists  $\rho > 0$  such that in every  $r$  neighborhood of  $f_{\lambda_0}(0)$  there exists an interval of size at least  $\rho r$  disjoint from  $K$ . Since  $\chi|_V \cap \mathbb{R}$  is quasisymmetric, this property is preserved: there exists  $\rho' > 0$  such that in every  $r$  neighborhood of  $\lambda_0$  there exists an interval of size at least  $\rho' r$  not intersecting  $X$ . This contradicts the hypothesis that  $\lambda_0$  is a density point of  $X$ .

(It is easy to see that this argument gives much more information on the size of  $X$ . One can see for instance that the Hausdorff dimension of  $X$  in  $\lambda_0$  (defined as the infimum of the Hausdorff dimension of  $X \cap \mathbb{D}_\epsilon(\lambda_0)$ ) is no greater than the Hausdorff dimension of  $K$  in  $f_{\lambda_0}(0)$ , which is known to be less than 1. Notice that  $X$  is essentially the set of non-regular non-recurrent parameters avoiding a definite neighborhood of 0. We should remark that these ideas show also that the Hausdorff dimension of the set of non-regular non-recurrent parameters is usually 1 except for some trivial situations.)  $\square$

### 9.3. Infinitely renormalizable maps.

**Lemma 9.5.** *Let  $f_\lambda$  be a non-trivial analytic family of unimodal maps. Then the set of infinitely renormalizable parameters has Lebesgue measure zero.*

*Proof.* Let  $X$  be the set of parameters  $\lambda$  such that  $f_\lambda$  is infinitely renormalizable, and let  $\lambda_0 \in X$  be a density point of  $X$ . By Lemma 5.4, there exists a nice interval  $T[\lambda]$ ,  $|\lambda - \lambda_0| < \delta$ , which is periodic (of period, say  $m$ ) such that  $f^m|_{T[\lambda]}$  has negative Schwarzian derivative. In particular, if  $A_\lambda : T[\lambda] \rightarrow I$  is affine,  $g_\lambda = A_\lambda \circ f_\lambda^m \circ A_\lambda^{-1}$  is an analytic family of unimodal maps, which is non-trivial (because  $f_\lambda$  is). By Theorem B of [ALM], for almost every  $\lambda$ ,  $g_\lambda$  is not infinitely renormalizable. It is clear that if  $\lambda \in X$  and  $|\lambda - \lambda_0| < \delta$  then  $g_\lambda$  is infinitely renormalizable, so  $\lambda_0$  is not a density point of  $X$ , contradiction.  $\square$

**9.4. Simple maps.** The following argument is adapted from the corresponding result of Lyubich for the quadratic family [L3].

**Lemma 9.6.** *Let  $f_\lambda$  be a non-trivial analytic family of unimodal maps. Then almost every parameter  $\lambda$  with  $f_\lambda \in \mathcal{F}$  is simple.*

*Proof.* If this is not the case, we could find  $C > 0$ ,  $\rho < 1$ ,  $\kappa \geq 0$ ,  $m \geq 0$  and a set  $X$  of parameters of positive measure such that if  $\lambda_0 \in X$  then

- (1)  $f_{\lambda_0} \in \mathcal{F}$  and is not simple (by hypothesis),
- (2)  $f_\lambda$  is transverse at  $\lambda_0$  (by Theorem 9.3),
- (3) The sequence of parameter windows  $J_n[\lambda_0]$  associated to  $\lambda_0$  are defined for  $n \geq m$  (by Theorem 8.5, item (1)),
- (4) If  $n_k(\lambda_0) - 1$  denotes the sequence of non-central levels of the principal nest of  $f_{\lambda_0}$  then for  $n_k(\lambda_0) \geq m$ ,  $|J_{n_k(\lambda_0)+1}[\lambda_0]|/|J_{n_k(\lambda_0)}[\lambda_0]| < C\rho^k$  (by Theorem 8.5).

Consider now the set  $X_k$ ,  $k \geq m$  of parameters  $\lambda_0 \in X$  such that the return of level  $n_k(\lambda_0)$  is central. Let  $\Delta_k$  be the union of  $J_{n_k(\lambda_0)}[\lambda_0]$ ,  $\lambda_0 \in X_k$  and  $\Pi_k$  be the union of  $J_{n_k(\lambda_0)+1}[\lambda_0]$ ,  $\lambda_0 \in X_k$ .

Then each connected component  $J_{n_k(\lambda_0)}[\lambda_0]$  of  $\Delta_k$  contains a single connected component  $J_{n_k(\lambda_0)+1}[\lambda_0]$  of  $\Pi_k$ , and thus  $|\Pi_k|/|\Delta_k| < C\rho^k$ , so that  $|X_k| \leq |\Pi_k| < C\rho^k|\Delta_k| \leq 2C\rho^k$ . On the other hand,  $X \subset \bigcap_{k_0 \geq m} \bigcup_{k \geq k_0} X_k$  and thus,  $|X| \leq \inf_{k_0 \geq m} \sum_{k \geq k_0} 2C\rho^k = 0$ , contradiction.  $\square$

## 10. PROOF OF THEOREM B

We will give now a proof of Theorem B using a parameter exclusion argument. The first proof of this result in [Av1] relied on the refined statistical argument of [AM1], but we will give a much simpler argument based on a modified version of the quasisymmetric capacities of [AM1], which allows us to get rid of distortion estimates and at the same time to work with a fixed quasisymmetric constant.

**10.1. Measure estimate.** Define the modified  $\gamma$ -qs capacity of a set  $X$  in an interval  $I$  as

$$p_\gamma(X|I) = \sup \frac{|h_1 \circ h_2(X \cap I)|}{|h_1 \circ h_2(I)|}$$

where  $h_1 : \mathbb{R} \rightarrow \mathbb{R}$  is  $\gamma$ -qs and  $h_2 : I \rightarrow \mathbb{R}$  has non-negative Schwarzian derivative.

Notice that if  $F : T_1 \rightarrow T_2$  is a diffeomorphism with non-positive Schwarzian derivative and  $X \subset T_1$  then

$$p_\gamma(X|T_1) \leq p_\gamma(F(X)|T_2).$$

This is the main advantage of modified quasisymmetric capacities over the traditional ones of [AM1].

By the Koebe Principle (see [MS]), if  $h : I \rightarrow I$  is a diffeomorphism and has non-positive Schwarzian derivative then  $h([-\epsilon, \epsilon]) = O(\epsilon)$ . By Hölder continuity of  $\gamma$ -qs maps, we get

$$p_\gamma([-\epsilon, \epsilon]|[-1, 1]) = O(\epsilon^k)$$

Reference for the Koebe Principle for some  $0 < k < 1$  depending on  $\gamma$ .

For a map  $f \in \mathcal{F}$  with principal nest  $\{I_n\}$ , let  $s$  be as in Lemma 5.5, and let

$$\alpha_n = p_\gamma(s(\cup I_n^j)|s(I_n)).$$

Let us consider the components  $T_n^k$  of  $(R_n|I_n^0)^{-1}(\cup I_n^j)$ . We reserve the index 0 for the component containig 0, and the indexes  $-1$  and  $1$  for the components of

$(R_n|I_n^0)^{-1}(I_n^0)$ . If  $|k| > 1$  then  $R_n|T_n^k$  is a diffeomorphism onto some  $I_n^j$ ,  $j \neq 0$  and  $R_n^2|T_n^k$  is a diffeomorphism onto  $I_n$ . Let

$$\epsilon_n = p_\gamma(s(\cup_{i=-1}^1 T_n^i)|s(I_{n+1})).$$

**Lemma 10.1.** *The following estimate holds for  $n$  big:*

$$(1 - \alpha_{n+1}) \geq (1 - \epsilon_n)(1 - \alpha_n)$$

*Proof.* Indeed, if  $|k| > 1$  then  $s(T_{n+1}^k)$  is taken to  $s(I_n)$  by  $s \circ R_n^2 \circ s^{-1}$  which has negative Schwarzian derivative for  $n$  big. In particular

$$p_\gamma(s(\cup I_{n+1}^j)|s(T_{n+1}^k)) \leq p_\gamma(s(\cup C_n^d)|s(I_n)) \leq \alpha_n.$$

Thus

$$p_\gamma(\cup I_{n+1}^j|I_{n+1}) \leq \epsilon_n + (1 - \epsilon_n)\alpha_n.$$

□

**Lemma 10.2.** *If  $f$  is simple then the  $\epsilon_n$  decay exponentially fast.*

*Proof.* Indeed, if  $f$  is simple then  $|s(I_{n+1})|/|s(I_n)|$  decays exponentially fast by Lemma 5.6. In particular, by the Koebe Principle, for each  $j$ , each of the connected components of  $s(I_{n+1} \setminus I_{n+1}^j)$  is exponentially (in  $n$ ) bigger than  $s(I_{n+1}^j)$ . This implies that, for each  $k$ , each component of  $s(I_{n+2} \setminus T_{n+2}^k)$  is exponentially bigger than  $T_{n+2}^k$  (using the Koebe Principle), so  $p_\gamma(s(T_{n+2}^k)|s(I_{n+2}))$  decays exponentially and so does  $\epsilon_n$ . □

**Lemma 10.3.** *If  $f$  does not admit a quasiquadratic renormalization then  $\cup I_n^j$  is not dense in  $I_n$ , for  $n$  sufficiently big.*

*Proof.* Indeed, up to considering a renormalization or unimodal restriction, we may assume that  $f$  is non-renormalizable and does not admit unimodal restriction. It is easy to see that if  $x \in I$  never enters  $I_1$  then  $x$  accumulate on a orientation preserving fixed point of  $f$ , and since  $f$  does not admit a unimodal restriction, we conclude that  $x \in \partial I$ .

Since  $f$  is not conjugate to a quadratic map, there exists an interval  $T$  whose orbit does not accumulate on the critical point (Lemma 5.8). Let  $n$  be biggest with the orbit of  $T$  intersecting  $I_n$  ( $T$  intersects  $I_1$  by the previous discussion). Of course, the set of points which land on  $I_{n+1}$  does not intersect the orbit of  $T$ , and so is not dense in  $I_n$ .

It is easy to see that if the set of points in  $I_m$  which eventually land in  $I_{m+1}$  is not dense in  $I_n$  then  $\cup I_{m+1}^j$  is not dense on  $I_{m+1}$ . In particular, by induction,  $\cup I_m^j$  is not dense in  $I_m$  for  $m \geq n$ . □

**Lemma 10.4.** *If  $f$  does not admit a quasiquadratic renormalization then for  $n$  large enough,  $\alpha_n < 1$ .*

*Proof.* Let  $n$  be large enough such that there exists an interval  $E \subset I_n$  disjoint from  $\cup I_n^j$ . For  $j \neq 0$ , let  $E^j = (R_n|I_n^j)^{-1}(E)$ . Notice that  $E^j$  does not intersect  $\cup C_n^d$ .

If  $h_1 : \mathbb{R} \rightarrow \mathbb{R}$  is a  $\gamma$ -qs map and  $h_2 : s(I_n) \rightarrow \mathbb{R}$  has non-negative Schwarzian derivative, we see that  $h_2|s(I_n^j)$  has bounded distortion by the Koebe Principle. Notice that this implies that for some  $\epsilon > 0$ , if  $j \neq 0$ ,  $|h_1 \circ h_2(s(E^j))| > \epsilon |h_1 \circ h_2(I_n^j)|$ . Thus

$$p_\gamma(s(I_n \setminus \cup E^j)|s(I_n)) \leq p_\gamma(s(I_n^0)|s(I_n)) + (1 - p_\gamma(s(I_n^0)|s(I_n)))(1 - \epsilon) < 1.$$

By a previous argument of the proof of Lemma 10.1,

$$p_\gamma(s(\cup I_{n+1}^j)|s(I_{n+1}^k)) \leq p_\gamma(s(\cup C_n^d)|s(I_n)) \leq p_\gamma(s(I_n \setminus \cup E^j)|s(I_n)).$$

Thus,

$$p_\gamma(s(\cup I_{n+1}^j)|s(I_{n+1})) \leq \epsilon_n + (1 - \epsilon_n)p_\gamma(s(I_n \setminus \cup E^j)|s(I_n)) < 1.$$

□

**Lemma 10.5.** *Let  $f_\lambda$  be a one-parameter non-trivial analytic family of unimodal maps satisfying the Phase-Parameter relation at a parameter  $\lambda_0$  (in particular,  $f = f_{\lambda_0}$  is simple). Assume that  $f$  does not admits quasiquadratic renormalization. Then  $\lambda_0$  is not a density point of non-hyperbolic parameters.*

*Proof.* Since  $|J_n| \rightarrow 0$ , and  $\lambda_0 \in \Xi_n(I_n^{\tau_n}) \subset J_n$ , it is enough to show that if  $n$  is big enough then  $|\Xi_n(\cup C_n^d \cap I_n^{\tau_n})| \leq \alpha_n |\Xi_n(I_n^{\tau_n})|$ . Indeed, if  $\lambda \notin \Xi_n(\cup C_n^d)$  then the critical point is non-recurrent. By Lemma 9.4, almost every non-recurrent parameter in  $f_\lambda$  is hyperbolic.

Fix  $1 < \hat{\gamma} < \gamma$ . By PhPa1,  $\Xi(K_n^\tau)$  has a  $\hat{\gamma}$ -qs extension (that we denote  $\Xi_n$ ) for  $n$  big enough. On the other hand,  $s^{-1}|s(I_n^{\tau_n})$  is essentially linear for  $n$  big (because  $s$  is analytic, and in particular  $C^1$ , and  $s(I_n^{\tau_n})$  is small), so  $\Xi_n \circ s^{-1}|s(I_n^{\tau_n})$  is  $\gamma$ -qs. In particular

$$\frac{|\Xi(\cup C_n^d \cap I_n^{\tau_n})|}{|\Xi(I_n^{\tau_n})|} \leq \frac{|\Xi \circ s^{-1}s(\cup C_n^d \cap I_n^{\tau_n})|}{|\Xi \circ s^{-1}s(I_n^{\tau_n})|} \leq p_\gamma(s(\cup C_n^d)|s(I_n^{\tau_n})) \leq \alpha_n.$$

By Lemmas 10.1, 10.2 and 10.3,  $\limsup \alpha_n < 1$ . □

For one-parameter families, Theorem B follows from Theorem A and Lemma 10.5.

10.1.1. *Many parameters.* The argument of Lemma 9.1 implies:

**Lemma 10.6.** *Let  $\{f_\lambda\}_{\lambda \in \Lambda}$  be a  $k$ -parameter non-trivial analytic family of unimodal maps. The set of parameters which are not Kupka-Smale or have a periodic or preperiodic critical point is contained in a countable union of analytic submanifolds of codimension at least 1 of  $\Lambda$  and so has Lebesgue measure zero.*

Let  $\{f_\lambda\}_{\lambda \in \Lambda}$  be a  $k$ -parameter analytic family of unimodal maps, and let  $\lambda_0 \in \text{int } \Lambda$  be a Kupka-Smale parameter. Let us consider a small ball  $B_\epsilon$  around  $\lambda_0$  of radius  $\epsilon$  contained in  $\Lambda$ . By Theorem 10.6, it is clearly enough to show Theorem B for the family  $f_\lambda$  restricted to  $B_\epsilon$ .

Using Theorem 8.1, if  $\epsilon$  is sufficiently small,  $\lambda \mapsto f_\lambda$  is an analytic map from  $B_\epsilon$  to some open set  $\mathcal{V}$  where the hybrid lamination  $\mathcal{L}$  is defined. Let  $\lambda_1 \in B_\epsilon$  be a regular parameter. If  $L$  is a line in  $\mathbb{R}^k$  through  $\lambda_1$ , then by Lemma 9.2  $L \cap B_\epsilon$  is not contained in the topological class of a non-regular parameter, and so regular parameters are dense in  $L \cap B_\epsilon$ . By the one-dimensional Theorem B, we see that almost every parameter in  $L \cap B_\epsilon$  is quasiquadratic. By Fubini's Theorem, almost every parameter in  $B_\epsilon$  is quasiquadratic.

## 11. PROOF OF COROLLARIES

**11.1. Some conditions related to good ergodic properties.** Let us first recall the conditions on the critical orbit stated in the introduction. Let  $f \in \mathbb{U}^2$ . We say that  $f$  is *Collet-Eckmann* if the lower Lyapunov exponent of the critical value is bigger than zero:

$$\liminf \frac{\ln |Df^n(f(0))|}{n} > 0.$$

We say that  $f$  has subexponential recurrence if

$$\limsup \frac{-\ln |f^n(0)|}{n} = 0.$$

We say that  $f$  has polynomial recurrence if

$$\gamma = \limsup \frac{-\ln |f^n(0)|}{\ln(n)} < \infty,$$

and in this case, we call  $\gamma$  the *exponent* of the recurrence.

We introduce the following additional condition: we say that  $f$  is *Weakly Regular* if

$$(11.1) \quad \lim_{\delta \rightarrow 0} \liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{\substack{1 \leq k \leq n \\ f^k(0) \in (-\delta, \delta)}} \ln |Df(f^k(0))| = 0.$$

Notice that polynomial recurrence is much stronger than subexponential recurrence.

*Remark 11.1.* Maps satisfying the Collet-Eckmann and the subexponential recurrence conditions have been intensively studied after the works of Benedicks and Carleson. Those two conditions give a very precise control of the critical orbit. They are not sufficient to show that  $f$  has good statistical properties however: one must also ask that  $f$  has a renormalization with all periodic orbits repelling (and so is conjugate to a quadratic polynomial). Under this additional assumption, it is possible to show that  $f$  has an absolutely continuous invariant measure (see [BY]).

In order to study further the properties of  $\mu$ , it is convenient to consider the smallest periodic nice interval  $T$  of  $f$  ( $f$  is not infinitely renormalizable, since it has an absolutely continuous invariant measure). The first return map  $f^m : T \rightarrow T$  can be then rescaled to a unimodal map  $\hat{f}$ , which also possess an absolutely continuous invariant measure  $\hat{\mu}$ .

Assuming that  $f$  is also Kupka-Smale and using Lemma 5.2, we see that the dynamics of  $f$  splits in a hyperbolic part, that describes points  $x \in I$  which never enter  $\text{int } T$ , and an interesting part described by  $\hat{f}$ .

The measurable dynamics of  $\hat{f}$  are described by  $\hat{\mu}$ : for almost every  $x \in I$  and any continuous function  $\phi : I \rightarrow \mathbb{R}$  we have

$$\frac{1}{n} \sum_{k=0}^{n-1} \phi(\hat{f}^k(x)) = \int \phi d\hat{\mu}.$$

Since  $\hat{f}$  is non-renormalizable, it follows that  $\hat{\mu}$  is supported on  $[\hat{f}^2(0), \hat{f}(0)]$ , and  $(\hat{f}, \hat{\mu})$  is exponentially mixing (see [Y]).

The condition of Weak Regularity is important to show that  $(\hat{f}, \hat{\mu})$  is stochastically stable (see [T2]). If we assume a little bit more smoothness,  $f \in \mathbb{U}^3$ , the

Weak Regularity condition is not necessary, and it is possible to show that  $(\hat{f}, \hat{\mu})$  is stochastically stable in a stronger sense (see [BV]).

**11.2. Analytic families.** We will actually prove the following result, which is a more precise form of Corollaries C and E:

**Theorem 11.1.** *Let  $f_\lambda$  be a non-trivial analytic family of unimodal maps. Then almost every non-regular parameter is Kupka-Smale and has a quasiquadratic renormalization which satisfies the Collet-Eckmann condition and is polynomially recurrent with exponent 1.*

*Proof.* We will prove the stated result for one-parameter families, the general case reducing to this one by the argument of §10.1.1.

By Theorems A and B of [AM1], the conclusion of the theorem holds for the quadratic family. However, the only properties of the quadratic family that are actually used in the proof is that it is an analytic family of quasiquadratic maps with negative Schwarzian derivative for which the Phase-Parameter relation holds at almost every parameter, see Remark 3.3 of that paper. Due to the work of Kozłowski, the hypothesis of negative Schwarzian derivative can also be removed (this can be checked directly using Lemma 5.5). Using our Theorem A, we conclude that the conclusion also holds for analytic families of quasiquadratic maps.

Let us now consider the general case. By Theorem A, almost every non-regular parameter is simple, and by Theorem B, almost every non-regular parameter has a quasiquadratic renormalization. Let us fix such a parameter  $\lambda_0$ .

Let  $T$  be the smallest periodic nice interval for  $f_{\lambda_0}$  (of period  $m$ ). For  $\lambda$  near  $\lambda_0$ , the interval  $T$  has a continuation  $T[\lambda]$ . Consider the analytic family  $g_\lambda = A[\lambda] \circ f_\lambda^m \circ A[\lambda]^{-1}$ ,  $|\lambda - \lambda_0| < \epsilon$ , where  $A[\lambda] : T[\lambda] \rightarrow I$  is affine. Then  $g_\lambda$  is  $C^\infty$  close to  $g_{\lambda_0}$ , which is quasiquadratic, so we conclude that for  $\epsilon > 0$  small,  $g_\lambda$ ,  $|\lambda - \lambda_0| < \epsilon$  is an analytic family of quasiquadratic maps.

In particular, for almost every  $\lambda$  near  $\lambda_0$ ,  $g_\lambda$  is either regular or satisfy the Collet-Eckmann condition and its critical point is polynomially recurrent with exponent 1. In particular, the same holds for  $f_\lambda$ , which concludes the proof of the theorem.  $\square$

*Remark 11.2.* Notice that the proof of Theorem A in [AM2] could not use directly the proof of [AM1] (the argument needs modifications which are dealt in the Appendix of [AM2]), since their main phase-parameter tool essentially amounts to comparing the phase-space of a non-trivial family with the parameter space of the quadratic family. This distorts the estimates and makes it impossible to obtain the exponent of the recurrence.

### 11.3. Smooth families.

**Theorem 11.2.** *Let  $f_\lambda$ ,  $\lambda \in \Lambda$  be a non-trivial family of unimodal maps. For every  $\epsilon > 0$  there exists a neighborhood  $\mathcal{V} \subset \mathbb{UF}^2(\Lambda)$  of  $f_\lambda$  such that if  $g_\lambda \in \mathcal{V}$  then, outside a set of parameters  $\lambda$  of measure at most  $\epsilon$ ,  $g_\lambda$  is either regular or is Kupka-Smale and has a renormalization with all periodic orbits repelling and satisfies the Collet-Eckmann, subexponential recurrence and Weak Regularity conditions.*

*Proof.* Using Vitali's covering Lemma, let  $\{B_i\}$ ,  $\{C_i\}$  be finite families of disjoint closed balls covering the parameter space up to a set of Lebesgue measure  $\epsilon/2$  such that:

- (1) For  $\lambda \in B_i$ ,  $f_\lambda$  is regular;
- (2) For  $\lambda \in C_i$ , there exists a nice interval  $T_i[\lambda]$ , which is periodic of period  $m_i$ , depending continuously on  $\lambda$  such that  $f^{m_i} : T_i[\lambda] \rightarrow T_i[\lambda]$  can be rescaled to a quasiquadratic map  $\hat{f}_\lambda^i$ .

It is easy to see that if  $g_\lambda$  is  $C^2$  close to  $f_\lambda$ , then for every  $\lambda \in \cup B_i$   $g_\lambda$  is regular, and for every  $\lambda \in \cup C_i$ , there exists a continuously depending interval  $T_i^g[\lambda]$ , close to  $T_i^g[\lambda]$  such that  $g_\lambda^{m_i} : T_i^g[\lambda] \rightarrow T_i^g[\lambda]$  can be rescaled to a unimodal map  $\hat{g}_\lambda^i$ , and the family  $\hat{g}_i[\lambda]$  is  $C^2$  close to  $\hat{f}_\lambda^i$ .

The family  $\hat{f}_\lambda^i$  is non-trivial, so by Theorem B of [ALM], the set of parameters in  $C_i$  such that  $\hat{g}_\lambda^i$  is non-regular or fails to satisfy conditions (1) and (2) has Lebesgue measure less than  $|C_i|\epsilon/2$ , provided  $\mathcal{V}$  is small enough. The result follows.  $\square$

*Remark 11.3.* In particular, if  $f_\lambda$  is a non-trivial analytic family of unimodal maps, almost every parameter is Weakly Regular.

Recall that if  $\Lambda \in \mathbb{R}^k$  is an bounded open connected domain with smooth boundary,  $\mathbb{UF}^r(\Lambda)$  is the space of  $C^r$  families of unimodal maps parametrized by  $\Lambda$ , and is a Baire space. Recall also that by Remark 8.1, non-trivial analytic families are dense in  $\mathbb{UF}^r(\Lambda)$ . Using Theorem 11.2 and an easy Baire argument we obtain the following precise version of Corollary D:

**Theorem 11.3.** *In a generic family  $f_\lambda$  in  $\mathbb{UF}^r(\Lambda)$ ,  $r = 2, \dots, \infty$  for almost every non-regular parameter  $\lambda_0 \in \Lambda$ ,  $f = f_{\lambda_0}$  is Kupka-Smale and has a renormalization which has all periodic orbits repelling and satisfies the Collet-Eckmann, subexponential recurrence and Weak Regularity conditions.*

## APPENDIX A. HYBRID CLASSES

In this section we will give a global characterization of the leaves of the lamination  $\mathcal{L}$  of Theorem 8.1.

Notice that the leaves of  $\mathcal{L}$  are claimed to coincide with topological classes only in the non-regular case: the partition in topological classes is not a lamination because regular topological classes are open sets. It turns out that the behavior of the regular leaves of  $\mathcal{L}$  can be quite arbitrary. In order to give a global characterization of the leaves of  $\mathcal{L}$ , we need to introduce once and for all an arbitrary, but fixed, way to refine the topological classes of regular maps. We shall call this refinement the *hybrid lamination*.

If  $f$  is non-regular, the hybrid class of  $f$  is just the set of all non-regular maps  $g$  which are topologically conjugate to  $f$ .

Let  $f$  be a regular map, and let  $A$  be the set of attracting periodic orbits of  $f$  and let  $B = \{x \in I | f^n(x) \rightarrow A\}$  denote the basins of the attracting periodic orbits of  $f$ . Notice that if  $f$  is a regular map, there exists a minimal  $m \geq 0$  such that  $f^m(0)$  belongs to a periodic connected component of  $B$ . It is possible to show that if  $f$  is quasiquadratic, then  $m = 0$ . It turns out that if  $m = 0$  (this case will be called essential), there is a natural way to refine the topological class of  $f$ : the hybrid class of  $f$  is the set of all regular maps  $g$  which are topologically conjugate to  $f$  and the multiplier of the periodic orbit that attracts 0 is the same for both maps (this definition agrees with the one of [ALM] in the quasiquadratic case).

In the non-essential case, there is no natural way to refine the topological class of  $f$ , so we fix an arbitrary way that works.

**Definition A.1.** Let  $f$  be a regular map. We say that a homeomorphism  $h : I \rightarrow \mathbb{C}$  is  $f$ -admissible if the following holds. Let  $T$  be a periodic component of  $B \setminus A$  which does not contain 0, and, writing  $T = (a, b)$  with  $|a| < |b|$ , we have that the interval  $[-a, a]$  is nice. Then  $h$  takes  $d = (a+b)/2$  to  $h(d) = (h(a) + h(b))/2$  and  $h|_{[d, f^q(d)]}$  is affine, where  $q$  is the period of  $T$ .

In the non-essential case, we let the hybrid class of  $f$  be defined as the set of all regular maps  $g$  such that there exists a  $f$ -admissible topological conjugacy between  $f$  and  $g$ .

The following lemma is elementary, and shows that the definition of hybrid class is at least adequate:

**Lemma A.1.** *Let  $f$  be a regular map. Then its hybrid class intersects  $\mathcal{U}_a$  in a codimension-one analytic submanifold.*

With this definition, it is possible to prove Theorem 8.1 in the case of hyperbolic maps  $f$ . The case of infinitely renormalizable  $f$  can be dealt by reduction to the quasiquadratic case using renormalization (dealt in Theorem A of [ALM]), see Lemma 5.4.

We will now explain the relation of  $f$ -admissible map with the main tool of [ALM]: the concept of persistent puzzle.

**A.1. Persistent puzzle.** Assume that  $f \in \mathcal{F}$ . As in §6.1, fix a level  $\mathbf{n}$  of the principal nest and assume that  $|I_{\mathbf{n}}|/|I_{\mathbf{n}-1}|$  is very small. Let us consider the first landing map to  $A^0 = I_{\mathbf{n}}$ , the connected components of its domain are denoted  $A^j$ . Let  $A^1$  be the component of  $f(0)$ , and let  $A^1 = (l, r)$ , with  $r > 0$ . Let  $V^j$  be the complexification of the  $A^j$  obtained as in Lemma 6.1. Let  $V$  be the union of all  $V^j$  such that  $V^j \cap \mathbb{R} \subset [-1, r]$ . We shall informally call  $V$  the *puzzle*.

Let  $\mathcal{V} \subset \mathcal{A}_a$  be a real-symmetric neighborhood of  $f$ . We will say that the puzzle *persists* in  $\mathcal{V}$  if there exists a real-symmetric holomorphic motion  $h$  over  $\mathcal{V}$  given by a family of transition maps  $h[f, g] = h_g : \mathbb{C} \rightarrow \mathbb{C}$ ,  $g \in \mathcal{V}$  such that:

- (1)  $h_g|_{\mathbb{C} \setminus \Omega_a} = \text{id}$ ;
- (2)  $g \circ h_g|_{V \setminus V^0} = h_g \circ f$ ,  $g \circ h_g|_{\partial V^0} = f$ .
- (3)  $h_g|_I$  is  $f$ -admissible and  $g \circ h_g|_{[-1, r] \setminus V} = h_g \circ f$ .

The following plays the role of Lemma 5.6 of [ALM].

**Lemma A.2.** *Let  $f \in \mathcal{F} \cap \mathcal{U}_a$ . If  $|I_{\mathbf{n}}|/|I_{\mathbf{n}-1}|$  is sufficiently small, then there exists a neighborhood of  $f$  where the puzzle persists.*

The proof is the same as of Lemma 5.6 of [ALM]. It follows a sequence of steps:

(1) One considers a holomorphic motion  $h'$  of  $[-1, r] \setminus V$  which is  $f$ -admissible and equivariant:  $g \circ h'_g = h'_f \circ f$  (this holomorphic motion exists because the dynamics of  $f|_{[-1, r] \setminus V}$  is hyperbolic). Over a small neighborhood of  $f$ .

(2) Using the Canonical extension Lemma, we extend  $h'$  to a holomorphic motion defined also on  $\partial f(V_0)$ . Considering a slightly smaller neighborhood  $\mathcal{V}'$  of  $f$  we may extend  $h'$  to  $\mathbb{C} \setminus \Omega$  as  $\text{id}$ .

(3) One considers a holomorphic motion  $h^0$  of  $\overline{V}^0$  such that  $g \circ h_g^0|_{\partial V^0} = h' \circ f$  over a neighborhood  $\mathcal{V}^0$  of  $f$ .

(4) One notices that for each  $V^i$ ,  $i \neq 0$ , we can define uniquely a holomorphic motion  $h^i$  on  $V^i$  by a finite lift of  $h^0|_{V^0}$  over a small neighborhood  $\mathcal{V}^i$  of  $f$ .



(5) The (countably many) holomorphic motions  $h'$ ,  $h^i$  are defined apriori over different neighborhoods of  $f$ , but using again hyperbolicity of  $f|[-1, r] \setminus V$ , one sees that all those holomorphic motions are defined over a definite neighborhood of  $f$ .

(6) An estimate of hyperbolic geometry shows that the several regions of definition of those different holomorphic motions cannot collide in a slightly smaller neighborhood of  $f$ , so they define a common holomorphic motion which can be completed using the Canonical Extension Lemma and satisfies automatically (1), (2), and (3).

*Remark A.1.* The last condition of the definition of persistence defines uniquely  $h_g$  in  $[-1, r] \setminus \overline{V}$ . This set is empty in the quasiquadratic case (and so this condition does not appear in [ALM]). This (obvious) observation concerning the first step is the only formal difference in the proof, the remaining steps do not need to be modified.

*Remark A.2.* If  $f$  is a Kupka-Smale, finitely-renormalizable, non-hyperbolic map, with a non-recurrent critical point, a similar construction can be made. In this case, we take  $T \subset T'$  nice intervals with preperiodic boundary such that 0 does not return to  $T'$  and  $|T|/|T'|$  is very small. We let  $A^0 = T$ , and put  $A^1$  as a domain of the first landing map to  $A^0$  which is contained in  $[f(0), f(0) + \epsilon]$ ,  $\epsilon$  very small.

*Remark A.3.* If  $g_1, g_2 \in \mathcal{V} \cap \mathcal{U}_a$  are regular maps in the same hybrid class then they are of non-essential type if and only if for all  $m$  sufficiently big,

$$h_{g_1}^{-1}(g_1^m(0)), h_{g_2}^{-1}(g_2^m(0)) \notin [-1, r] \setminus \overline{V}$$

(use the Schwarz Lemma). The definition of hybrid class implies

$$h_{g_1}^{-1}(g_1^m(0)) = h_{g_2}^{-1}(g_2^m(0)).$$

This is important for the application of the several pullback arguments of [ALM].

One obtains Theorem 8.1 by repetition of the proof of Theorem A of [ALM], taking into consideration the above remarks.

## APPENDIX B. NON-RENORMALIZABLE PARAMETERS IN THE MANDELBROT SET

The aim of this appendix is to show how the idea of the proof of Theorem B can be coupled with Lyubich's result of [L3] to obtain the following theorem:

**Theorem B.1.** *Let  $\mathcal{NR}$  be the set of non-renormalizable quadratic parameters with recurrent critical point and no indifferent periodic orbits in the boundary of the Mandelbrot set. Then  $\mathcal{NR}$  has Lebesgue measure 0.*

This theorem implies easily the following result due to Shishikura:

**Theorem B.2.** *The set of parameters in the boundary of the Mandelbrot set which are not infinitely renormalizable has Lebesgue measure 0.*

*Remark B.1.* The reduction of Theorem B.2 to Theorem B.1 is obtained using the following three steps:

(1) It is easy to pass from the non-renormalizable case to the finitely renormalizable case using renormalization techniques: the (countably many) little copies of the Mandelbrot set are related by renormalization to the original Mandelbrot set by a quasiconformal (and thus absolutely continuous) transformation, see [L4]. Alternatively, we can also repeat the proofs for the little Mandelbrot copies.

(2) Quadratic polynomials with a neutral fixed points are contained in the boundary of the main cardioid of the Mandelbrot set, which is an analytic curve (with one singularity) and thus has Lebesgue measure zero.

(3) The case of non-recurrent non-renormalizable polynomial without neutral fixed points can be treated easily using holomorphic motions, see our proof of Lemma 9.4 (it is enough to use that under those conditions the set of points that never enter a small neighborhood of 0 is a hyperbolic set and thus persistent<sup>7</sup>).

To prove Theorem B.1 we will make use of the Complex Phase-Parameter relation (Theorem 3.1) and Lyubich's parapuzzle estimate (Theorem 4.3). Then, we will redo the estimates of Theorem B in the complex setting to show that non-renormalizable parameters have Lebesgue measure zero, because the critical point has a tendency to fall in the basin of infinity (in the same way that in the real setting the critical point has a tendency to fall in the basin of non-essential attractors).

*Remark B.2.* Lyubich has another proof of Theorem B.1, also based on [L3] and estimates on the area of the set of points that return to deep puzzle pieces.

**B.1. Parapuzzle notation.** Let us fix  $c_0 \in \mathcal{NR}$ . By Theorem 4.3, there exists a neighborhood  $\Lambda_1 \subset \mathbb{C}$  of  $c_0$  and domains  $0 \in U_1[\lambda] \subset \mathbb{C}$ ,  $\lambda \in \Lambda_1$  such that the first return map to  $U_1[\lambda]$  by  $p_\lambda$  induces a full  $\mathcal{R}$ -family over  $\Lambda_1$ .

To prove Theorem B.1, it is clearly sufficient to show that  $\Lambda_1 \cap \mathcal{NR}$  has Lebesgue measure zero.

For  $\lambda \in \mathcal{NR} \cap \Lambda_1$ , we can define a  $\mathcal{R}$ -chain over  $\lambda$  since the critical point is recurrent. Let us denote the parameter domains of this chain by  $\Lambda_i[\lambda]$ .

Let  $\mathcal{NR}^\infty \subset \mathcal{NR} \cap \Lambda_1$  be the set of parameters  $\lambda$  such that the chain  $\mathcal{R}_i$  over  $\lambda$  has infinitely many central levels, and let  $\mathcal{NR}^0$  be the complementary set in  $\mathcal{NR} \cap \Lambda_1$ .

By Theorem 4.4, there exists a constant  $C(\lambda) > 0$ ,  $\lambda \in \mathcal{NR} \cap \Lambda_1$  such that  $\text{mod}(\Lambda_{n_k}[\lambda] \setminus \overline{\Lambda_{n_k+1}[\lambda]}) > C(\lambda)k$ , where  $n_k - 1$  counts the non-central levels of the chain. If  $\lambda \in \mathcal{NR}^0$ , we actually have linear growth of moduli (without passing through a subsequence), and by Theorem 3.1, the complex phase-parameter relation holds.

**B.2. Finitely many central cascades.** The argument of Lyubich which shows that almost every real quadratic maps in  $\mathcal{F}$  is simple applies in the complex setting and gives:

**Lemma B.3.**  $|\mathcal{NR}^\infty| = 0$ .

*Proof.* Let  $\mathcal{NR}_\epsilon^\infty$  be the set of parameters  $\lambda \in \mathcal{NR}^\infty$  such that  $C(\lambda) \geq \epsilon$ . If  $\mathcal{NR}^\infty$  has positive Lebesgue measure then we can select  $\epsilon$  such that  $\mathcal{NR}_\epsilon^\infty$  also has positive Lebesgue measure. Let  $\mathcal{NR}_\epsilon^\infty(k) \subset \mathcal{NR}_\epsilon^\infty$  be the set of parameters such that the  $n_k$  level is central. If  $\lambda \in Z^k$ ,

$$\mathcal{NR}_\epsilon^\infty(k) \cap \Lambda_{n_k}[\lambda] \subset \Lambda_{n_k}^0[\lambda]$$

thus

$$|\mathcal{NR}_\epsilon^\infty(k) \cap \Lambda_{n_k}[\lambda]| \leq |\Lambda_{n_k}^0[\lambda]|.$$

---

<sup>7</sup>This actually holds for any non-renormalizable polynomial without neutral fixed points.

Since  $C(\lambda) \geq \epsilon$ , there exists  $\delta$  and  $k_0$  which only depend on  $\epsilon$  such that if  $k > k_0$  then  $\Lambda_{n_k}[\lambda] \setminus \overline{\Lambda_{n_k}^0[\lambda]}$  contains a round annuli of moduli  $k\delta$ . This implies that

$$|\Lambda_{n_k}^0[\lambda]| \leq e^{-k\delta'} |\Lambda_{n_k}[\lambda]|$$

for some  $\delta'$  depending on  $\delta$ . For each  $k$ , the domains  $\Lambda_{n_k}[\lambda]$ ,  $\lambda \in \mathcal{NR}_\epsilon^\infty(k)$  are either equal or disjoint, and their union has Lebesgue measure at most  $|\Lambda_1|$ , so  $|\mathcal{NR}_\epsilon^\infty(k)|$  decays exponentially on  $k$ . It follows immediately that

$$\mathcal{NR}_\epsilon^\infty = \cap_{k \geq 1} \cup_{n \geq k} \mathcal{NR}_\epsilon^\infty(k)$$

has Lebesgue measure zero, contradiction.  $\square$

**B.3. Area estimate.** Let  $U$  be a bounded open set of  $\mathbb{C}$  and  $Z$  be a measurable set of  $\mathbb{C}$ . Let

$$c_\gamma(Z|U) = \sup \frac{|h(Z \cap U)|}{|h(U)|}$$

where  $h$  ranges over all quasiconformal homeomorphisms  $h : U \rightarrow \mathbb{C}$  with dilatation bounded by  $\gamma$  and such that  $h(U)$  is bounded. The following two properties are immediate:

(1) If  $V^j \subset U$  are disjoint open subsets and  $Z \subset \cup V^j$  then

$$c_\gamma(Z|U) \leq \sup_j c_\gamma(Z|V^j) c_\gamma(\cup V^j|U).$$

(2) If  $A, B \subset U$  are disjoint open subsets and  $Z \subset A \cup B$  then

$$c_\gamma(Z|U) \leq c_\gamma(B|U) + (1 - c_\gamma(B|U)) c_\gamma(Z|B).$$

Denote by  $V_n^k[\lambda]$  the connected components of the preimages of

$$(R_{n-1}[\lambda]|U_n^0[\lambda])^{-1}(\cup U_{n-1}^j[\lambda]).$$

We reserve the index 0 for the component of 0, so that  $0 \in V_n^0$ . We also reserve the indexes  $-1$  and  $1$  for the components of the preimages of  $U_n[\lambda]$ .

Fix some  $\gamma > 1$ . Let

$$\epsilon_n(\lambda) = c_\gamma(\cup_{|k| \leq 1} V_n^k[\lambda]|U_n[\lambda])$$

$$\alpha_n(\lambda) = c_\gamma(\cup_j U_n^j[\lambda]|U_n[\lambda]).$$

**Lemma B.4.** *Let  $\lambda \in \mathcal{NR}^0$ . Then  $\alpha_2 < 1$ .*

*Proof.* Notice that  $\cup U_1^j[\lambda]$  is not dense in  $U_1[\lambda]$  (otherwise the filled-in Julia set of  $p_\lambda$  would have to contain  $U_1[\lambda]$ ). Thus, there exists a domain  $U_1^0[\lambda] \subset D[\lambda] \subset U_1[\lambda]$  such that  $U_1[\lambda] \setminus \overline{D[\lambda]}$  is an annulus, and a non-empty open set  $E[\lambda] \subset U_1[\lambda] \setminus \cup U_1^j[\lambda]$ . By the Koebe distortion Lemma, if  $h : U_1[\lambda] \rightarrow \mathbb{C}$  is a  $\gamma$ -qc map with bounded image then  $|h(E[\lambda])| > C|h(U_1^0[\lambda])|$  for some constant  $C > 0$ .

To each  $\underline{d} \in \Omega$  we associate  $E^{\underline{d}}[\lambda] = (R_1^{\underline{d}}[\lambda])^{-1}(E[\lambda])$ . We conclude that if  $h : U_1[\lambda] \rightarrow \mathbb{C}$  is a  $\gamma$ -qc map then  $|h(\cup E^{\underline{d}}[\lambda])| > C|h(\cup W_1^{\underline{d}}[\lambda])|$ , so  $c_\gamma(\cup W_1^{\underline{d}}[\lambda]|U_1[\lambda]) < 1$ .

If  $|k| > 1$  then  $R_1^2[\lambda]|V_2^k[\lambda]$  is a diffeomorphism onto  $U_1[\lambda]$  and we conclude that

$$c_\gamma(\cup U_2^j[\lambda]|V_2^k[\lambda]) = c_\gamma(\cup W_1^{\underline{d}}[\lambda]|U_1[\lambda]).$$

Thus

$$c_\gamma(\cup U_2^j[\lambda]|U_2[\lambda]) \leq \epsilon_2 + (1 - \epsilon_2) c_\gamma(\cup W_1^{\underline{d}}[\lambda]|U_1[\lambda]) < 1.$$

$\square$

**Lemma B.5.** *If  $\lambda \in \mathcal{NR}^0$  then  $\epsilon_n(\lambda) \rightarrow 0$  exponentially fast.*

*Proof.* Notice that if  $R_{n-1}[\lambda](V_n^k[\lambda]) = U_{n-1}^j[\lambda]$  then

$$\begin{aligned} \text{mod}(U_n[\lambda] \setminus \overline{V_n^k[\lambda]}) &\geq \text{mod}(U_{n-1}[\lambda] \setminus \overline{U_{n-1}^j[\lambda]})/3, \\ \text{mod}(U_{n-1}[\lambda] \setminus \overline{U_{n-1}^j[\lambda]}) &\geq \text{mod}(U_{n-2}[\lambda] \setminus \overline{U_{n-2}^0[\lambda]})/2. \end{aligned}$$

For  $\lambda \in \mathcal{NR}^0$ ,  $\text{mod}(U_{n-2}[\lambda] \setminus \overline{U_{n-2}^0[\lambda]})$  grows linearly in  $n$ , so  $\inf_k \text{mod}(U_n[\lambda] \setminus \overline{V_n^k[\lambda]})$  also grows linearly, and this implies exponential decay of  $\sup_k c_\gamma(V_n^k[\lambda]|U_n[\lambda])$ , which implies exponential decay of  $\epsilon_n$ .  $\square$

**Lemma B.6.** *If  $\lambda \in \mathcal{NR}^0$  then  $\alpha(\lambda) = \sup_{n \geq 2} \alpha_n(\lambda) < 1$ .*

*Proof.* Indeed, if  $|k| > 1$  then  $R_n^2[\lambda]|V_{n+1}^k[\lambda]$  is a diffeomorphism onto  $U_n[\lambda]$ . In particular,

$$c_\gamma(\cup U_{n+1}^j[\lambda]|V_{n+1}^k[\lambda]) \leq c_\gamma(\cup U_n^j[\lambda]|U_n[\lambda]) = \alpha_n(\lambda).$$

Thus

$$c_\gamma(\cup U_{n+1}^j[\lambda]|U_{n+1}[\lambda] \setminus \overline{\cup_{|k| \leq 1} V_{n+1}^k[\lambda]}) \leq \alpha_n(\lambda)$$

which implies

$$\alpha_{n+1}(\lambda) \leq \epsilon_n(\lambda) + (1 - \epsilon_n(\lambda))\alpha_n(\lambda)$$

and

$$1 - \alpha_{n+1}(\lambda) \geq (1 - \epsilon_n(\lambda))(1 - \alpha_n(\lambda)).$$

If  $\lambda \in \mathcal{NR}^0$ ,  $\epsilon_n(\lambda)$  decays exponentially (Lemma B.5) and  $\alpha_2(\lambda) < 1$  (Lemma B.4), so the result follows.  $\square$

If  $\mathcal{NR}^0$  has positive measure, there exists  $\alpha > 0$ ,  $k > 0$  and a positive measure set  $X$  such that for  $\lambda \in X$ ,  $\alpha(\lambda) < \alpha$  and for  $n > k$  the estimate CPhPa1 of the Complex Phase-Parameter relation is valid with a constant smaller than  $\gamma$ .

Let  $Y \supset X$  be an open set such that  $\alpha|Y| < |X|$ . For every parameter  $\lambda \in X$ , let  $\mu(\lambda)$  be the smallest  $j > k$  such that  $\lambda \in Z[\lambda] = \Lambda_j^{\tau_j(\lambda)}[\lambda] \subset Y$  (such a  $j$  exists since  $\cap \Lambda_j[\lambda] = \{\lambda\}$ ). The resulting collection of parameter domains  $Z[\lambda]$ ,  $\lambda \in X$  are either disjoint or equal. To reach a contradiction, it is enough to show that  $\alpha|Z[\lambda]| \geq |X \cap Z[\lambda]|$ , for in this case  $\alpha|Y| \geq |X|$ . But this is an immediate consequence of CPhPa1, for

$$\frac{|X \cap Z[\lambda]|}{|Z[\lambda]|} \leq c_\gamma(\cup W_{\mu(\lambda)}^d |U_{\mu(\lambda)}^{\tau_{\mu(\lambda)}(\lambda)}) \leq c_\gamma(\cup U_{\mu(\lambda)}^j |U_{\mu(\lambda)}) \leq \alpha,$$

since  $\tau_{\mu(\lambda)} \neq 0$  by hypothesis (notice that we even have  $|\mathcal{M} \cap Z[\lambda]|/|Z[\lambda]| \leq \alpha$ , that is, a definite proportion of parameters in  $Z[\lambda]$  have escaping critical point).

## REFERENCES

- [Av1] A. Avila. Bifurcations of unimodal maps: the topologic and metric picture. IMPA Thesis (2001) (<http://www.math.sunysb.edu/~artur/>).
- [Av2] A. Avila. Bifurcations d'applications unimodales. To appear in Comptes Rendus Mathematiques (<http://www.math.sunysb.edu/~artur/>).
- [ALM] A. Avila, M. Lyubich and W. de Melo. Regular or stochastic dynamics in real analytic families of unimodal maps. Preprint (<http://www.math.sunysb.edu/~artur/>).
- [AM1] A. Avila, C. G. Moreira. Statistical properties of unimodal maps: the quadratic family. Preprint (<http://www.arXiv.org>).
- [AM2] A. Avila and C. G. Moreira. Statistical properties of unimodal maps: smooth families with negative Schwarzian derivative. Preprint (<http://www.arXiv.org>).

- [AM3] A. Avila and C. G. Moreira. Statistical properties of unimodal maps: physical measures, periodic orbits and pathological laminations. In preparation.
- [BV] V. Baladi and M. Viana. Strong stochastic stability and rate of mixing for unimodal maps. *Ann. scient. Éc. Norm. Sup.*, v. 29 (1996), 483-517.
- [BY] M. Benedicks and L.-S. Young. Absolutely continuous invariant measures and random perturbations for certain one-dimensional maps. *Ergodic Theory Dynam. Systems* 12 (1992), no. 1, 13-37.
- [BR] L. Bers & H.L. Royden. Holomorphic families of injections. *Acta Math.*, v. 157 (1986), 259-286.
- [D] A. Douady. Prolongement de mouvements holomorphes (d'après Ślodkowski et autres). *Asterisque*, v. 227, 7-20.
- [DH1] A. Douady & J.H. Hubbard. On the dynamics of polynomial-like maps. *Ann. Sc. Éc. Norm. Sup.*, v. 18 (1985), 287-343.
- [DH2] A. Douady and J. H. Hubbard. Étude dynamique des polynômes complexes. Parties I et II. *Publications Mathématiques d'Orsay*, 84-2 & 85-4.
- [GSS] J. Graczyk, D. Sands and G. Świątek. Schwarzian derivative in unimodal dynamics. *C. R. Acad. Sci. Paris*.
- [K1] O. S. Kozlovski. Getting rid of the negative Schwarzian derivative condition. *Ann. of Math.* (2) 152 (2000), no. 3, 743-762.
- [K2] O.S. Kozlovsky. Structural stability in one-dimensional dynamics. Thesis (1998).
- [L1] M. Lyubich. Combinatorics, geometry and attractors of quasi-quadratic maps. *Ann. Math.* **140** (1994), 347-404. A remark on parabolic towers. Manuscript 1999.
- [L2] M. Lyubich. Dynamics of quadratic polynomials, I-II. *Acta Math.*, **178** (1997), 185-297.
- [L3] M. Lyubich. Dynamics of quadratic polynomials, III. Parapuzzle and SBR measure. *Asterisque*, v. 261 (2000), 173 - 200.
- [L4] M. Lyubich. Feigenbaum-Coullet-Tresser Universality and Milnor's Hairiness Conjecture. *Ann. Math.* v. 149 (1999), 319 - 420.
- [L5] M. Lyubich. Almost every real quadratic map is either regular or stochastic. Preprint IMS at Stony Brook, # 1997/8. To appear in *Ann. Math.*
- [M] R. Mañé. Hyperbolicity, sinks and measures for one-dimensional dynamics. *Comm. Math. Phys.* 100 (1985), 495-524.
- [MSS] R. Mañé, P. Sad & D. Sullivan. On the dynamics of rational maps, *Ann. scient. Ec. Norm. Sup.*, **16** (1983), 193-217.
- [MS] W. de Melo & S. van Strien. One-dimensional dynamics. Springer, 1993.
- [MT] J. Milnor & W. Thurston. On iterated maps of the interval. "Dynamical Systems", *Proc. U. Md.*, 1986-87, ed. J. Alexander, *Lect. Notes Math.*, v. 1342 (1988), 465-563.
- [NS] T. Nowicki and D. Sands. Non-uniform hyperbolicity and universal bounds for  $S$ -unimodal maps. *Invent. Math.* 132 (1998), no. 3, 633-680.
- [S] D. Sands. Misiurewicz maps are rare. *Comm. Math. Phys.* 197 (1998), no. 1, 109-129.
- [S1] Z. Ślodkowski. Holomorphic motions and polynomial hulls. *Proc. Amer. Math. Soc.*, **111** (1991), 347-355.
- [T1] M. Tsujii. Positive Lyapunov exponents in families of one dimensional dynamical systems. *Invent. Math.* 111 (1993), 113-137.
- [T2] M. Tsujii. Small random perturbations of one dimensional dynamical systems and Margulis-Pesin entropy formula. *Random & Comput. Dynamics*. Vol.1 No.1 59-89, (1992).
- [Y] L.-S. Young. Statistical properties of dynamical systems with some hyperbolicity. *Ann. Math.* (2) 147, No.3, 585-650 (1998).

COLLÈGE DE FRANCE – 3 RUE D'ULM, 75005 PARIS – FRANCE.

*E-mail address:* avila@impa.br