



Unknotting Number One Knots are Prime: A New Proof

Xingru Zhang

Proceedings of the American Mathematical Society, Vol. 113, No. 2 (Oct., 1991),
611-612.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9939%28199110%29113%3A2%3C611%3AUNOKAP%3E2.0.CO%3B2-3>

Proceedings of the American Mathematical Society is currently published by American Mathematical Society.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

UNKNOTTING NUMBER ONE KNOTS ARE PRIME: A NEW PROOF

XINGRU ZHANG

(Communicated by Frederick R. Cohen)

ABSTRACT. An alternative proof for unknotting number one knots being prime is given.

The unknotting number of a knot $K \subset S^3$, denoted by $u(K)$, is the minimum number of crossing changes required to unknot K . Obviously $u(K)$ is a knot invariant, but surprisingly little is known about it. The following theorem of M. Scharlemann proves a long standing conjecture.

Theorem [S, Theorem]. *A knot $K \subset S^3$ with $u(K) = 1$ is prime.*

To prove the above theorem, Scharlemann developed certain combinatorics dealing with planar graphs coming from an intersection of two special planar surfaces. Later in [ST], Scharlemann and Thompson gave another proof of the theorem [ST, Corollary 3.4], that is based on a delicate application of the sutured manifold structure theory. In this note we point out a new proof, applying only some existing results. In fact the proof follows immediately from the following three known lemmas.

Lemma 1 [L, Lemma 1]. *Let K be a knot in S^3 with $u(K) = 1$, and let M_K be the double cover of S^3 branched over K . Then M_K can be obtained by $n/2$ -surgery on some knot in S^3 , n being an odd integer.*

Lemma 2 [GL, Theorem 1]. *Let K be a knot in S^3 , and let $K(m/l)$ denote the manifold obtained by m/l -surgery on K . Then $K(m/l)$ is a prime manifold if $|l| \neq 1$.*

Lemma 3 [KT, Corollary 4]. *Let K be a knot in S^3 . Then the double cover M_K of S^3 branched over K is a prime manifold iff K is a prime knot.*

Proof of Theorem. Since $u(K) = 1$, M_K is a prime manifold by Lemmas 1 and 2. Hence K is a prime knot by Lemma 3.

The author has learned that a similar approach was pointed out by C. Gordon in a lecture given at Santa Barbara.

Received by the editors June 5, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 57M25.

©1991 American Mathematical Society
 0002-9939/91 \$1.00 + \$.25 per page

REFERENCES

- [GL] C. McA. Gordon and J. Luecke, *Only integral Dehn surgery can yield reducible manifolds*, Math. Proc. Cambridge Philos. Soc. **102** (1987), 97–101.
- [KT] P. K. Kim and J. L. Tollefson, *Splitting the PL involutions of nonprime 3-manifolds*, Michigan Math. J. **27** (1980), 259–274.
- [L] W. B. R. Lickorish, *The unknotting number of a classical knot*, Contemp. Math., vol. 44, 1985, pp. 117–121.
- [S] M. G. Scharleemann, *Unknotting number one knots are prime*, Invent. Math. **82** (1985), 37–55.
- [ST] M. G. Scharleemann and A. Thompson, *Unknotting number, genus, and companion tori*, Math. Ann. **280** (1988), 191–205.

MATHEMATICS DEPARTMENT, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BRITISH COLUMBIA, CANADA, V6T 1Y4