

An infinite crowd of mathematicians enters a bar. The first one orders a pint, the second one a half pint, the third one a quarter pint... "I understand", says the bartender - and pours two pints.



MAT 132

8.7 Taylor and Maclaurin Series
8.8 Applications of Taylor Polynomials

Suppose we have a function defined by a series

$$f(x) = c_0 + c_1 x + c_2 x^2 \dots \text{ with convergence radius } R > 0.$$

- We know that f has derivatives of all orders on $(-R, R)$. Find those derivatives.
- Can you determine the values of the coefficients c_n in terms of f ?
- If $f(x) = e^x$, find an expression of f as a power series. What is the radius of convergence?

Suppose we have a function defined by a series

$$f(x) = c_0 + c_1 (x - a) + c_2 (x - a)^2 \dots \text{ with convergence radius } R > 0.$$

- We know that f has derivatives of all orders on $(a - R, a + R)$. Find those derivatives.
- Can you determine the values of the coefficients c_n in terms of f ?
- If $f(x) = \sin(x)$, can you find an expression of f as a power series centered at $\pi/2$?

5 Theorem If f has a power series representation (expansion) at a , that is, if

$$f(x) = \sum_{n=0}^{\infty} c_n (x - a)^n \quad |x - a| < R$$

then its coefficients are given by the formula

$$c_n = \frac{f^{(n)}(a)}{n!}$$

The series $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$

$$= f(a) + \frac{f'(a)}{1!} (x - a) + \frac{f''(a)}{2!} (x - a)^2 + \frac{f'''(a)}{3!} (x - a)^3 + \dots$$

is called the Taylor series of f at a .

When $a=0$, the series (below) is called the Maclaurin series

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = f(0) + \frac{f'(0)}{1!} x + \frac{f''(0)}{2!} x^2 + \dots$$

- The powers series representation of a function a given "a" is unique.
- If f is a function and we know that f has a representation as a power series, then the Taylor series converges to f in the interval of convergence.
- If we are given a function f (we don't know whether f has a representation as a power series,) the theorem below gives necessary conditions that imply that f has a representation as a power series:

$$T_n(x) = \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} (x - a)^i$$

$$= f(a) + \frac{f'(a)}{1!} (x - a) + \frac{f''(a)}{2!} (x - a)^2 + \dots + \frac{f^{(n)}(a)}{n!} (x - a)^n$$

$$R_n(x) = f(x) - T_n(x)$$

8 Theorem If $f(x) = T_n(x) + R_n(x)$, where T_n is the n th-degree Taylor polynomial of f at a and

$$\lim_{n \rightarrow \infty} R_n(x) = 0$$

for $|x - a| < R$, then f is equal to the sum of its Taylor series on the interval $|x - a| < R$.

8 Theorem If $f(x) = T_n(x) + R_n(x)$, where T_n is the n th-degree Taylor polynomial of f at a and

$$\lim_{n \rightarrow \infty} R_n(x) = 0$$

for $|x - a| < R$, then f is equal to the sum of its Taylor series on the interval $|x - a| < R$.

$$T_n(x) = \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} (x - a)^i$$

$$= f(a) + \frac{f'(a)}{1!} (x - a) + \frac{f''(a)}{2!} (x - a)^2 + \dots + \frac{f^{(n)}(a)}{n!} (x - a)^n$$

$$R_n(x) = f(x) - T_n(x)$$

9 Taylor's Inequality If $|f^{(n+1)}(x)| \leq M$ for $|x - a| \leq d$, then the remainder $R_n(x)$ of the Taylor series satisfies the inequality

$$|R_n(x)| \leq \frac{M}{(n+1)!} |x - a|^{n+1} \quad \text{for } |x - a| \leq d$$

This implies that the Taylor series converges in this case


- Find the Taylor series for $f(x)=e^x$ at $a=-3$.
 - Prove that $f(x)$ is equal to the series.
 - Use the series to approximate $e^{-3.1}$ correct to four decimal places.(you are given e^{-3})
- Find the Maclaurin series for $f(x)=\cos(x)$ and prove the series represents $f(x)$ for all values of x .
- Find the Maclaurin series for $f(x)=\sin(x)$ and prove the series represents $f(x)$ for all values of x .

REVIEW The alternating series test

If we have a sequence $\{a_n\}$, where $a_n > 0$, $a_n \geq a_{n+1}$, and $a_n \rightarrow 0$ when $n \rightarrow \infty$ then the series

$$\sum_{n=1}^{\infty} (-1)^n a_n$$

converges



Alternating Series Estimation Theorem If $s = \sum (-1)^n b_n$ is the sum of an alternating series that satisfies

$$(i) b_{n+1} \leq b_n \quad \text{and} \quad (ii) \lim_{n \rightarrow \infty} b_n = 0$$

$$\text{then} \quad |R_n| = |s - s_n| \leq b_{n+1}$$

- Find the Maclaurin series for $f(x)=(1+x)^k$, where k is any real number.

17 The Binomial Series If k is any real number and $|x| < 1$, then

$$(1+x)^k = \sum_{n=0}^{\infty} \binom{k}{n} x^n = 1 + kx + \frac{k(k-1)}{2!} x^2 + \frac{k(k-1)(k-2)}{3!} x^3 + \dots$$

- Prove that the binomial series converges when $|x| < 1$.
- Use the binomial series to expand $1/(1+x)^{0.5}$, as a power series. Use this to estimate $1/\sqrt{1.01}$ correct to three decimal places.

$$1 - \frac{1}{2}x + \frac{1 \cdot 3}{2^2 \cdot 2!} x^2 - \frac{1 \cdot 3}{2^3 \cdot 3!} x^3 + \dots + (-1)^n \frac{1 \cdot 3 \dots (2n-1)}{2^n \cdot n!} x^n + \dots$$

8.8.1

- Find the Taylor polynomials of $\cos(x)$ degree up to 6 centered at $a=0$.
- Evaluate these polynomials at $x=0, 0.1, \pi/4, \pi/2$, and π .
- Comment on how the polynomials converge to $\cos(x)$.

- 8.7.50 Use series to approximate the definite integral within the indicated accuracy

$$\int_0^{0.5} x^2 e^{-x^2} dx$$

correct to four decimal places.

- Use series to evaluate the limit

$$\lim_{x \rightarrow 0} \frac{\sin(x) - x + \frac{1}{6} x^3}{x^5}$$

1/120

$$\begin{aligned} \frac{1}{1-x} &= \sum_{n=0}^{\infty} x^n = 1 + x + x^2 + x^3 + \dots & R = 1 \\ e^x &= \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots & R = \infty \\ \sin x &= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots & R = \infty \\ \cos x &= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots & R = \infty \\ \tan^{-1} x &= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots & R = 1 \\ \ln(1+x) &= \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots & R = 1 \\ (1+x)^k &= \sum_{n=0}^{\infty} \binom{k}{n} x^n = 1 + kx + \frac{k(k-1)}{2!} x^2 + \frac{k(k-1)(k-2)}{3!} x^3 + \dots & R = 1 \end{aligned}$$

Recall, given a function f , and a number a , we define

$$T_n(x) = \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} (x-a)^i$$
$$= f(a) + \frac{f'(a)}{1!} (x-a) + \frac{f''(a)}{2!} (x-a)^2 + \cdots + \frac{f^{(n)}(a)}{n!} (x-a)^n$$

- We want to approximate $f(x)$ by $T_n(x)$ (for some n). Thus we need to know.
- For a given n , how good is our approximation?
- If we want the "error" to be less than a given number, how large does n have to be?

The remainder ("error") is $R_n(x) = f(x) - T_n(x)$.

We saw two methods for estimating $|R_n(x)|$ (each needs hypothesis)

- If the series is alternating, by the Alternating Series Estimation Theorem.
- If $|f^{(n)}(x)| \leq M$, then $|R_n(x)| \leq \frac{M}{(n+1)!} |x-a|^{n+1}$ for $|x-a| \leq d$

- Evaluate the integral below, correct up to two decimal places.

$$\int_{\frac{1}{2}}^1 \frac{\sin(x)}{x} dx$$