

MATH 533, SPRING 2020, HW6

DUE IN CLASS, MARCH 9

**Problem 1.** Let  $\mu, \nu$  be Radon measures on  $X, Y$ , not necessarily  $\sigma$ -finite. If  $f$  is a nonnegative l.s.c. function on  $X \times Y$ , show that  $x \rightarrow \int f_x d\nu$  and  $y \rightarrow \int f^y d\mu$  are Borel measurable and  $\int f d(\mu \hat{\times} \nu) = \iint f d\mu d\nu = \iint f d\nu d\mu$ .

**Problem 2.** The following gives an example of a smooth function not equal to its Taylor expansion at 0. Let  $f(t) = e^{-1/t}$  for  $t > 0$ ,  $f(t) = 0$  for  $t \leq 0$ . Check that

- (1) For  $k \in \mathbb{N}$  and  $t > 0$ ,  $f^{(k)}(t) = P_k(1/t)e^{-1/t}$  where  $P_k$  is a polynomial.
- (2)  $f^{(k)}(0)$  exists and is equal to 0 for all  $k$ .

**Problem 3.** If  $f \in L^\infty$  and  $\|f^y - f\|_\infty \rightarrow 0$  as  $y \rightarrow 0$ , then  $f$  agrees a.e. with a uniformly continuous function.