

MATH 533, SPRING 2020, HW7

DUE IN CLASS, APRIL 6

**Problem 1.** Suppose  $f \in L^p(\mathbb{R})$ . If there exists  $h \in L^p(\mathbb{R})$  such that

$$\lim_{y \rightarrow 0} \|y^{-1}(f^{-y} - f) - h\|_p = 0,$$

we call  $h$  the *strong  $L^p$  derivative* of  $f$  and write  $h = df/dx$ . If  $f \in L^p(\mathbb{R}^n)$ ,  $L^p$  derivatives of  $f$  are defined similarly. If  $p$  and  $q$  are conjugate exponents,  $f \in L^p$ ,  $g \in L^q$ , and the  $L^p$  derivative  $\partial_j f$  exists, then prove  $\partial_j(f * g)$  exists in the ordinary sense and equals  $(\partial_j f) * g$ .

**Problem 2.** Let  $\phi \in L^1(\mathbb{R}^n)$  satisfy  $|\phi(x)| \leq C(1+|x|)^{-n-\epsilon}$  for some  $C, \epsilon > 0$ , and  $\int \phi(x)dx = a$ . For  $t > 0$ ,  $\phi_t(x) = t^{-n}\phi(\frac{x}{t})$ . If  $f \in L^p$  define the  $\phi$ -maximal function of  $f$  to be  $M_\phi f(x) = \sup_{t>0} |f * \phi_t(x)|$ . The *Hardy-Littlewood maximal function*  $Hf$  is  $M_\phi|f|$  where  $\phi$  is the characteristic function of the unit ball, divided by the volume of the ball. Show that there is a constant  $C$ , independent of  $f$ , such that  $M_\phi f \leq CHf$ .

**Problem 3.** Young's inequality shows that  $L^1$  is a Banach algebra with convolution as multiplication.

- (1) If  $\mathcal{I}$  is an ideal in the algebra  $L^1$ , prove that its closure is, also.
- (2) If  $f \in L^1$ , the smallest closed ideal in  $L^1$  containing  $f$  is the smallest closed subspace of  $L^1$  containing translates of  $f$ .

**Problem 4.** Show that if  $f \in L^1(\mathbb{R}^n)$ ,  $f$  is continuous at 0, and  $\hat{f} \geq 0$ , then  $\hat{f} \in L^1$ .

**Problem 5.** Let  $f$  be a function on  $\mathbb{T}^1$  and  $A_r f$  the  $r$ th Abel mean of the Fourier series of  $f$ . Check that

- (1)  $A_r f = f * P_r$  where  $P_r(x) = \sum_{-\infty}^{\infty} r^{|k|} e^{2\pi i k x}$  is the Poisson kernel for  $\mathbb{T}^1$ .

- (2)  $P_r(x) = \frac{1-r^2}{1+r^2-2r \cos 2\pi x}$ .

**Problem 6.** Given  $f \in L^1(\mathbb{T}^1)$ , let  $S_m f(x) = \sum_{-m}^m \hat{f}(k) e^{2\pi i k x}$  and

$$\sigma_m f(x) = \sum_{-m}^m \hat{f}(k) \left(1 - \frac{|k|}{m+1}\right) e^{2\pi i k x}.$$

Prove the following.

- (1)  $\sigma_m f = \frac{1}{m+1} \sum_0^m S_k f$ .
- (2) If  $D_k$  is the  $k$ th Dirichlet kernel, we have  $\sigma_m f = f * F_m$  where  $F_m = \frac{1}{m+1} \sum_0^m D_k$ .  $F_m$  is the  $m$ th Fejér kernel on  $\mathbb{T}^1$ .
- (3)  $F_m(x) = \frac{\sin^2(m+1)\pi x}{(m+1)\sin^2 \pi x}$ .

**Problem 7.** Prove the following.

- (1) If  $D_m$  is the  $m$ th Dirichlet kernel,  $\|D_m\|_1 \rightarrow \infty$  as  $m \rightarrow \infty$ .
- (2) The Fourier transform is not surjective from  $L^1(\mathbb{T}^1)$  to  $C_0(\mathbb{Z})$ .