

Algebra & Number Theory

Volume 16
2022
No. 6

Saturation bounds for smooth varieties

Lawrence Ein, Huy Tài Hà and Robert Lazarsfeld

Saturation bounds for smooth varieties

Lawrence Ein, Huy Tài Hà and Robert Lazarsfeld

We prove bounds on the saturation degrees of homogeneous ideals (and their powers) defining smooth complex projective varieties. For example, we show that a classical statement due to Macaulay for zero-dimensional complete intersection ideals holds for any smooth variety. For curves, we bound the saturation degree of powers in terms of the regularity.

Introduction

We prove some saturation bounds for the ideals of nonsingular complex projective schemes and their powers.

We begin with some background. Consider the polynomial ring $S = \mathbb{C}[x_0, \dots, x_r]$ in $r + 1$ variables, and fix homogeneous polynomials

$$f_0, f_1, \dots, f_p \in S \quad \text{with} \quad \deg(f_i) = d_i.$$

We assume that $d_0 \geq d_1 \geq \dots \geq d_p$, and we denote by

$$J = (f_0, f_1, \dots, f_p) \subseteq S$$

the ideal that the polynomials span. Suppose now that J is primary for the irrelevant maximal ideal $\mathfrak{m} = (x_0, \dots, x_r)$ or equivalently that $\dim_{\mathbb{C}} S/J < \infty$. In this case J contains all monomials of sufficiently large degree, and it is a classical theorem of Macaulay [Carlson et al. 2003, Theorem 7.4.1] that

$$J_t = S_t \quad \text{for} \quad t \geq d_0 + \dots + d_r - r. \quad (1)$$

Moreover, this bound is (always) sharp when $p = r$. Although less well known, a similar statement holds for powers of J :

$$(J^a)_t = S_t \quad \text{for} \quad t \geq ad_0 + d_1 + \dots + d_r - r. \quad (2)$$

This again is always sharp when $p = r$.

It is natural to ask whether there are analogous results for more general homogeneous ideals J , in particular when

$$X =_{\text{def}} \text{Zeroes}(J) \subseteq \mathbb{P}^r$$

Ein partially supported by NSF grant DMS-1801870. Hà partially supported by Louisiana BOR grant LEQSF(2017-19)-ENH-TR-25. Lazarsfeld partially supported by NSF grant DMS-1739285.

MSC2020: 13B02, 14F99.

Keywords: saturation degree, symbolic powers of ideals, Castelnuovo–Mumford regularity.

is a smooth complex projective scheme. Of course if J has nontrivial zeroes, then it does not contain any power of the maximal ideal. However, if one interprets (1) and (2) as saturation bounds, then the question makes sense more generally. Specifically, recall that the *saturation* of a homogeneous ideal J is defined by

$$J^{\text{sat}} = \{f \in S \mid \mathfrak{m}^k \cdot f \subseteq J \text{ for some } k \geq 0\}.$$

The quotient J^{sat}/J has finite length, and in particular,

$$(J^{\text{sat}})_t = J_t \quad \text{for } t \gg 0.$$

The least such integer t is called the *saturation degree* $\text{sat. deg}(J)$ of J . Observing that $J^{\text{sat}} = S$ if and only if J is \mathfrak{m} -primary, statements (1) and (2) are equivalent to estimates for the saturation degrees of J and J^a . So the problem becomes to bound the saturation degree of an ideal in terms of the degrees of its generators.

It is instructive to consider some examples. Let $X \subseteq \mathbb{P}^r$ be a hyperplane defined by a linear form $\ell \in S$, and set

$$f_i = x_i^{d-1} \cdot \ell \quad \text{and} \quad J = (f_0, \dots, f_r) \subseteq S. \quad (3)$$

Then $J^{\text{sat}} = (\ell)$, and it follows from Macaulay's theorem that

$$\text{sat. deg}(J) = (r+1)(d-1) - r + 1 = (r+1)d - 2r,$$

which is very close to the bound (1). On the other hand, it is not the case that the saturation degree of an arbitrary ideal is bounded linearly in the degrees of its generators. For instance, the ideals

$$J = (x^d, y^d, xz^{d-1} - yw^{d-1}) \subseteq \mathbb{C}[x, y, z, w]$$

considered by Caviglia [2004, Example 4.2.1] have $\text{sat. deg}(J) \approx d^2$.

Our first main result asserts that for ideals defining smooth varieties, the Macaulay bounds remain true without modification.

Theorem A. *As above, suppose that*

$$J = (f_0, f_1, \dots, f_p) \subseteq S$$

is generated by forms of degrees $d_0 \geq \dots \geq d_p$, and assume that the projective scheme

$$X =_{\text{def}} \text{Zeroes}(J) \subseteq \mathbb{P}^r$$

cut out by the f_i is nonsingular. Then $\text{sat. deg}(J) \leq d_0 + \dots + d_r - r$, and more generally

$$\text{sat. deg}(J^a) \leq ad_0 + d_1 + \dots + d_r - r. \quad (4)$$

(If $p < r$, one takes $d_{p+1} = \dots = d_r = 0$.) We do not know whether the stated bound is best possible, but in any event it is asymptotically sharp. Indeed, if J is the ideal considered in (3), then Theorem A predicts that $\text{sat. deg}(J^a) \leq (a+r)d - r$, whereas in fact $\text{sat. deg}(J^a) = (a+r)d - 2r$.

Given a reduced algebraic set $X \subseteq \mathbb{P}^r$ denote by $I_X \subseteq S$ the saturated homogeneous ideal of X . Recall that the *symbolic powers* of I_X are

$$I_X^{(a)} = \{f \in S \mid \text{ord}_x(f) \geq a \text{ for general (or every) } x \in X\}.$$

Evidently $I_X^a \subseteq I_X^{(a)}$, and there has been a huge amount of interest in recent years in understanding the connections between actual and symbolic powers; see [Ein et al. 2001; Hochster and Huneke 2002; Bocci and Harbourne 2010; Dao et al. 2018]. If X is nonsingular, then $I_X^{(a)} = (I_X^a)^{\text{sat}}$. Therefore, Theorem A implies:

Corollary B. *Assume that $X \subseteq \mathbb{P}^r$ is smooth, and that I_X is generated in degrees $d_0 \geq d_1 \geq \cdots \geq d_p$. Then*

$$(I_X^{(a)})_t = (I_X^a)_t \quad \text{for } t \geq ad_0 + d_1 + \cdots + d_r - r.$$

For example, suppose that $X \subseteq \mathbb{P}^2$ consists of the three coordinate points, so that $I_X = (xy, yz, zx) \subseteq \mathbb{C}[x, y, z]$. Corollary B guarantees that I_X^a and $I_X^{(a)}$ agree in degrees $\geq 2a + 2$, whereas in reality $\text{sat. deg}(I_X^a) = 2a$. So here again the statement is asymptotically but not precisely sharp.

In the case of finite sets, results of Geramita, Gimigliano and Pitteloud [Geramita et al. 1995], Chandler [1997] and Sidman [2002] provide an alternative bound that is often best-possible. Recall that a scheme $X \subseteq \mathbb{P}^r$ is said to be m -regular in the sense of Castelnuovo–Mumford if its ideal sheaf $\mathcal{I}_X \subseteq \mathcal{O}_{\mathbb{P}^r}$ satisfies the vanishings

$$H^i(\mathbb{P}^r, \mathcal{I}_X(m - i)) = 0 \quad \text{for } i > 0.$$

This is equivalent to asking that I_X be generated in degrees $\leq m$, that the first syzygies among minimal generators of I_X appear in degrees $\leq m + 1$, the second syzygies in degrees $\leq m + 2$, and so on.¹ The authors just cited show that if $X \subseteq \mathbb{P}^r$ is an m -regular finite set, then

$$\text{sat. deg}(I_X^a) \leq am.$$

This is optimal for the example of the three coordinate points in \mathbb{P}^2 .

Our second main result asserts that the same statement holds when $\dim X = 1$.

Theorem C. *Let $X \subseteq \mathbb{P}^r$ be a smooth m -regular curve. Then*

$$(I_X^a)_t = (I_X^{(a)})_t \quad \text{for } t \geq am.$$

In fact, for the saturation bound it suffices that the curve X be reduced. The statement is optimal (for all a) for instance when $X \subseteq \mathbb{P}^4$ is a rational normal curve. We also show that if $X \subseteq \mathbb{P}^r$ is a reduced surface, then $\text{reg}(\mathcal{I}_X^a) \leq a \cdot \text{reg}(\mathcal{I}_X)$. We do not know any examples where the analogous statements fail for smooth varieties of higher dimension.

¹For saturated ideals, Castelnuovo–Mumford regularity of I_X agrees with an algebraic notion of regularity introduced by Eisenbud and Goto [1984] that we propose to call *arithmetic regularity*. An arbitrary ideal $J \subseteq S$ is arithmetically m -regular if and only if J^{sat} is m -regular and $\text{sat. deg}(J) \leq m$. Given that we are interested in establishing bounds on saturation degree, unless otherwise stated we always refer to regularity in the geometric sense.

Returning to the setting of [Theorem A](#), the first and third authors showed with Bertram some years ago [[Bertram et al. 1991](#)] that if $X \subseteq \mathbb{P}^r$ is a smooth complex projective variety of codimension e cut out as a scheme by homogeneous polynomials of degrees $d_0 \geq \cdots \geq d_p$, then \mathcal{I}_X^a is $(ad_0 + d_1 + \cdots + d_{e-1} - e)$ -regular in the sense of Castelnuovo–Mumford. Note however that this does not address the questions of saturation required to control the arithmetic (Eisenbud–Goto) regularity of I_X^a .² In fact, one can view [Theorem A](#) as promoting the results of [[Bertram et al. 1991](#)] to statements about arithmetic regularity:

Corollary D. *Assume that $J \subseteq S$ satisfies the hypotheses of [Theorem A](#). Then*

$$\text{arith. reg}(J^a) \leq ad_0 + (d_1 + \cdots + d_r - r).$$

It is known [[Kodiyalam 2000](#); [Cutkosky et al. 1999](#)] that if $J \subseteq S$ is an arbitrary homogeneous ideal then

$$\text{arith. reg}(J^a) = ad + b \quad \text{when } a \gg 0,$$

where d is the maximal degree needed to generate a reduction of J — which coincides with the generating degree of J when it is equigenerated — and b is some constant. However, computing the constant term b has proven elusive, and [Corollary D](#) gives a bound in the case at hand.

The proofs of these results revolve around using complexes of sheaves to study the image in $H_*^0(\mathbb{P}^r, \mathcal{I}_X^a) = (I_X^a)^{\text{sat}}$ of the powers of the ideal spanned by generators of I_X or J : this approach was inspired in part by geometrizing the arguments of [[Cooper et al. 2020](#)] for codimension-two subvarieties. Specifically, suppose that

$$\varepsilon : U_0 =_{\text{def}} \bigoplus \mathcal{O}_{\mathbb{P}^r}(-d_i) \rightarrow \mathcal{I}_X$$

is the surjective map of sheaves determined by generators of I_X or J . If X is m -regular, then this sits in an exact complex U_\bullet of bundles

$$0 \rightarrow U_{r-1} \rightarrow U_{r-2} \rightarrow \cdots \rightarrow U_1 \rightarrow U_0 \xrightarrow{\varepsilon} \mathcal{I}_X \rightarrow 0,$$

where $\text{reg}(U_i) \leq m + i$. Weyman [[1979](#)] (see also [[Tchernev 1996](#)]) constructs a new complex $L_\bullet = \text{Sym}^a(U_\bullet)$ that takes the form

$$\cdots \rightarrow L_2 \rightarrow L_1 \rightarrow S^a(U_0) \rightarrow \mathcal{I}_X^a \rightarrow 0,$$

where $\text{reg}(L_i) \leq am + i$. This complex is exact only off X , but as in [[Gruson et al. 1983](#)] when $\dim X = 1$ one can still read off the surjectivity of

$$H^0(\mathbb{P}^r, S^a(U_0)(t)) \rightarrow H^0(\mathbb{P}^r, \mathcal{I}_X^a(t))$$

for $t \geq am$. This gives [Theorem C](#).

Turning to [Theorem A](#), a natural idea is to start with the Koszul complex

$$\cdots \rightarrow \Lambda^3 U_0 \rightarrow \Lambda^2 U_0 \rightarrow U_0 \rightarrow \mathcal{I}_X \rightarrow 0.$$

²In particular, the proof of Proposition 2.2 in [[Arsie and Vatne 2003](#)] seems to be erroneous.

As established by Buchsbaum and Eisenbud [1975], this determines a new complex

$$\cdots \rightarrow S^{a,1^2}(U_0) \rightarrow S^{a,1}(U_0) \rightarrow S^a(U_0) \rightarrow \mathcal{I}_X^a \rightarrow 0, \quad (*)$$

where $S^{a,1^k}(U_0)$ denotes the Schur power of U_0 corresponding to the Young diagram $(a, 1^k)$. We observe that

$$\operatorname{reg}(S^{a,1^i}(U_0)) \leq ad_0 + d_1 + \cdots + d_i,$$

so if $(*)$ were exact then the statement of [Theorem A](#) would follow immediately. Unfortunately $(*)$ is exact only if X is a complete intersection, but by blowing-up X this construction yields an exact complex whose cohomology groups one can control with some effort. At the end of the day, the computation boils down to using Kodaira–Nakano vanishing on X to prove a vanishing statement for symmetric powers of the normal bundle to X in \mathbb{P}^r :

Proposition E. *Let $X \subseteq \mathbb{P}^r$ be a smooth complex projective variety, and denote by $N = N_{X/\mathbb{P}^r}$ the normal bundle to X in \mathbb{P}^r . Then*

$$H^i(X, S^k N \otimes \det N \otimes \mathcal{O}_X(\ell)) = 0 \quad \text{for } i > 0,$$

every $k \geq 0$ and every $\ell \geq -r$.

(Similar but slightly different vanishings were established by Schneider and Zintl [1993].) We hope that some of these ideas may find other applications in the future.³

The paper is organized as follows. [Section 1](#) is devoted to [Theorem C](#). We collect in [Section 2](#) some preliminary results towards the Macaulay-type bounds. Specifically, we discuss the Buchsbaum–Eisenbud powers of Koszul complexes, the computation of some pushforwards from a blowing-up, and [Proposition E](#). The proof of [Theorem A](#) occupies [Section 3](#).

Concerning our assumptions: we work throughout over the complex numbers. As the referee points out, the main results stated above do not require X to be irreducible or even pure-dimensional. However, the essential ideas occur for irreducible varieties, and we generally leave it to the reader to think through this technical improvement.

1. Saturation and regularity

The present section is devoted to the proof of [Theorem C](#) from the [Introduction](#).

We start with some general remarks. Let $X \subseteq \mathbb{P}^r$ be a complex projective scheme, with ideal sheaf $\mathcal{I}_X \subseteq \mathcal{O}_{\mathbb{P}^r}$ and homogeneous ideal $I_X \subseteq S$. Denote by U_\bullet the locally free resolution of \mathcal{I}_X obtained by sheafifying a minimal graded free resolution of I_X :

$$0 \rightarrow U_r \rightarrow U_{r-1} \rightarrow \cdots \rightarrow U_1 \rightarrow U_0 \xrightarrow{\varepsilon} \mathcal{I}_X \rightarrow 0. \quad (1-1)$$

³We remark that some of the auxiliary results appearing here — for example, [Proposition E](#) — were known to the first and third authors some years ago in connection with their work on [\[Bertram et al. 1991\]](#). However, they were put aside in favor of the simpler arguments with vanishing theorems that eventually appeared in that paper.

Thus each U_i is a direct sum of line bundles, and we recover the original resolution as the complex $H_*^0(\mathbb{P}^r, U_\bullet)$ obtained from U_\bullet by taking global sections of all twists.

Consider now the surjective homomorphism of sheaves

$$S^a(\varepsilon) : S^a U_0 \rightarrow \mathcal{I}_X^a.$$

For any $t \geq 0$ one has

$$H^0(\mathbb{P}^r, \mathcal{I}_X^a(t)) = ((I_X^a)^{\text{sat}})_t.$$

On the other hand, the fact that U_0 is constructed from minimal generators of I_X implies that

$$\text{im}(H^0(\mathbb{P}^r, S^a(U_0)(t)) \rightarrow H^0(\mathbb{P}^r, \mathcal{I}_X^a(t))) = (I_X^a)_t.$$

Therefore:

Lemma 1.1. *The degree t pieces of I_X^a and $(I_X^a)^{\text{sat}}$ coincide if and only if the homomorphism*

$$H^0(\mathbb{P}^r, S^a(U_0)(t)) \rightarrow H^0(\mathbb{P}^r, \mathcal{I}_X^a(t))$$

determined by $S^a(\varepsilon)$ is surjective. □

The plan is to study $S^a(\varepsilon)$ by realizing it as the last map of a complex $S^a(U_\bullet)$.

Specifically, consider a smooth variety M , a subvariety $X \subseteq M$, and a locally free resolution U_\bullet of $\mathcal{I}_X \subseteq \mathcal{O}_M$ as above:

$$0 \rightarrow U_r \rightarrow U_{r-1} \rightarrow \cdots \rightarrow U_1 \rightarrow U_0 \xrightarrow{\varepsilon} \mathcal{I}_X \rightarrow 0. \quad (1-2)$$

As explained by Weyman [1979] and Tchernev [1996], U_\bullet determines for fixed $a \geq 1$ a new complex $L_\bullet = S^a(U_\bullet)$ having the shape

$$\cdots \rightarrow L_4 \rightarrow L_3 \rightarrow (S^{a-2}U_0 \otimes \Lambda^2 U_1) \oplus (S^{a-1}U_0 \otimes U_2) \rightarrow S^{a-1}U_0 \otimes U_1 \rightarrow S^a U_0 \rightarrow \mathcal{I}_X^a \rightarrow 0. \quad (1-3)$$

The last map on the right is $S^a(\varepsilon)$, and the homomorphism $S^{a-1}U_0 \otimes U_1 \rightarrow S^a U_0$ is the natural one arising as the composition

$$S^{a-1}U_0 \otimes U_1 \rightarrow S^{a-1}U_0 \otimes U_0 \rightarrow S^a U_0.$$

The L_i are determined by setting

$$C^k(U_j) = \begin{cases} S^k U_j & \text{if } j \text{ is even,} \\ \Lambda^k U_j & \text{if } j \text{ is odd,} \end{cases} \quad (1-4)$$

and then taking

$$L_i = \bigoplus_{\substack{k_0 + \cdots + k_r = a \\ k_1 + 2k_2 + \cdots + rk_r = i}} C^{k_0}(U_0) \otimes C^{k_1}(U_1) \otimes \cdots \otimes C^{k_r}(U_r). \quad (1-5)$$

It follows from [Weyman 1979, Theorem 1] or [Tchernev 1996, Theorem 2.1] that

$$\text{the complex (1-3) is exact away from } X. \quad (1-6)$$

In general one does not expect exactness at points of X , but when X is smooth the right-most terms at least are well-behaved:

Lemma 1.2. *Assume that X is nonsingular. Then the sequence*

$$S^{a-1}U_0 \otimes U_1 \rightarrow S^a U_0 \rightarrow \mathcal{I}_X^a \rightarrow 0$$

is exact.

Proof. The question being local, we can work over the local ring $\mathcal{O} = \mathcal{O}_{M,x}$ of M at a point $x \in X$. Since X is smooth, $\mathcal{I} = \mathcal{I}_{X,x} \subseteq \mathcal{O}$ is generated by a regular sequence of length $e = \text{codim } X$. Thus \mathcal{I} has a minimal presentation

$$\Lambda^2 \mathcal{U} \rightarrow \mathcal{U} \rightarrow \mathcal{I} \rightarrow 0$$

given by the beginning of a Koszul complex, where $\mathcal{U} = \mathcal{O}^e$ is a free module of rank e . Here one checks by hand the exactness of

$$S^{a-1}\mathcal{U} \otimes \Lambda^2 \mathcal{U} \rightarrow S^a \mathcal{U} \rightarrow \mathcal{I}^a \rightarrow 0.$$

(Compare (2-3) below.) An arbitrary free presentation of \mathcal{I} then has the form

$$\Lambda^2 \mathcal{U} \oplus \mathcal{A} \oplus \mathcal{B} \rightarrow \mathcal{U} \oplus \mathcal{A} \rightarrow \mathcal{I} \rightarrow 0,$$

where \mathcal{A} is a free module mapping to zero in \mathcal{I} , \mathcal{B} is a free module mapping to zero in $\mathcal{U} \oplus \mathcal{A}$, and the left-hand map is the identity on \mathcal{A} . It suffices to verify the exactness of

$$S^{a-1}(\mathcal{U} \oplus \mathcal{A}) \otimes (\Lambda^2 \mathcal{U} \oplus \mathcal{A}) \rightarrow S^a(\mathcal{U} \oplus \mathcal{A}) \rightarrow \mathcal{I}^a \rightarrow 0,$$

and this is clear upon writing $S^a(\mathcal{U} \oplus \mathcal{A}) = S^a \mathcal{U} \oplus \mathcal{A} \otimes S^{a-1}(\mathcal{U} \oplus \mathcal{A})$. \square

With these preliminaries out of the way, we now prove (a slight strengthening of) [Theorem C](#) from the [Introduction](#).

Theorem 1.3. *Let $X \subseteq \mathbb{P}^r$ be a reduced (but possibly singular) curve, and assume that X is m -regular in the sense of Castelnuovo–Mumford. Denote by $I_X \subseteq S$ the homogeneous ideal of X . Then*

$$\text{sat. deg}(I_X^a) \leq am.$$

Proof. The m -regularity of X means that we can take a resolution U_\bullet of \mathcal{I}_X as in (1-1) where U_i is a direct sum of line bundles of degrees $\geq -m - i$, i.e., $\text{reg}(U_i) \leq m + i$. Consider the resulting Weyman complex $L_\bullet = S^a(U_\bullet)$:

$$\rightarrow L_3 \rightarrow L_2 \rightarrow L_1 \rightarrow L_0 \rightarrow \mathcal{I}_X^a \rightarrow 0, \quad (*)$$

where the last map is the surjection $S^a(\varepsilon) : L_0 = S^a U_0 \rightarrow \mathcal{I}_X$. In view of [Lemma 1.1](#), the issue is to establish the surjectivity of the homomorphism

$$H^0(\mathbb{P}^r, L_0(t)) \rightarrow H^0(\mathbb{P}^r, \mathcal{I}_X^a(t)) \quad (**)$$

for $t \geq am$. To this end, observe first from (1-4) and (1-5) that

$$\text{reg}(L_i) \leq am + i.$$

Consider next the homology sheaves $\mathcal{H}_i = \mathcal{H}_i(L_\bullet \rightarrow \mathcal{I}_X^a)$ of the augmented complex $(*)$. (So for $i = 0$ we understand $\mathcal{H}_0 = \ker(L_0 \rightarrow \mathcal{I}_X^a)/\text{im}(L_1 \rightarrow L_0)$.) Thanks to (1-6), these are all supported on the one-dimensional set X . Moreover, it follows from Lemma 1.2 that \mathcal{H}_0 is supported on the finitely many singular points of X . Therefore the required surjectivity $(**)$ is a consequence of the first statement of the following lemma. \square

Lemma 1.4. *Consider a complex L_\bullet of coherent sheaves on \mathbb{P}^r sitting in a diagram*

$$\cdots \rightarrow L_3 \rightarrow L_2 \rightarrow L_1 \rightarrow L_0 \xrightarrow{\varepsilon} \mathcal{F} \rightarrow 0, \quad (1-7)$$

and denote by $\mathcal{H}_i = \mathcal{H}_i(L_\bullet \rightarrow \mathcal{F})$ the i -th homology sheaf of the augmented complex (1-7).⁴ Assume that ε is surjective, and let p be an integer with the property that L_i is $(p+i)$ -regular for every i .

(i) *If each \mathcal{H}_i is supported on a set of dimension $\leq i$, then the homomorphism*

$$H^0(\mathbb{P}^r, L_0(t)) \rightarrow H^0(\mathbb{P}^r, \mathcal{F}(t))$$

is surjective for $t \geq p$.

(ii) *If each \mathcal{H}_i is supported on a set of dimension $\leq i + 1$, then \mathcal{F} is p -regular.*

Proof. This is established by chopping L_\bullet into short exact sequences in the usual way and chasing through the resulting diagram. (Compare [Lazarsfeld 2004, B.1.2, B.1.3], but note that the sheaf \mathcal{H}_0 there should refer to the augmented complex, as above.) \square

Remark 1.5. The argument just completed shows that Theorem 1.3 remains true if X has several irreducible components, as well as possibly isolated points.

We conclude this section by observing that the same argument proves that Castelnuovo–Mumford regularity of surfaces behaves submultiplicatively in powers. For curves, this has been known for some time [Chandler 1997; Sidman 2002].

Proposition 1.6. *Let $X \subseteq \mathbb{P}^r$ be a reduced (but possibly singular) surface, and denote by $\mathcal{I}_X \subseteq \mathcal{O}_{\mathbb{P}^r}$ the ideal sheaf of X . If \mathcal{I}_X is m -regular, then \mathcal{I}_X^a is am -regular.*

Sketch of proof. One argues just as in the proof of Theorem 1.3, reducing to statement (ii) of the previous lemma. \square

2. Macaulay-type bounds: preliminaries

This section is devoted to some preliminary results that will be used in the proof of Theorem A from the Introduction. In the first subsection, we discuss symmetric powers of a Koszul complex. The second is devoted to the computation of some direct images from a blowup. Finally Section 2C gives the proof of Proposition E from the Introduction. We will focus mostly on the case when X is a variety.

2A. Powers of Koszul complexes. In this subsection we review the construction of symmetric powers of a Koszul complex. In the local setting this (and much more) appears in [Buchsbaum and Eisenbud 1975],

⁴So as above, the group of zero-cycles used to compute \mathcal{H}_0 is $\ker(\varepsilon)$.

and it was revisited by Srinivasan [1989]. However, for the convenience of the reader, we give here a quick sketch of the particular facts we require. We continue to work over the complex numbers.

Let M be a smooth algebraic variety, and let V be a vector bundle of rank e on M . Fix integers $a, k \geq 1$. We denote by $S^{a,1^{k-1}}(V)$ the Schur power of V corresponding to the partition $(a, 1, \dots, 1)$ ($k-1$ repetitions of 1). It follows from Pieri's rule that

$$S^{a,1^{k-1}}(V) = \ker(\Lambda^{k-1}V \otimes S^aV \rightarrow \Lambda^{k-2}V \otimes S^{a+1}V) = \operatorname{im}(\Lambda^kV \otimes S^{a-1}V \rightarrow \Lambda^{k-1}V \otimes S^aV). \quad (2-1)$$

Remark 2.1 (properties of $S^{a,1^{k-1}}(V)$). We collect some useful observations concerning this Schur power.

(i) If $k = 1$ then $S^{a,1^{k-1}}(V) = S^aV$, while if $a = 1$ then $S^{a,1^{k-1}}(V) = \Lambda^kV$. Moreover,

$$S^{a,1^{k-1}}(V) = 0 \quad \text{when } k > \operatorname{rank} V.$$

(ii) The bundle $S^{a,1^{k-1}}(V)$ is actually a summand of $S^{a-1}V \otimes \Lambda^kV$. In fact, Pieri shows that

$$S^{a-1}V \otimes \Lambda^kV = S^{a,1^{k-1}}(V) \oplus S^{a-1,1^k}(V).$$

(iii) If L is a line bundle on M , then it follows from (2-1) or (ii) that

$$S^{a,1^{k-1}}(V \otimes L) = S^{a,1^{k-1}}(V) \otimes L^{\otimes a+k-1}.$$

(iv) Suppose that $M = \mathbb{P}^r$ and

$$V = \mathcal{O}_{\mathbb{P}^r}(-d_0) \oplus \cdots \oplus \mathcal{O}_{\mathbb{P}^r}(-d_p)$$

with $d_0 \geq \cdots \geq d_p$. Then it follows from (ii) that $S^{a,1^{k-1}}(V)$ is a direct sum of line bundles of degrees $\geq -(ad_0 + d_1 + \cdots + d_{k-1})$ and, moreover, a summand of this degree appears. In other words,

$$\operatorname{reg}(S^{a,1^{k-1}}(V)) = ad_0 + d_1 + \cdots + d_{k-1}.$$

One can also realize $S^{a,1^{k-1}}(V)$ geometrically, à la Kempf [1970].

Lemma 2.2. *Let $\pi : \mathbb{P}(V) \rightarrow M$ be the projective bundle of one-dimensional quotients of V , and denote by F the kernel of the canonical quotient $\pi^*V \rightarrow \mathcal{O}_{\mathbb{P}(V)}(1)$, so that F sits in the short exact sequence*

$$0 \rightarrow F \rightarrow \pi^*V \rightarrow \mathcal{O}_{\mathbb{P}(V)}(1) \rightarrow 0 \quad (*)$$

of bundles on $\mathbb{P}(V)$. Then

$$S^{a,1^{k-1}}(V) = \pi_*(\Lambda^{k-1}F \otimes \mathcal{O}_{\mathbb{P}(V)}(a)).$$

Proof. In fact, $(*)$ gives rise to a long exact sequence

$$0 \rightarrow \Lambda^{k-1}F \otimes \mathcal{O}_{\mathbb{P}(V)}(a) \rightarrow \Lambda^{k-1}(\pi^*V) \otimes \mathcal{O}_{\mathbb{P}(V)}(a) \rightarrow \Lambda^{k-2}(\pi^*V) \otimes \mathcal{O}_{\mathbb{P}(V)}(a+1) \rightarrow \cdots.$$

The assertion follows from (2-1) upon taking direct images. □

Now suppose given a map of bundles

$$\varepsilon : V \rightarrow \mathcal{O}_M \quad (2-2)$$

whose image is the ideal sheaf $\mathcal{I} \subseteq \mathcal{O}_M$ of a subscheme $Z \subseteq X$: equivalently, ε is dual to a section $\mathcal{O}_M \rightarrow V^*$ whose zero-scheme is Z . We allow the possibility that ε is surjective, in which case $\mathcal{I} = \mathcal{O}_M$ and $Z = \emptyset$.

If Z has the expected codimension $e = \text{rank}(V)$, then \mathcal{I} is resolved by the Koszul complex associated to ε . The following result of Buchsbaum and Eisenbud gives the resolution of powers of \mathcal{I} .

Proposition 2.3 [Buchsbaum and Eisenbud 1975, Theorem 3.1; Srinivasan 1989, Theorem 2.1]. *Fix $a \geq 1$. Then ε determines a complex*

$$\cdots \rightarrow S^{a,1^2}(V) \rightarrow S^{a,1}(V) \rightarrow S^a V \xrightarrow{S^a(\varepsilon)} \mathcal{I}^a \rightarrow 0 \quad (2-3)$$

of vector bundles on M . This complex is exact provided that either ε is surjective or Z has codimension equal to $\text{rank}(V)$.

Observe from Remark 2.1(i) that this complex has the same length as the Koszul complex of ε .

Proof. Returning to the setting of Lemma 2.2, denote by $\tilde{\varepsilon} : F \rightarrow \mathcal{O}_{\mathbb{P}(V)}$ the composition of the inclusion $F \hookrightarrow \pi^* V$ with $\pi^* \varepsilon : \pi^* V \rightarrow \pi^* \mathcal{O}_M$. The zero-locus of $\tilde{\varepsilon}$ defines the natural embedding of $\mathbb{P}(\mathcal{I})$ in $\mathbb{P}(V)$. Now consider the Koszul complex of $\tilde{\varepsilon}$. After twisting by $\mathcal{O}_{\mathbb{P}(V)}(a)$ this has the form

$$\cdots \rightarrow \Lambda^2 F \otimes \mathcal{O}_{\mathbb{P}(V)}(a) \rightarrow F \otimes \mathcal{O}_{\mathbb{P}(V)}(a) \rightarrow \mathcal{O}_{\mathbb{P}(V)}(a) \rightarrow \mathcal{O}_{\mathbb{P}(\mathcal{I})}(a) \rightarrow 0. \quad (*)$$

In view of Lemma 2.2, (2-3) arises by taking direct images. If ε is surjective, or defines a regular section of V^* , then the Koszul complex $(*)$ is exact. Since the higher direct images of all the terms vanish, $(*)$ pushes down to an exact complex. Furthermore, in this case $\pi_* \mathcal{O}_{\mathbb{P}(\mathcal{I})}(a) = \mathcal{I}^a$ (see [Fulton and Lang 1985, Theorem IV.2.2]), and the exactness of (2-3) follows. \square

Example 2.4 (Macaulay's theorem). Suppose as in the Introduction that $f_0, \dots, f_p \in \mathbb{C}[x_0, \dots, x_r]$ are homogeneous polynomials of degrees $d_0 \geq \dots \geq d_p$ that generate a finite colength ideal J . This gives rise to a surjective map

$$V = \bigoplus \mathcal{O}_{\mathbb{P}^r}(-d_i) \rightarrow \mathcal{O}_{\mathbb{P}^r} \rightarrow 0$$

of bundles on projective space. Keeping in mind Remark 2.1(iv), Macaulay's statements (1) and (2) follow by looking at the cohomology of the resulting complex (2-3). When $p = r$ this complex has length $r + 1$, so one can also read off the nonsurjectivity of

$$H^0(\mathbb{P}^r, S^a V(t)) \rightarrow H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(t))$$

when $t < ad_0 + d_1 + \dots + d_r - r$.

Example 2.5 (complete intersection ideals). Suppose that $Z \subseteq \mathbb{P}^r$ is a complete intersection of dimension ≥ 0 . Applying (2-3) to the Koszul resolution of its homogeneous ideal I_Z , one sees that I_Z^a is saturated for every $a \geq 1$. This is a result of Zariski.

2B. Pushforwards from a blowing-up. We compute here the direct images of multiples of the exceptional divisor under the blowing-up of a smooth subvariety.

Consider then a smooth variety M and a nonsingular subvariety $X \subseteq M$ having codimension $e \geq 2$ and ideal sheaf $\mathcal{I} = \mathcal{I}_X \subseteq \mathcal{O}_M$. We consider the blowing-up

$$\mu : M' = \text{Bl}_X(M) \rightarrow M$$

of M along X . Write $E \subseteq M'$ for the exceptional divisor of M' , so that $\mathcal{I} \cdot \mathcal{O}_{M'} = \mathcal{O}_{M'}(-E)$. We recall that if $a > 0$ then

$$\mu_* \mathcal{O}_{M'}(-aE) = \mathcal{I}^a \quad \text{and} \quad R^j \mu_* \mathcal{O}_{M'}(-aE) = 0 \quad \text{for } j > 0. \quad (2-4)$$

The following proposition gives the analogous computation for positive multiples of E .

Proposition 2.6. *Fix $a > 0$. Then*

$$R^j \mu_* \mathcal{O}_{M'}(aE) = \text{Ext}_{\mathcal{O}_M}^j(\mathcal{I}^{a-e+1}, \mathcal{O}_M).^5 \quad (2-5)$$

In particular, $\mu_ \mathcal{O}_{M'}(aE) = \mathcal{O}_M$, $R^j \mu_* \mathcal{O}_{M'}(aE) = 0$ if $j \neq 0, e-1$, and*

$$R^{e-1} \mu_* \mathcal{O}_{M'}(aE) = \text{Ext}_{\mathcal{O}_M}^{e-1}(\mathcal{I}^{a-e+1}, \mathcal{O}_M).$$

Proof. This is a consequence of duality for μ , which asserts that

$$R\mu_* R\mathcal{H}om_{\mathcal{O}_{M'}}(\mathcal{F}, \omega_\mu) = R\mathcal{H}om_{\mathcal{O}_M}(R\mu_* \mathcal{F}, \mathcal{O}_M) \quad (*)$$

for any sheaf \mathcal{F} on M' , where ω_μ denotes the relative dualizing sheaf for μ [Huybrechts 2006, (3.19) on page 86]. We apply this with

$$\mathcal{F} = \mathcal{O}_{M'}((e-1-a)E).$$

Then $R\mu_* \mathcal{F} = \mathcal{I}^{a-e+1}$ thanks to (2-4) (and a direct computation when $0 < a < e-1$), and $\omega_\mu = \mathcal{O}_{M'}((e-1)E)$. Therefore the first assertion of the proposition follows from (*). The vanishing of $\text{Ext}_{\mathcal{O}_M}^j(\mathcal{I}^{a-e+1}, \mathcal{O}_M)$ for $j \neq 0, e-1$ follows from the perfection of powers of the ideal of a smooth variety (which in turn is a consequence, e.g., of Proposition 2.3). \square

Remark 2.7 (generalization to multiplier ideal sheaves). Let $\mathfrak{b} \subseteq \mathcal{O}_M$ be an arbitrary ideal sheaf, and let $\mu : M' \rightarrow M$ be a log resolution of \mathfrak{b} , with $\mathfrak{b} \cdot \mathcal{O}_{M'} = \mathcal{O}_{M'}(-E)$. A completely parallel argument shows that for $a > 0$,

$$R^j \mu_* \mathcal{O}_{M'}(aE) = \text{Ext}_{\mathcal{O}_M}^j(\mathcal{J}(\mathfrak{b}^a), \mathcal{O}_M),$$

where $\mathcal{J}(\mathfrak{b}^a)$ is the multiplier ideal of \mathfrak{b}^a . The formula (2-5) is a special case of this.

Corollary 2.8. *Continuing to work in characteristic zero, fix $a \geq 1$ and denote by $N = N_{X/M}$ the normal bundle to X in M . If $a \leq e-1$, then*

$$R^{e-1} \mu_* \mathcal{O}_{M'}(aE) = 0.$$

⁵When $0 < a < e-1$ we take $\mathcal{I}^{a-e+1} = \mathcal{O}_M$.

If $a \geq e$, then $R^{e-1}\mu_*\mathcal{O}_{M'}(aE)$ has a filtration with successive quotients

$$S^k N \otimes \det N \quad \text{for } 0 \leq k \leq a - e.$$

Proof. The first statement follows directly from the previous proposition. For the second, recall first that if E is any locally free \mathcal{O}_X -module, then — X being nonsingular of codimension e in M —

$$\mathcal{E}xt_{\mathcal{O}_M}^e(E, \mathcal{O}_M) = E^* \otimes \det N,$$

while all the other $\mathcal{E}xt^j$ vanish. The claim then follows from [Proposition 2.6](#), using the exact sequences

$$0 \rightarrow \mathcal{I}^{k+1} \rightarrow \mathcal{I}^k \rightarrow S^k N^* \rightarrow 0$$

together with the isomorphism $(S^k(N^*))^* = S^k N$ valid in characteristic zero. \square

Remark 2.9. Recalling that $E = \mathbb{P}(N^*)$, one can inductively prove [Corollary 2.8](#) directly, circumventing [Proposition 2.6](#), by pushing forward the exact sequences

$$0 \rightarrow \mathcal{O}_{M'}((k-1)E) \rightarrow \mathcal{O}_{M'}(kE) \rightarrow \mathcal{O}_E(kE) \rightarrow 0.$$

However, it seemed to us that [Proposition 2.6](#) may be of independent interest.

2C. A vanishing theorem for normal bundles. This final subsection is devoted to the proof of:

Proposition 2.10. *Let $X \subseteq \mathbb{P}^r$ be a smooth complex projective variety of dimension n , and denote by $N = N_{X/\mathbb{P}^r}$ the normal bundle to X . Then*

$$H^i(X, S^k N \otimes \det N \otimes \mathcal{O}_X(\ell)) = 0$$

for all $i > 0$, $k \geq 0$ and $\ell \geq -r$.

Here $\mathcal{O}_X(k)$ denotes $\mathcal{O}_{\mathbb{P}^r}(k)|_X$. We remark that similar statements were established by Schneider and Zintl [1993], but this particular vanishing does not seem to appear there. Other vanishings for normal bundles played a central role in [Ein and Lazarsfeld 1993].

Proof of Proposition 2.10. We use the abbreviation $\mathbb{P} = \mathbb{P}^r$. Starting from the exact sequence $0 \rightarrow TX \rightarrow T\mathbb{P}|_X \rightarrow N \rightarrow 0$, we get a long exact sequence

$$\dots \rightarrow S^{k-2}T\mathbb{P}|_X \otimes \Lambda^2 TX \rightarrow S^{k-1}T\mathbb{P}|_X \otimes TX \rightarrow S^k T\mathbb{P}|_X \rightarrow S^k N \rightarrow 0. \quad (*)$$

By adjunction, $\det N \otimes \mathcal{O}_X(\ell) = \omega_X \otimes \mathcal{O}_X(\ell + r + 1)$. So after twisting through by $\det N \otimes \mathcal{O}_X(\ell)$ in $(*)$, we see that the proposition will follow if we prove

$$H^i(X, S^{k-j}T\mathbb{P}|_X \otimes \Lambda^j TX \otimes \omega_X \otimes \mathcal{O}_X(\ell + r + 1)) = 0 \quad \text{for } i \geq j + 1 \quad (**)$$

when $0 \leq j \leq k$ and $\ell \geq -r$. It follows from the Euler sequence that $S^m T\mathbb{P}|_X$ has a presentation of the form

$$0 \rightarrow \bigoplus \mathcal{O}_X(m-1) \rightarrow \bigoplus \mathcal{O}_X(m) \rightarrow S^m T\mathbb{P}|_X \rightarrow 0,$$

so for (**) it suffices in turn to verify that

$$H^i(X, \Lambda^j TX \otimes \omega_X \otimes \mathcal{O}_X(\ell_1)) = 0$$

for $i \geq j + 1$ and $\ell_1 > 0$. But $\Lambda^j TX \otimes \omega_X = \Omega_X^{n-j}$, so finally we're asking that

$$H^i(X, \Omega_X^{n-j} \otimes \mathcal{O}_X(\ell_1)) = 0 \quad \text{for } i \geq j + 1 \text{ and } \ell_1 > 0,$$

and this follows from Nakano vanishing. □

3. Proof of Theorem A

We now turn to the proof of Theorem A from the Introduction.

Consider then a nonsingular scheme $X \subseteq \mathbb{P}^r$ that is cut out as a scheme by hypersurfaces of degrees $d_0 \geq \dots \geq d_p$. Equivalently, we are given a surjective homomorphism of sheaves,

$$\varepsilon : U \rightarrow \mathcal{I}_X, \quad U = \bigoplus \mathcal{O}_{\mathbb{P}^r}(-d_i).$$

Let $\mu : \mathbb{P}' = \text{Bl}_X(\mathbb{P}^r) \rightarrow \mathbb{P}^r$ be the blowing-up of X , with exceptional divisor $\mathbf{E} \subseteq \mathbb{P}'$, so that $\mathcal{I}_X \cdot \mathcal{O}_{\mathbb{P}'} = \mathcal{O}_{\mathbb{P}'}(-\mathbf{E})$. Write H for the pullback to \mathbb{P}' of the hyperplane class on \mathbb{P}^r , and set $U' = \mu^*U$. Thus on \mathbb{P}' we have a surjective map of bundles,

$$\varepsilon' : U' \rightarrow \mathcal{O}_{\mathbb{P}'}(-\mathbf{E}). \tag{3-1}$$

Noting that

$$H^0(\mathbb{P}', \mathcal{O}_{\mathbb{P}'}(tH - a\mathbf{E})) = H^0(\mathbb{P}^r, \mathcal{I}_X^a \otimes \mathcal{O}_{\mathbb{P}^r}(t)),$$

one sees as in Lemma 1.1 that the question is to prove the surjectivity of

$$H^0(\mathbb{P}', S^a U' \otimes \mathcal{O}_{\mathbb{P}'}(tH)) \rightarrow H^0(\mathbb{P}', \mathcal{O}_{\mathbb{P}'}(tH - a\mathbf{E})) \tag{3-2}$$

for $t \geq ad_0 + d_1 + \dots + d_r - r$.

To this end, we pass to the Buchsbaum–Eisenbud complex (2-3) constructed from

$$U' \otimes \mathcal{O}_{\mathbb{P}'}(\mathbf{E}) \xrightarrow{\varepsilon'} \mathcal{O}_{\mathbb{P}'} \rightarrow 0.$$

Twisting through by $\mathcal{O}_{\mathbb{P}'}(tH - a\mathbf{E})$, we arrive at a long exact sequence of vector bundles on \mathbb{P}' having the following form:

$$\begin{array}{ccccccc} \dots \rightarrow & S^{a,1^2} U' \otimes \mathcal{O}_{\mathbb{P}'}(tH+2\mathbf{E}) & \rightarrow & S^{a,1} U' \otimes \mathcal{O}_{\mathbb{P}'}(tH+\mathbf{E}) & \rightarrow & S^a U' \otimes \mathcal{O}_{\mathbb{P}'}(tH) & \rightarrow \mathcal{O}_{\mathbb{P}'}(tH-a\mathbf{E}) \rightarrow 0 \\ & \parallel & & \parallel & & \parallel & \\ & C_2 & & C_1 & & C_0 & \end{array} \tag{3-3}$$

With indexing as indicated, the i -th term of this sequence is given by

$$C_i = S^{a,1^i}(U') \otimes \mathcal{O}_{\mathbb{P}'}(tH + i\mathbf{E}).$$

In order to establish the surjectivity (3-2) it suffices upon chasing through (3-3) to prove that

$$H^i(\mathbb{P}', C_i) = 0 \quad \text{for } 1 \leq i \leq r \quad (3-4)$$

provided that $t \geq ad_0 + d_1 + \cdots + d_r - r$. But now recall (Remark 2.1) that if $i \leq r$ then $S^{a,1^i}(U')$ is a sum of line bundles $\mathcal{O}_{\mathbb{P}'}(mH)$ with

$$m \geq -ad_0 - d_1 - \cdots - d_i \geq -ad_0 - d_1 - \cdots - d_r.$$

Hence, when $t \geq ad_0 + d_1 + \cdots + d_r - r$, C_i is a sum terms of the form

$$\mathcal{O}_{\mathbb{P}'}(\ell H + iE) \quad \text{with } \ell \geq -r.$$

Therefore (3-4)—and with it Theorem A—is a consequence of:

Proposition 3.1. *If $\ell \geq -r$, then*

$$H^i(\mathbb{P}', \mathcal{O}_{\mathbb{P}'}(\ell H + iE)) = 0 \quad \text{for } i > 0.$$

Proof. Thanks to the Leray spectral sequence, it suffices to show

$$H^j(\mathbb{P}^r, R^k \mu_* \mathcal{O}_{\mathbb{P}'}(\ell H + iE)) = 0 \quad \text{when } j + k = i > 0. \quad (*)$$

For $k = 0$, observe that $\mu_* \mathcal{O}_{\mathbb{P}'}(\ell H + iE) = \mathcal{O}_{\mathbb{P}^r}(\ell)$, and these sheaves have no higher cohomology when $\ell \geq -r$. Suppose that $k > 0$. Suppose first that X is smooth and irreducible of codimension e . By Proposition 2.6, the only nonvanishing higher direct images are the $R^{e-1} \mu_* \mathcal{O}_{\mathbb{P}'}(\ell H + iE)$, which do not appear when $i \leq e - 1$. So (*) holds when $j = 0$ and $k = e - 1$. It remains to consider the case $k = e - 1$ and $i \geq e$, so that $j = i - (e - 1) > 0$. Here Corollary 2.8 implies that the R^{e-1} have a filtration with quotients

$$S^\alpha N \otimes \det N \otimes \mathcal{O}_X(\ell),$$

where as above $N = N_{X/\mathbb{P}^r}$ is the normal bundle to X in \mathbb{P}^r . But since we are assuming $\ell \geq -r$, Proposition 2.10 guarantees that these sheaves have vanishing higher cohomology. This completes the proof when X is irreducible. When X has several components of possibly different dimensions one argues similarly, one component at a time: we leave the details to the interested reader. \square

Remark 3.2. Observe that if X is defined by $p < r$ equations, then the argument just completed goes through taking $d_{p+1} = \cdots = d_r = 0$.

Acknowledgements

We are grateful to Sankhaneel Bisui, David Eisenbud, Eloísa Grifo and Claudia Miller for valuable remarks and correspondence.

References

- [Arsie and Vatne 2003] A. Arsie and J. E. Vatne, “A note on symbolic and ordinary powers of homogeneous ideals”, *Ann. Univ. Ferrara Sez. VII (N.S.)* **49** (2003), 19–30. [MR](#) [Zbl](#)
- [Bertram et al. 1991] A. Bertram, L. Ein, and R. Lazarsfeld, “Vanishing theorems, a theorem of Severi, and the equations defining projective varieties”, *J. Amer. Math. Soc.* **4**:3 (1991), 587–602. [MR](#) [Zbl](#)
- [Bocci and Harbourne 2010] C. Bocci and B. Harbourne, “Comparing powers and symbolic powers of ideals”, *J. Algebraic Geom.* **19**:3 (2010), 399–417. [MR](#) [Zbl](#)
- [Buchsbaum and Eisenbud 1975] D. A. Buchsbaum and D. Eisenbud, “Generic free resolutions and a family of generically perfect ideals”, *Adv. Math.* **18**:3 (1975), 245–301. [MR](#) [Zbl](#)
- [Carlson et al. 2003] J. Carlson, S. Müller-Stach, and C. Peters, *Period mappings and period domains*, Cambridge Stud. Adv. Math. **85**, Cambridge Univ. Press, 2003. [MR](#) [Zbl](#)
- [Caviglia 2004] G. Caviglia, *Koszul algebras, Castelnuovo–Mumford regularity, and generic initial ideals*, Ph.D. thesis, University of Kansas, 2004, available at <https://www.proquest.com/docview/305172202>.
- [Chandler 1997] K. A. Chandler, “Regularity of the powers of an ideal”, *Comm. Algebra* **25**:12 (1997), 3773–3776. [MR](#) [Zbl](#)
- [Cooper et al. 2020] S. Cooper, G. Fatabbi, E. Guardo, A. Lorenzini, J. Migliore, U. Nagel, A. Seceleanu, J. Szpond, and A. Van Tuyl, “Symbolic powers of codimension two Cohen–Macaulay ideals”, *Comm. Algebra* **48**:11 (2020), 4663–4680. [MR](#) [Zbl](#)
- [Cutkosky et al. 1999] S. D. Cutkosky, J. Herzog, and N. V. Trung, “Asymptotic behaviour of the Castelnuovo–Mumford regularity”, *Compos. Math.* **118**:3 (1999), 243–261. [MR](#) [Zbl](#)
- [Dao et al. 2018] H. Dao, A. De Stefani, E. Grifo, C. Huneke, and L. Núñez-Betancourt, “Symbolic powers of ideals”, pp. 387–432 in *Singularities and foliations: geometry, topology and applications* (Salvador, Brazil, 2015), edited by R. N. Araújo dos Santos et al., Springer Proc. Math. Stat. **222**, Springer, 2018. [MR](#) [Zbl](#)
- [Ein and Lazarsfeld 1993] L. Ein and R. Lazarsfeld, “Syzygies and Koszul cohomology of smooth projective varieties of arbitrary dimension”, *Invent. Math.* **111**:1 (1993), 51–67. [MR](#) [Zbl](#)
- [Ein et al. 2001] L. Ein, R. Lazarsfeld, and K. E. Smith, “Uniform bounds and symbolic powers on smooth varieties”, *Invent. Math.* **144**:2 (2001), 241–252. [MR](#) [Zbl](#)
- [Eisenbud and Goto 1984] D. Eisenbud and S. Goto, “Linear free resolutions and minimal multiplicity”, *J. Algebra* **88**:1 (1984), 89–133. [MR](#) [Zbl](#)
- [Fulton and Lang 1985] W. Fulton and S. Lang, *Riemann–Roch algebra*, Grundlehren der Math. Wissenschaften **277**, Springer, 1985. [MR](#) [Zbl](#)
- [Geramita et al. 1995] A. V. Geramita, A. Gimigliano, and Y. Pitteloud, “Graded Betti numbers of some embedded rational n -folds”, *Math. Ann.* **301**:2 (1995), 363–380. [MR](#) [Zbl](#)
- [Gruson et al. 1983] L. Gruson, R. Lazarsfeld, and C. Peskine, “On a theorem of Castelnuovo, and the equations defining space curves”, *Invent. Math.* **72**:3 (1983), 491–506. [MR](#) [Zbl](#)
- [Hochster and Huneke 2002] M. Hochster and C. Huneke, “Comparison of symbolic and ordinary powers of ideals”, *Invent. Math.* **147**:2 (2002), 349–369. [MR](#) [Zbl](#)
- [Huybrechts 2006] D. Huybrechts, *Fourier–Mukai transforms in algebraic geometry*, Oxford Univ. Press, 2006. [MR](#) [Zbl](#)
- [Kempf 1970] G. R. Kempf, *The singularities of certain varieties in the Jacobian of a curve*, Ph.D. thesis, Columbia University, 1970.
- [Kodiyalam 2000] V. Kodiyalam, “Asymptotic behaviour of Castelnuovo–Mumford regularity”, *Proc. Amer. Math. Soc.* **128**:2 (2000), 407–411. [MR](#) [Zbl](#)
- [Lazarsfeld 2004] R. Lazarsfeld, *Positivity in algebraic geometry, I: Classical setting: line bundles and linear series*, Ergebnisse der Mathematik (3) **48**, Springer, 2004. [MR](#) [Zbl](#)
- [Schneider and Zintl 1993] M. Schneider and J. Zintl, “The theorem of Barth–Lefschetz as a consequence of Le Potier’s vanishing theorem”, *Manuscripta Math.* **80**:3 (1993), 259–263. [MR](#) [Zbl](#)

- [Sidman 2002] J. Sidman, “On the Castelnuovo–Mumford regularity of products of ideal sheaves”, *Adv. Geom.* **2**:3 (2002), 219–229. [MR](#) [Zbl](#)
- [Srinivasan 1989] H. Srinivasan, “Algebra structures on some canonical resolutions”, *J. Algebra* **122**:1 (1989), 150–187. [MR](#) [Zbl](#)
- [Tchernev 1996] A. B. Tchernev, “Acyclicity of symmetric and exterior powers of complexes”, *J. Algebra* **184**:3 (1996), 1113–1135. [MR](#) [Zbl](#)
- [Weyman 1979] J. Weyman, “Resolutions of the exterior and symmetric powers of a module”, *J. Algebra* **58**:2 (1979), 333–341. [MR](#) [Zbl](#)

Communicated by Shunsuke Takagi

Received 2021-04-02 Revised 2021-07-13 Accepted 2021-10-05

ein@uic.edu

*Department of Mathematics, University of Illinois, Chicago, Chicago, IL,
United States*

tha@tulane.edu

Department of Mathematics, Tulane University, New Orleans, LA, United States

robert.lazarsfeld@stonybrook.edu

*Department of Mathematics, Stony Brook University, Stony Brook, NY,
United States*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Antoine Chambert-Loir
Université Paris-Diderot
France

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Raman Parimala	Emory University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Irena Peeva	Cornell University, USA
Brian D. Conrad	Stanford University, USA	Jonathan Pila	University of Oxford, UK
Samit Dasgupta	Duke University, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Bjorn Poonen	Massachusetts Institute of Technology, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Victor Reiner	University of Minnesota, USA
Sergey Fomin	University of Michigan, USA	Peter Sarnak	Princeton University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Wee Teck Gan	National University of Singapore	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	Shunsuke Takagi	University of Tokyo, Japan
Ben J. Green	University of Oxford, UK	Pham Huu Tiep	Rutgers University, USA
Christopher Hacon	University of Utah, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Akshay Venkatesh	Institute for Advanced Study, USA
János Kollár	Princeton University, USA	Melanie Matchett Wood	Harvard University, USA
Michael J. Larsen	Indiana University Bloomington, USA	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2022 is US \$450/year for the electronic version, and \$655/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

ANT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2022 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 16 No. 6 2022

Square-free OM computation of global integral bases	1327
JORDI GUÀRDIA and ENRIC NART	
Explicit computation of symmetric differentials and its application to quasihyperbolicity	1377
NILS BRUIN, JORDAN THOMAS and ANTHONY VÁRILLY-ALVARADO	
Zero-sum subsets in vector spaces over finite fields	1407
COSMIN POHOATA and DMITRIY ZAKHAROV	
Automorphisms of Cartan modular curves of prime and composite level	1423
VALERIO DOSE, GUIDO LIDO and PIETRO MERCURI	
Topological spectrum and perfectoid Tate rings	1463
DIMITRI DINE	
On the geometry and representation theory of isomeric matrices	1501
ROHIT NAGPAL, STEVEN V. SAM and ANDREW SNOWDEN	
Saturation bounds for smooth varieties	1531
LAWRENCE EIN, HUY TÀI HÀ and ROBERT LAZARSFELD	