

Math 125

Solutions to Second Midterm(pineapple)

1. Compute each of the derivatives below as indicated.

4 points

(a) $f(x) = 3x^8 - 5x^4 + 4x - e^3$.

Solution: $f'(x) = 24x^7 - 20x + 4$.

Don't forget that e^3 is a constant just a bit larger than 20, so its derivative is zero.

4 points

(b) $f(x) = e^{4x} \tan x$

Solution: We need both the product rule and the chain rule here:

$$f'(x) = 4e^{4x} \tan x + e^{4x} \sec^2 x.$$

4 points

(c) $f(x) = \frac{8x^3 - 5x}{\sec(\pi x) + x^2}$

Solution: By the quotient rule,

$$f'(x) = \frac{(24x^2 - 5)(\sec(\pi x) + x^2) - (8x^3 - 5x)(\pi \sec(\pi x) \tan(\pi x) + 2x)}{(\sec(\pi x) + x^2)^2}$$

This mess doesn't really simplify much, so why bother trying?

4 points

(d) $f(x) = \arcsin(e^x)$

Solution: Using the chain rule, $f'(x) = \frac{e^x}{\sqrt{1 - e^{2x}}}$.

2. Calculate the indicated derivatives.

4 points

(a) $\frac{d}{d\theta} \sin(4\theta) \cos(2\theta)$

Solution: By the product chain rules, we get $4 \cos(4\theta) \cos(2\theta) - 2 \sin(4\theta) \sin(2\theta)$.

4 points

(b) Calculate the second derivative of $x^2 e^{2x}$ with respect to x .

Solution: If $f(x) = x^2 e^{2x}$, then using the product rule we get $f'(x) = 2x e^{2x} + 2x^2 e^{2x}$ or, if we want to factor it, $2(x + x^2) e^{2x}$. Thus,

$$f''(x) = 2(1 + 2x)e^{2x} + 4(x + x^2)e^{2x} = (2 + 8x + 4x^2)e^{2x}.$$

You could also leave it as $2e^{2x} + 4xe^{2x} + 8xe^{2x} + 4x^2e^{2x}$.

4 points

(c) $\frac{d^{10}}{dt^{10}} 11t^9.$

Solution: If you didn't immediately realize that the tenth derivative of a ninth degree polynomial is zero, maybe you'd see the pattern once you started taking derivatives:

$$f(t) = 11t^9, f'(t) = 9 \cdot 11t^8, f''(t) = 8 \cdot 9 \cdot 11t^7, f'''(t) = 7 \cdot 8 \cdot 9 \cdot 11t^6, \dots, f^{(9)}(t) = 1 \cdot 2 \cdot 3 \cdots 8 \cdot 9 \cdot 11.$$

Since the 9th derivative is a constant, $\frac{d^{10}}{dt^{10}} 11t^9 = 0.$

3. Let $f(x) = x \ln(x^6).$

5 points

(a) Calculate $f'(x).$

Solution: Just apply the product rule and the chain rule to get

$$f'(x) = \ln(x^6) + \frac{x}{x^6} \cdot 6x^5 = \ln(x^6) + 6$$

5 points


(b) For what values of x is $f(x)$ decreasing? If there are none, write "NONE"; otherwise, describe *all* such x .

Solution: The function will be decreasing when $f'(x) < 0$. From the first part, we have $f'(x) = \ln(x^6) + 6$, but remember that $\ln(x^6) = 6 \ln|x|$, so $f'(x) = 6(\ln|x| + 1)$. (Since the power of x inside the log is even, both $x < 0$ and $x > 0$ are in the domain, so we need to include the absolute value.)

Now, we want to know when $4(\ln|x| + 1) < 0$. This happens exactly when $\ln|x| < -1$. Exponentiating both sides gives

$$|x| < e^{-1} \quad \text{or} \quad -\frac{1}{e} < x < \frac{1}{e}.$$

At right is a graph of $f(x)$, and you can see that this corresponds to what we found.



10 points

4. Find the slope of the line tangent to the curve $\sin(xy) = x^2 + y^2 - \pi$ at the point $(0, \sqrt{\pi})$.

Solution: To do this, we use implicit differentiation. Differentiating both sides with respect to x (remembering that y is some unknown function of x and using y' to represent the derivative of y with respect to x), we get


$$\cos(xy)(y + xy') = 2x + 2yy'$$

Now, we can substitute $x = 0$ and $y = \sqrt{\pi}$, then solve for y' .

$$\cos(0)(\sqrt{\pi}) = 0 + 2\sqrt{\pi}y' \quad \text{so} \quad 1/2 = y'.$$

Thus, the slope at the desired point is $1/2$.

The set of points satisfying $\sin(xy) = x^2 + y^2 - \pi$ is shown at right, together with the desired tangent line.



10 points

5. Jimi Chiu makes “designer” shoes¹ that he sells at \$250 a pair. He knows that the number of pairs of shoes he sells is a function of the price he charges; let’s denote this by $N(p)$, where p is the price per pair. Market research tells him that $N'(250)$ is about -20 ; that is, if he raises the price by one dollar, he should expect to sell 20 fewer pairs. The amount of revenue $R(p)$ he makes at a given price will be given by $R(p) = p \cdot N(p)$.

If he typically sells 2000 pairs of shoes at \$250 each, what is $R'(250)$? Should he raise the price a little?

Solution: $N(p)$ is some unknown function, but we know that $N(250) = 2000$ (since he sells 2000 pairs at a price of \$250 each, and we know that $N'(250) = -20$).

We want to calculate $R'(250)$ where $R(p) = p \cdot N(p)$. By the product rule, we have $R'(p) = N(p) + pN'(p)$, so

$$R'(250) = 2000 + 250 \cdot (-20) = 2000 - 5000 = -3000.$$

Since R' is negative, he will lose revenue if he raises the price. So he shouldn’t charge more—in fact, he should lower the price to make more money.

¹No relation to Jimmy Choo shoes, unless you don’t look very closely. Mr. Chiu is also fond of Rollex watches.

10 points

6. Let $f(x) = 4x^3 - x - 1$. Find the equation of a line which passes through the origin and is also tangent to the curve $y = f(x)$ at some point (a, b)

Solution:

While you don't need the graph to do this problem, at right is a graph of what we are trying to solve, to help you keep in mind what is going on. We are looking for the blue line.

There are a couple of ways to do this problem; here's one. Note that since the line we are looking for goes through the origin, it is of the form $y = mx$. Since this line is also tangent to the graph of $f(x)$ at some point (a, b) , we have

$$m = f'(a) = 12a^2 - 1$$

and $b = f(a) = 4a^3 - a - 1$.

So we have to find a so that


$$4a^3 - a - 1 = (12a^2 - 1)(a).$$

$$4a^3 - a - 1 = 12a^3 - a$$

$$-1 = 8a^3.$$

Thus, $a = -1/2$, and so $m = 12/4 - 1 = 2$.

The line we want is $y = 2x$.



10 points

7. Suppose $y = (1 + \cos(x))^{(1+\sin(x))}$. What is $\frac{dy}{dx}$ when $x = \pi/2$ and $y = 1$?

Solution: To do this, we use logarithmic differentiation. Taking the log of both sides gives us

$$\ln(y) = \ln((1 + \cos(x))^{(1+\sin(x))}) = (1 + \sin(x)) \ln(1 + \cos(x)),$$

and then differentiating both sides with respect to x gives us

$$\frac{y'}{y} = (\cos(x)) \ln(1 + \cos(x)) + (1 + \sin(x)) \cdot \frac{1}{1 + \cos(x)} \cdot (-\sin(x)).$$

Substituting $x = \pi/2$ and $y = 1$ (remembering that $\cos(\pi/2) = 0$ and $\sin(\pi/2) = 1$) yields

$$\frac{y'}{1} = 0 \cdot \ln(1 + 0) + (1 + 1) \left(\frac{1}{1 + 0} \right) (-1) = 0 - 2 = -2.$$

A graph of the function with a tangent line at the desired point is shown below.

