

MAT 125 Solutions to Midterm 2 (Fezzik)

1. For each of the functions $f(x)$ given below, find $f'(x)$.

4 points

(a) $f(x) = \frac{1 + 2x^2}{1 + x^4}$

Solution: This is a straightforward quotient rule problem:

$$f'(x) = \frac{(4x)(1 + x^4) - (1 + 2x^2)(4x^3)}{(1 + x^4)^2} = \frac{4x - 4x^3 - 4x^5}{(1 + x^4)^2}$$

The simplification is not required.

4 points

(b) $f(x) = \sin(2x) \tan(x)$

Solution: Apply the product rule, with a chain rule for the $\sin(2x)$ term to get

$$f'(x) = 2 \cos(2x) \tan(x) + \sin(2x) \sec^2(x).$$

4 points

(c) $f(x) = \arctan(\sqrt{1 + 3x})$

Solution: Applying the chain rule, we get

$$\frac{1}{1 + (\sqrt{1 + 3x})^2} \cdot \frac{1}{2} (1 + 3x)^{-1/2} \cdot (3) = \frac{3}{2(2 + 3x)\sqrt{1 + 3x}}$$

2. Compute each of the following derivatives as indicated:

4 points

(a) $\frac{d}{du} \left[\frac{u^3}{4} + \frac{4}{u^3} \right]$

Solution: Write this as $\frac{1}{4}u^3 + 4u^{-3}$ and apply the power rule to get

$$\frac{3}{4}u^2 - 12u^{-4}.$$

4 points

(b) $\frac{d}{dx} [e^x - \pi^3]$

Solution: Remember that π^3 is a constant and so its derivative is zero. Thus, we have $\frac{d}{dx} [e^x - \pi^3] = e^x$.

4 points

(c) $\frac{d}{dw} \left[\sqrt{1 + \sqrt{1 + w}} \right]$

Solution: View this as $\frac{d}{dw} \left[\left(1 + (1 + w)^{1/2} \right)^{1/2} \right]$ and apply the chain rule:

$$\frac{1}{2} \left(1 + (1 + w)^{1/2} \right)^{-\frac{1}{2}} \cdot \frac{1}{2} (1 + w)^{-\frac{1}{2}} = \frac{1}{4\sqrt{1 + w} \sqrt{1 + \sqrt{1 + w}}}$$


12 points

3. The set of points (x, y) which satisfy the relationship

$$y^2(y^2 - 9) = x^2(x^2 - 10)$$

lie on what is known as a “devil’s curve”, shown at right.

Write the equation of the line tangent to the given devil’s curve at the point $(\sqrt{10}, 3)$.



Solution:

First, we use implicit differentiation to determine the slope of the tangent line. This will be slightly easier if we rewrite the equation as $y^4 - 9y^2 = x^4 - 10x^2$ first. Differentiating with respect to x gives

$$4y^3y' - 9 \cdot 2y \cdot y' = 4x^3 - 10 \cdot 2x \quad \text{and so} \quad y' = \frac{x(2x^2 - 10)}{y(2y^2 - 9)}.$$

At our desired point, $x = \sqrt{10}$ and $y = 3$, and so the slope is

$$y' = \frac{\sqrt{10} \cdot 10}{3 \cdot 9} = \frac{10\sqrt{10}}{27}.$$

This means the desired line is

$$y - 3 = \frac{10\sqrt{10}}{27}(x - \sqrt{10}).$$

4. Let $f(x) = x \ln(2x)$

4 points

(a) Calculate $f'(x)$

Solution: Applying the product rule (and the chain rule) gives

$$f'(x) = \ln(2x) + x \frac{1}{2x} \cdot 2 = \ln(2x) + 1.$$

4 points

(b) Calculate $f''(x)$

Solution: Taking the derivative of the above, we get $f''(x) = \frac{1}{x}$.

3 points

(c) For what values of x is $f(x)$ increasing?

Solution: As we all know, $f(x)$ is increasing when $f'(x) > 0$. Thus, using our answer from part (a) tells us that we need to know when

$$\ln(2x) + 1 > 0 \quad \text{or, equivalently,} \quad \ln(2x) > -1.$$

Exponentiating both sides gives $2x > e^{-1}$, so we know that

$$f(x) \text{ is increasing for } x > \frac{1}{2e}.$$

3 points

(d) For what values of x is $f(x)$ concave down?

Solution: We need to determine when $f''(x) < 0$. From part (b), this means

$$\frac{1}{x} < 0 \quad \text{that is,} \quad x < 0.$$

However, remember that $\ln(3x)$ is only defined for $x > 0$. Thus $f(x)$ is concave up for all values of x in its domain. There are no values of x where $f(x)$ is concave down.

12 points

5. Give the x and y coordinates of the (absolute) maximum and minimum values of the function

$$y = x^4 - 8x^2 - 1 \quad \text{where} \quad -3 \leq x \leq 1.$$

Solution: First, we locate the critical points. Since the function is a polynomial, $f'(x)$ is defined everywhere, so we only need concern ourselves with the x for which $f'(x) = 0$.

Since $f'(x) = 4x^3 - 16x = 4x(x^2 - 4) = 4x(x - 2)(x + 2)$, we have the critical points

$$x = 0 \quad x = 2 \quad x = -2$$

However, since we are concerned only with $-3 \leq x \leq 1$, we discard $x = 2$.

Now we evaluate f at each of the critical points, and the endpoints:

- $f(0) = -1$.
- $f(-2) = 16 - 32 - 1 = -17$.
- $f(-3) = 81 - 72 - 1 = 8$.
- $f(1) = 1 - 8 - 1 = -8$.

The largest value of the above occurs at $x = -3, y = 8$. This is our absolute maximum.

The smallest occurs when $x = -2$ and $y = -17$, which is our absolute minimum.

12 points

6. Calvin's family is visiting a winery in Cutchogue, and he wanders off into the fermenting room and dives into one of the large cylindrical[†] wine vats. The vat has a diameter of 6 feet and is 8 feet tall. The vintner hears the splash and quickly opens the taps to drain the vat, which drains at a rate of 5 cubic feet per minute. How quickly is the height of wine in the tank dropping when the wine is 6 feet deep?

[†]The volume of a cylinder of height h and radius r is $\pi r^2 h$ and its surface area (excluding top and bottom) is $2\pi r h$. The density of the wine is about .98 kg/L or 61 pounds per cubic foot. 5 cubic feet is about 38 gallons or 142 liters. The wine is a rather sweet Riesling, but is probably less sweet after Calvin has been in it.



Solution: We have the formula for the volume of a cylinder $V = \pi r^2 h$. In our case, $r = 3$ since the diameter is 6, so we have $V = 9\pi h$. We want to know dh/dt .

Since the vat is draining at a rate of 5 cubic feet per minute, we have $dV/dt = -5$.


Using implicit differentiation, we get $\frac{dV}{dt} = 9\pi \frac{dh}{dt}$. So, we see that

$$\frac{-5}{9\pi} = \frac{dh}{dt}.$$


12 points

7. For each of the 4 functions graphed in the left column, find the corresponding derivative function among any of the 8 choices on the right (not just on the same row) and put its letter in the corresponding box. If the graph does not occur, use the letter X.


F




A:




B:




H




C:




D:




G




E:




F:



E



G:



H:

