

1. For each of the following sequences, determine whether it converges or diverges. If the sequence converges, give its limit. Justify your answer.

5 pts

$$(a) \left\{ \frac{n^4 + \cos(n\pi)}{(1+3n)(2+n^3)} \right\}_{n=0}^{\infty} = -\frac{1}{2}, \frac{1}{6}, \frac{3}{70}, \dots$$

Solution: For n large, the $\cos(n\pi)$ term is irrelevant, as are the lower powers of n in the denominator. Thus

$$\lim_{n \rightarrow \infty} \frac{n^4 + \cos(n\pi)}{(1+3n)(2+n^3)} = \lim_{n \rightarrow \infty} \frac{n^4}{3n^4} = \frac{1}{3}$$

5 pts

$$(b) \left\{ \frac{\ln(2n)}{n} \right\}_{n=1}^{\infty} = \ln 2, \ln 2, \frac{\ln 6}{3}, \dots$$

Solution: $\lim_{n \rightarrow \infty} \frac{\ln(2n)}{n}$ is of the form ∞/∞ , so we can use L'Hôpital's rule. Thus,

$$\lim_{n \rightarrow \infty} \frac{\ln(2n)}{n} = \lim_{n \rightarrow \infty} \frac{1/n}{2} = 0$$

5 pts

$$(c) \{\pi + (-1)^n\}_{n=0}^{\infty} = \pi + 1, \pi - 1, \pi + 1, \dots$$

Solution: The sequence diverges, because the limit as $n \rightarrow \infty$ does not exist; it alternates between $\pi + 1$ and $\pi - 1$.

10 pts

2. What value of N do we need to ensure that the sum $\sum_{n=0}^N \frac{(-1)^n}{n^2 + 4}$ is within $1/100$ of the limit? Keep in mind that N must be a whole number.

Solution: Since this is an alternating series, we know that the remainder R_N is less than the absolute value of the $N + 1$ -st term. This means we want N so that

$$\frac{1}{(N+1)^2 + 4} < \frac{1}{100} \quad \text{or, equivalently} \quad 100 < (N+1)^2 + 4$$

This holds for $N = 9$.


- 15 pts 3. Find the volume of the solid whose base is the half-circle

$$x^2 + y^2 = 4 \quad \text{with } y \geq 0$$

and whose cross-sections perpendicular to the y -axis are squares.


Solution: Note that since the cross-sections perpendicular to the y -axis are squares, we want to integrate dy . (If we tried to integrate dx , we would have very complicated slices.)

So, we write the base as $x = \pm\sqrt{4 - y^2}$, and observe that if we take a slice at a particular y value, the square cross section will stretch from $x = -\sqrt{4 - y^2}$ to $x = +\sqrt{4 - y^2}$. This means the side length of the square is $2\sqrt{4 - y^2}$, and its area is $4(4 - y^2)$.



The volume of the solid is then given by integrating the cross-sectional area as y ranges from 0 to 2.

$$Vol = \int_0^2 4(4 - y^2) dy = 4(4y - y^3/3) \Big|_0^2 = 4(8 - \frac{8}{3}) = \frac{64}{3}.$$



- 15 pts 4. Find the area that lies inside the polar curve $r = \sqrt{1 - \sin \theta}$

but outside the circle $r = -\sqrt{2} \sin \theta$.


Solution: It is easiest to do this by calculating the outer area first, and then subtract off the inner area. The two curves meet at the origin and along the negative y -axis.

The area inside the cardioid-like shape is

$$\begin{aligned} \frac{1}{2} \int_0^{2\pi} (\sqrt{1 - \sin \theta})^2 d\theta &= \frac{1}{2} \int_0^{2\pi} 1 - \sin \theta d\theta \\ &= \frac{1}{2} (\theta + \cos \theta) \Big|_0^{2\pi} = \frac{1}{2} ((2\pi + 1) - 1) \\ &= \pi. \end{aligned}$$

The circle has radius $\sqrt{2}/2$, so its area is $\pi/2$.

This means the shaded area is $\pi - \pi/2 = \frac{\pi}{2}$.



15 pts

5. Find the volume of the solid obtained by rotating the region between the two curves

$$y = 2x \quad \text{and} \quad y = x^2$$

about the y -axis.

- (a) Write an integral which represents the volume.


Solution: Note that the curves cross at $(0, 0)$ and $(2, 4)$.

We can integrate dy or dx , as we wish. If we integrate with respect to x , we have cylindrical shells. The height of each cylinder runs from $y = x^2$ to $y = 2x$, so it is $2x - x^2$. The radius of the cylinder is x . This means the volume is given by

$$\int_0^2 2\pi x(2x - x^2) dx = 2\pi \int_0^2 2x^2 - x^3 dx.$$

If instead we want to integrate dy , note that such a slice will be a “washer” with inner radius $x = y/2$ and outer radius $x = \sqrt{y}$. This means we have the integral

$$\int_0^4 \pi(\sqrt{y})^2 - \pi(y/2)^2 dy = \pi \int_0^4 y - \frac{y^2}{4} dy$$



- (b) Evaluate the integral in (a).

Solution: For the first integral, we have

$$2\pi \int_0^2 2x^2 - x^3 dx = 2\pi \left(\frac{2}{3}x^3 - \frac{1}{4}x^4 \right) \Big|_0^2 = 2\pi \left(\frac{16}{3} - \frac{16}{4} \right) = \frac{8\pi}{3}.$$

The second integral gives

$$\pi \int_0^4 y - \frac{y^2}{4} dy = \pi \left(\frac{y^2}{2} - \frac{y^3}{12} \right) \Big|_0^4 = \pi \left(\frac{16}{2} - \frac{64}{12} \right) = \frac{8\pi}{3}.$$

Of course, they both evaluate to the same value.

15 pts

6. Determine the interval of convergence for the power series
- $\sum_{n=1}^{\infty} \frac{(-1)^n(4x-1)^n}{\sqrt{n+5}}$
- . Don't forget to establish convergence or divergence at the endpoints!

Solution: First, we apply the ratio test, calculating $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$:

$$\lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1}(4x-1)^{n+1}}{\sqrt{n+1+5}} \cdot \frac{\sqrt{n+5}}{(-1)^n(4x-1)^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{\sqrt{n+5}}{\sqrt{n+6}} \cdot (4x-1) \right| = |4x-1|$$

Solution: (continued) So, for the series to converge, we must have $|4x - 1| < 1$, that is, $-1 < 4x - 1 < 1$. Adding 1 to both sides yields $0 < 4x < 2$, or equivalently $0 < x < \frac{1}{2}$. (If you prefer, observe that the center is $\frac{1}{4}$ and the radius of convergence is also $\frac{1}{4}$.)

Now we need to establish what happens at the endpoints.

When $x = 0$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n(-1)^n}{\sqrt{n+5}}$, or, equivalently, $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n+5}}$. This series diverges. If you like, you can use the integral test, or we can use limit comparison with a p -series with $p = 1/2$:

$$\lim_{n \rightarrow \infty} \frac{1/\sqrt{n+5}}{1/\sqrt{n}} = 1,$$

so the two series do the same thing. Since $\sum 1/\sqrt{n}$ diverges, so does the original.

When $x = 1/2$, the series becomes $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n+5}}$. This is an alternating series. Since the terms are decreasing and $\lim 1/\sqrt{n+5} = 0$, the series converges.

Consequently, the interval of convergence is $\left(0, \frac{1}{2}\right]$.

7. For each of the infinite sums below, state whether it converges or diverges. Justify your answer completely.

5 pts

(a) $\sum_{n=1}^{\infty} \frac{\ln(2n)}{n} = \ln 2 + \ln 2 + \frac{\ln 6}{3} + \dots$

Solution: Diverges by the integral test, or by comparison with $\sum \frac{1}{n}$ (The series is larger than a divergent series, and so diverges).

5 pts

(b) $\sum_{n=3}^{\infty} \frac{n^2 + 5}{(n^2 - 1)(n^2 - 4)} = \frac{7}{20} + \frac{7}{60} + \frac{1}{16} + \dots$

Solution: Since $\frac{n^2 + 5}{(n^2 - 1)(n^2 - 4)} < \frac{n^2}{n^4} = \frac{1}{n^2}$, the series converges by comparison with a convergent p -series ($p = 2$).

5 pts

(c) $\sum_{n=5}^{\infty} \frac{5^n - 3^n}{7^{n+2}} = \frac{2882}{82543} + \frac{304}{117649} + \frac{75938}{40353607} + \dots$

Solution: Observe that $\sum \frac{5^n - 3^n}{7^{n+2}} = \frac{1}{49} \left(\sum \frac{5^n}{7^n} - \sum \frac{3^n}{7^n} \right)$. So this is the difference of two convergent geometric series (the ratios are both less than one), and it converges (to $\frac{1}{49} \left(\frac{7}{2} - \frac{7}{4} \right) = 1/28$).

5 pts

(d) $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}n}{\sqrt{n^2 + 4}} = \frac{1}{\sqrt{5}} - \frac{1}{\sqrt{2}} + \frac{3}{\sqrt{13}} - \dots$

Solution: This diverges, since the limit of a_n is not zero.

15 pts

8. Jack has a goose that lays golden eggs, one each day. Unfortunately each egg is only $\frac{4}{5}$ the mass of the previous one. Jack needs to obtain 120 grams of gold to ransom his sister from the evil monkey-king. If the first egg weighed 20 grams, does Jack ever get enough gold? If so, how long must he wait? Justify your answer.

Solution: The total amount of gold (in grams) that Jack will get is

$$20 + 20 \cdot \frac{4}{5} + 20 \cdot \frac{4^2}{5^2} + 20 \cdot \frac{4^3}{5^3} + \dots = 20 \sum_{n=0}^{\infty} \frac{4^n}{5^n}$$

This is a geometric series, and the sum is

$$\frac{20}{1 - 4/5} = \frac{20}{1/5} = 100$$

Sadly, no matter how long he waits, he will never even get 100 grams of gold.

15 pts

9. Recall that Hooke's law says that the amount of force required to stretch a spring x units beyond its natural length is kx , where k is a constant depending on the spring.

A giant spring designed to hold the gates of Mordor closed has a natural length of 10 meters. If it takes 1800 Joules¹ to stretch the spring from 10 meters to a length of 13 meters, how much work will it take to stretch the spring from 13 meters to a length of 15 meters?

Solution: First, we need to determine the spring constant. Since the amount of work to stretch the spring from 10 to 13 is 1800 J, we have

$$1800 = \int_0^3 kx \, dx = \frac{kx^2}{2} \Big|_0^3 = \frac{9k}{2},$$

and so $k = 1800 \cdot 2/9 = 400$. (We integrated from 0 to 3 because the natural length of the spring is 10 meters).

Now the required work to stretch the spring from 13 to 15 is given by

$$\int_3^5 400x \, dx = \frac{400x^2}{2} \Big|_3^5 = 200(25 - 9) = 200 \cdot 16 = 3200 \, J$$

¹A Joule is the amount of work needed to apply a force of 1 Newton over the distance of 1 meter.