

13. (*expires 3/1*) Similar to the class discussion on [Feb. 20](#), the worksheet [sliderfit.mw](#) contains an interactive slider to demonstrate how the interpolating polynomial changes when the  $y$ -value of one of the data points is moved.

Modify this worksheet to add another slider which allows the user to also move the  $x$ -value of the same point.

Also modify it to include the graph of the corresponding cubic spline in the plot.

Hint: you will need to modify the “startup code” at the top, and right-clicking on the slider to select “**Edit Value Changed Code...**” will be useful.

14. (*expires 3/1*) Fit the points  $(-1.9, -4.7), (-0.8, 1.2), (0.1, 2.8), (1.4, -1.2), (1.8, -3.5)$  with a quadratic function  $f(x) = ax^2 + bx + c$ , using the least square method.

You can load these data points from the web via the link at [fitquad.txt](#) which defines a list `fitquad` containing them.

15. (*expires 3/1*) The file [fitexp.txt](#) defines a list `expdata` with 21 data points approximating an exponential curve of the form  $y = ae^{bx}$ .

Find  $a$  and  $b$  by taking an appropriate logarithm, then use `CurveFitting[LeastSquares]` to find the resulting “best” line. Then transform this line appropriately to get an exponential curve which approximates the given data. The `map` command might be helpful.

Plot the exponential and the points from `expdata` on the same axes, and write the approximating exponential in the form  $y = ae^{bx}$ .

16. (*expires 3/1*) Fit the set of points  $(1.021, -4.30), (1.001, -2.12), (0.99, 0.52), (1.03, 2.51), (1.00, 3.34), (1.02, 5.30)$  with a line, using the least square method. Plotting these points and the line on the same graph shows that this is not a good fit. (This is most apparent if you have a plot with, say,  $0 < x < 2$  and  $-5 < y < 6$ .) Think of a better way to find a line which *is* a good fit and use Maple to do it. Explain in your solution why you think your better way is indeed an improvement. In case you don’t want to retype them, the file [badfit.txt](#) defines a list `fitme` containing these points.

17. (*expires 3/1*) In this problem we will estimate the charge of the [electron](#).

If an electron of energy  $E$  is thrown into a magnetic field  $B$  which is perpendicular to its velocity, the electron will be deflected into a circular trajectory of radius  $r$ . The relation between these three quantities is:

$$B r e = \frac{E^2}{m^2 c^4} \sqrt{E^2 - m^2 c^4}, \quad (1)$$

where  $e$  and  $m$  are, respectively, the charge and the mass of the electron, and  $c$  is the [speed of light](#) ( $2.9979 \times 10^8 \frac{\text{meter}}{\text{sec}}$ ). The rest mass of the electron is defined by  $E_0 = mc^2$ , and is about equal to  $8.817 \times 10^{-14}$  [Joules](#). In our experimental set-up the energy of the emitted electrons is set to be  $E = 2.511E_0$ .

The file [electron.txt](#) defines a list called `electron`. Each element of the list is a pair of the form  $[B_i, r_i]$ ; these quantities are expressed in [Teslas](#) and meters. Use least square fitting to determine the best value for  $e$ .

*Hint: Notice that the right hand side of eqn (1) is just a constant—calculate it once and for all and give it a name. Then eqn (1) becomes an equation which is linear in the unknown  $e$ . To verify your solution:  $e \approx 1.602 \times 10^{-19}$  [Coulomb](#). Physical constants courtesy of [N.I.S.T](#).*