


Volumes of simple bodies ← 3D domain, enclosed inside a surface

Consider a body which is analogous to a domain of type I in the plane



$$z = g(x, y)$$


$$z = f(x, y)$$

Body B is between the two surfaces

$$B: (x, y) \in D$$

$$f(x, y) \leq z \leq g(x, y)$$

Our goal is to compute the volume of this body.



divide body up
into thin vertical
columns.

The volume of one of these columns is approx

(height of column) $\cdot$ (Area of Base)

$$= (g(x, y) - f(x, y)) \, dA.$$


Summing up the little columns and taking
the area of the base to zero, we have

$$V(B) = \iint_D (g(x, y) - f(x, y)) \, dA$$


If we are lucky, this reduces to iterated integrals. (2)

Ex: Volume of tetrahedron

$$B: \begin{aligned} x &\geq 0, y \geq 0, z \geq 0 \\ x + y + z &\leq 1 \end{aligned}$$



$$D: \begin{aligned} x &\geq 0, y \geq 0 \\ x + y &\leq 1 \end{aligned}$$



$$h(x, y) = 1 - x - y$$

$$g(x, y) = 0$$

$$V(B) = \iint_D (1 - x - y) dA$$

$$= \int_0^1 \left(\int_0^{1-x} (1 - x - y) dy \right) dx$$

$$= \int_0^1 \left[(1-x)^2 - \frac{1}{2} (1-x)^2 \right] dx = \frac{1}{2} \int_0^1 (1-x)^2 dx$$

$$= \frac{1}{2} \int_0^1 x^2 dx = \frac{1}{2} \cdot \frac{1}{3} = \frac{1}{6}.$$


Volume of unit Ball

$$B = x^2 + y^2 + z^2 \leq 1$$

Base (projection on x - y plane)

$$D = x^2 + y^2 \leq 1$$

= unit disk



$$z^2 + x^2 + y^2 = 1$$

$$\Rightarrow z = \pm \sqrt{1 - x^2 - y^2}$$


$$B: x^2 + y^2 \leq 1$$

$$-\sqrt{1 - x^2 - y^2} \leq z \leq \sqrt{1 - x^2 - y^2}$$

$f(x, y)$ $g(x, y)$

$$V(B) = \iint_D 2\sqrt{1 - x^2 - y^2} \, dA$$

$$= \int_{-1}^1 dx \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} 2\sqrt{1 - x^2 - y^2} \, dy$$




$$I = \int_{-a}^a 2\sqrt{a^2 - y^2} dy$$

= Area of disk
of radius a

$$= \pi a^2 = \pi(1-x^2)$$

$a = \sqrt{1-x^2}$
length of segment.
 $2\sqrt{a^2 - y^2}$



Thus

$$V(B) = \pi \int_{-1}^1 (1-x^2) dx = \pi \left(2 - \frac{2}{3}\right) = \frac{4\pi}{3}.$$

Volume of unit ball (radius 1) is $\frac{4\pi}{3}$.

Archimedes did this 2.5 millennia ago.

He essentially did the same computation without machinery of calculus.

Lamina : what is it?




D plane, solid plate D
with variable density
 $f(x,y)$ (thickness of material) $\left[\frac{\text{kg}}{\text{m}^2} \right]$

example of lamina is a door.

Mass of lamina

$$M = \iint_D p(x,y) dA$$

Moments w.r.t. x-axis




Imagine x-axis is a solid rod. Balance D on it. Because of gravity there is a force acting on it.

$$M_x = \iint_D y \rho(x,y) dA$$

← moment is product of the force and distance to axis of rotation

Similarly moment w.r.t. y-axis

$$M_y = \iint_D x \rho(x,y) dA$$



Center of mass: $(\bar{x}, \bar{y})$

If you put it on the x-axis then the total moment is zero

$$\bar{x} = \frac{M_x}{M} \quad \bar{y} = \frac{M_y}{M}$$




you can balance the lamina with a support at center of mass

$\curvearrowright$ at equilibrium

Class of problems: given a domain D and density ρ , find all these things.

Example: $\rho(x, y) = x^2$



$$M = \iint_D x^2 dA = \int_0^1 dx \left(\int_{-1-x}^{1-x} x^2 dy \right)$$

$$= \int_0^1 dx \left(x^2 (2 - 2x) \right)$$

$$= \int_0^1 (2x^2 - 2x^3) dx$$

$$= \frac{2}{3} - \frac{2}{4} = \frac{1}{6}$$

Now we find moments

$$M_x = \iint_D y \rho(x, y) dA$$

$$= \int_0^1 dx \left(\int_{-1+x}^{1-x} y x^2 dy \right) = 0$$

y is odd function
integral is symmetric

$$M_y = \iint_D x \rho(x, y) dA$$

$$= \int_0^1 dx \left(\int_{-1+x}^{1-x} x^3 dy \right)$$



$$= \int_0^1 x^3 (2 - 2x) dx = \int_0^1 (2x^3 - 2x^4) dx$$

$$= \frac{1}{2} - \frac{2}{5} = \frac{1}{10}$$

Center of Mass:

$$(\bar{x}, \bar{y}) = \left(\frac{\frac{1}{10}}{\frac{1}{6}}, 0 \right) = (0.6, 0)$$

Moment of Inertia



angular velocity ω ↓

$$\text{Kinetic Energy} = \frac{1}{2} I \omega^2$$

moment of inertia depends on axis of rotation

(speed)² for rotation

$$I_x = \iint_D \underbrace{y^2 \rho(x, y)}_{\text{kinetic energy}} dA$$

$$I_y = \iint_D x^2 \rho(x, y) dA$$

Example:

$$I_x = \iint_D y^2 x^1 dA = \int_0^1 dx \int_{-1+x}^{1-x} y^2 x^2 dy$$

$$= \int_0^1 dx \left(x^2 \cdot \frac{2}{3} (1-x)^3 \right) = \int_0^1 dx \frac{2}{3} (x^2 - 3x^3 + 3x^4 - x^5)$$


$$= \frac{2}{3} \left(\frac{1}{3} - \frac{3}{4} + \frac{3}{5} - \frac{1}{6} \right) = \frac{1}{90}$$

$$I_y = \iint_D x^2 x^2 dA$$

$$= \int_0^1 dx \int_{-1+x}^{1-x} x^4 dy = \int_0^1 dx (2-2x) x^4 dx$$

$$= \frac{2}{5} - \frac{2}{6} = \frac{2}{30} = \frac{1}{15}$$

$$I_z = I_x + I_y$$




rotating around z-axis
kinetic energy of
each particle is

$$dm \omega |\vec{r}|^2 \\ = dm \omega^2 (x^2 + y^2)$$

$$I_z = \iint_D (x^2 + y^2) dA = I_x + I_y.$$

In our example $I_z = \frac{1}{90} + \frac{1}{15} = \frac{7}{90}$

Double integral in polar coordinates



polar coordinates of (x, y)
(length of $\vec{r}$, angle θ)

$$\begin{aligned} r &= \|\vec{r}\| \\ x &= r \cos \theta \\ y &= r \sin \theta \end{aligned}$$



$$r = \sqrt{x^2 + y^2}$$

$$\theta = \arctan\left(\frac{y}{x}\right)$$

works in the
right half plane

modify to $\arccot\left(\frac{x}{y}\right)$ in
upper half plane etc.



In cartesian coordinates, simplest domains
are rectangles.

In polar coordinates simplest domains
are $a \leq r \leq b$ $\alpha \leq \theta \leq \beta$



Consider $\iint_D f(r, \theta) dA$ -

What is dA ?




$$dA = r dr d\theta.$$

Thus

$$\iint_D f(r, \theta) dA = \int_a^b \int_a^b f(r, \theta) r dr d\theta.$$

Example




$$D: \quad 0 \leq r \leq 1 \\ -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

$$\iint_D x \, dA = \iint_D r \cos \theta \, r \, dr \, d\theta$$

$$= \int_{-\pi/2}^{\pi/2} \left(\int_0^1 r \cos \theta \, r \, dr \right) d\theta = \int_{-\pi/2}^{\pi/2} \cos \theta \int_0^1 r^2 \, dr \, d\theta$$

$$= \frac{1}{3} \int_{-\pi/2}^{\pi/2} \cos \theta \, d\theta = \frac{2}{3}.$$



$$D = \begin{aligned} &a \leq r \leq b \\ &-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2} \end{aligned}$$

$$\rho(r, \theta) = 1$$

Center of mass of horse shoe $(\bar{x}, \bar{y})$

$$M = \iint_D \rho \, dA = \int_{-\pi/2}^{\pi/2} \int_a^b r \, dr \, d\theta$$

$$= \pi \int_a^b r \, dr = \frac{\pi}{2} (b^2 - a^2)$$

$$M_x = \iint_D \rho y \, dA = \int_{-\pi/2}^{\pi/2} \int_a^b r \sin \theta \, r \, dr \, d\theta$$

$$= \int_{-\pi/2}^{\pi/2} \sin \theta \int_a^b r^2 \, dr \, d\theta = \int_{-\pi/2}^{\pi/2} \sin \theta \left(\frac{b^3 - a^3}{3} \right) d\theta$$

$$= \frac{b^3 - a^3}{3} \int_{-\pi/2}^{\pi/2} \sin \theta \, d\theta = 0$$

$$M_y = \iint_D \rho x \, dA = \int_{-\pi/2}^{\pi/2} \int_a^b r \cos \theta \, r \, dr \, d\theta$$

$$= \int_{-\pi/2}^{\pi/2} \cos \theta \left(\frac{b^3 - a^3}{3} \right) d\theta$$


$$= \frac{2}{3} (b^3 - a^3)$$

$$\bar{x} = \frac{M_y}{M} = \frac{4}{3} \frac{b^3 - a^3}{b^2 - a^2}$$

$$\bar{y} = 0$$

Volume of ball

$$B: x^2 + y^2 + z^2 \leq 1$$



$$D: x^2 + y^2 \leq 1 \quad \text{in polar}$$

$$D: \begin{aligned} 0 \leq r \leq 1 \\ 0 \leq \theta \leq 2\pi \end{aligned}$$

$$z = \pm \sqrt{1 - x^2 - y^2}$$

$$V(B) = \iint_D 2\sqrt{1 - x^2 - y^2} \, dA$$

$$= \int_0^{2\pi} \int_0^1 2\sqrt{1 - r^2} \, r \, dr \, d\theta$$

$$= 4\pi \int_0^1 \sqrt{1 - r^2} \, r \, dr$$

$$= 2\pi \int_0^1 \sqrt{1 - s} \, ds = 2\pi \int_0^1 \sqrt{s} \, ds \quad \begin{aligned} s &= r^2 \\ ds &= 2r \, dr \end{aligned}$$

$$= -2\pi \left. \frac{2}{3} (1 - s)^{3/2} \right|_0^1 = \frac{4\pi}{3}$$

$$V(\text{Ball of radius } R) = \frac{4\pi}{3} R^3$$